REVIEW

The role of thyroid hormones in depression

Carsten Kirkegaard and Jens Faber

Department of Internal Medicine E, Hillerød Sygehus, and Department of Endocrinology E, Frederiksberg Hospital, Denmark

(Correspondence should be addressed to C Kirkegaard, Department of Internal Medicine E, Hillerød Sygehus, DK-3400 Hillerød, Denmark)

Introduction

During the last 30 years a huge number of scientific articles have appeared on the subject of relationships between psychiatric disease and thyroid hormones. These studies have demonstrated the presence of numerous changes in the hypothalamo–pituitary–thyroid (HPT) axis, mainly in patients with depression, but also in patients with other psychiatric diseases. Simultaneously, many studies have been published on the possible therapeutic effects in depression of the hormones involved in the HPT axis.

Despite great efforts to standardize the classification of depression, this is still a less well-defined disease, possibly including several subtypes with different pathogenesis and biochemical abnormalities. Furthermore the classification has changed over time, making it difficult to interpret previously published data. The most accepted classifications are the DSM-IV and ICD-10. These cover a spectrum from minor depression (neurotic depression) through major depression to melancholic (endogenous and psychotic) depression. Furthermore major and melancholic depressions are divided into uni- and bipolar depressions, the latter also demonstrating episodes of mania. Rapid cycling bipolar psychosis is a subgroup of the bipolar depression, demonstrating four or more episodes in any year of observation.

The present review focuses on the concentrations of hormones of the HPT-axis seen in uni- and bipolar depression, their relation to relapse after antidepressive treatment, and a survey of therapeutic trials with thyrotropin-releasing hormone (TRH), thyroid-stimulating hormone (TSH) and thyroid hormones, as well as the effect of the various drugs used in depression on the HPT-axis. The hypothesis is put forward that the changes seen in the HPT-axis during depression might be explained by cerebral serotonin deficiency, and that tri-iodothyronine (T_3) treatment, to some degree, can revert this deficiency.

Iodothyronines in depression

An overall picture of thyroid-related disturbances seen in depression is given in Table 1.

Thyroxine (T_4)

Serum T_4 levels, both total and non-protein-bound (free), are consistently found as normal to increased in

groups of depressed patients (reviewed in 1, 2), approximately 25% of the patients having levels above the reference range. The different findings might be explained by the different severity of depression among the patients studied, since some studies have found a correlation between the severity of depression and serum T_4 levels (2). An additional explanation might be that the depressed patient is often in a state of semi-starvation, and thus may present changes in the HPT-axis similar to those seen in patients with nonthyroidal somatic illness, elevated serum T_4 and 3,3',5'-tri-iodothyronine (rT_3), and reduced serum T_3 .

Turnover studies using radiolabeled T_4 in a small group of depressed patients have demonstrated that also the daily production rate (PR) of T_4 is significantly increased, by 30% (3). This finding contrasts with the unaltered T_4 PR found in groups of patients with different nonthyroidal illnesses (4). These studies used a noncompartmental kinetic approach for the evaluation of PRs. This method correctly estimates the PR of T_4 . Increased PR of T_4 in depressed patients thus suggests that the thyroid gland is stimulated abnormally in the depressed patient.

The concentration of free T_4 in cerebrospinal fluid (CSF) seems relatively increased during depression, since recovery is followed by a reduction (5). The ratio between CSF and serum free T_4 levels has been found to be 0.6, both before and after recovery of depression due to electro-convulsive treatment (ECT) (5). This suggests that CSF T_4 concentrations follow systemic levels, and that transport of T_4 into CSF is restricted. An alternative explanation is that T_4 is taken up directly into brain tissues, and partly undergoes deiodination, thus yielding less T_4 into the CSF.

T_3

Serum T_3 levels in depressed patients are often found normal, but several studies have found reduced levels, typically in more severely depressed patients (1, 2). Serum T_3 levels are influenced by numerous factors which all may be present in the depressed patient: starvation, concomitant somatic illness and medication, and changes (increase) in cortisol levels. When present these factors all tend to decrease serum T_3 levels (4, 6). This makes interpretation of serum T_3 concentrations difficult. Serum free T_3 levels have been found both

Table 1 Changes in total and free serum concentrations, CSF concentrations, and production rates of various hormones in patients with depression. Relevant references are given in parentheses.

	Total serum	Free serum	CSF	Production rate
T ₄	Increased (1, 2)	Increased (1, 2)	Increased (5)	Increased (3)
T ₃	Unaltered (1, 2) Decreased (1, 2)	Unaltered (2, 9) Decreased (2)	—	Unaltered (3)
rT ₃	Increased (2, 11) Unaltered (5, 9)	Unaltered (2) Increased (5, 11)	Increased (5, 12)	—
TSH	Decreased, but elevated in comparison to T ₄ production (1, 3, 13–17). In TCA-resistance and RCBP, increased TSH possible (27–30)	—	—	—
Δmax TSH	Decreased (1, 2, 13) Increased (27–30)	—	—	—
TRH	—	—	Increased (20, 21) Unaltered (22)	—
Somatostatin	—	—	Decreased (24–26)	—

TCA = tricyclic antidepressants; RCBP = rapid cycling of bipolar psychosis.

normal and reduced (2). However, methods for the measurement of free T₃ in serum often give spurious, mainly reduced levels, in diseased patients (7). Ultra-filtration seems the method of choice (7, 8), and using this technique we have found unaltered free T₃ levels in depression (9). The daily PR of T₃ in unmedicated, moderately depressed patients has been studied using tracer turnover techniques, and T₃ PR was found normal (3). This was quite different from patients with various nonthyroidal illnesses, in whom a 40% reduction in T₃ PR has been demonstrated (4). The combination of an increased T₄ PR and an unaltered T₃ PR in depression suggests a reduced deiodination of T₄ into T₃, as also seen in nonthyroidal illness, but with more substrate (T₄) availability than in somatic nonthyroidal illnesses. However, correct interpretation of noncompartmental tracer studies is based on the assumption that the tracer injected is distributed freely into all tissue compartments (4). If anything, some degree of underestimation of T₃ PR might take place when performing T₃ kinetics in humans (4), but conflicting results are available on this issue (10).

The reduced conversion of T₄ into T₃ seen during depression (3) might be due to reduced deiodination enzyme activity. However, in which compartment of the human body this takes place is at present unknown. This could in theory be the brain, but unfortunately we are not aware of any data on intracerebral T₃ content or CSF levels of T₃ in depression.

rT₃

Changes in serum total as well as free rT₃ levels in depression seem to follow those seen for serum total and free T₄, respectively (2, 5, 11).

CSF levels of rT₃ have been studied in different types of depression, and have been found highest in the endogenous type (12). Similar to changes in serum, CSF

free rT₃ concentrations seem to follow CSF free T₄ levels (5), but the ratio of CSF to serum free rT₃ has been found to be approximately 26, which is quite different from that of T₄ (0.6) (5). This ratio for rT₃ did not change after recovery from the depression, and this suggests that intracerebral concentration of rT₃ is high in humans, and that rT₃ in brain is mainly derived from local production from T₄. This enzymatic production seems not changed in depression, and this argues against rT₃ as a pathogenic factor in depression.

TSH in depression

Previously, changes in circulating levels of TSH in depressed patients were evaluated using the TSH response to intravenous TRH, due to lack of sufficiently sensitive TSH assays. The response in TSH has usually been evaluated as the peak value minus basal TSH (ΔTSH). Normal ΔTSH to 200 μg TRH i.v. is typically 2–16 mU/l. Patients with depression, as a group, have reduced ΔTSH, and approximately 25% have ΔTSH below 2 mU/l in endogenous (melancholic) depression (1). Using newer, and more sensitive TSH methods, basal TSH values correlate closely to peak TSH as well as ΔTSH (13–15). According to this, depressed patients seem to have some degree of reduced basal serum TSH, but within the normal range (13, 15). Basal as well as ΔTSH using a sensitive TSH assay have been studied in various types of depression, and those patients with endogenous depression had the lowest levels of both basal TSH and ΔTSH (13), ΔTSH being a mean of 3.6 mU/l.

The diurnal variation of serum TSH is, in healthy man, associated with a surge in TSH around midnight. In untreated depression this diurnal variation seems attenuated (16). After complete recovery from depression the diurnal TSH variation seemed to be re-established, but

no or incomplete recovery still harbored an attenuated diurnal variation (17).

In one study the interrelation between serum TSH concentrations and the daily production of T_4 and T_3 in endogenous depression was evaluated (3). Six depressed patients were compared with seven subjects with L - T_4 -treated hypothyroidism. These two groups had quite similar T_4 and T_3 PRs, both having an equal T_4 PR/ T_3 PR ratio. Serum TSH level using a sensitive assay was a median 0.11 mU/l among L - T_4 -treated hypothyroid subjects, but 0.90 mU/l in depressed patients, significantly higher than in the L - T_4 -treated hypothyroid group (3). This suggests an inappropriate secretion of TSH (in relation to the elevated thyroidal production of T_4) in patients with endogenous depression, compatible with some degree of central over-stimulation of the thyroid in the untreated depressed state.

The apparent paradox that serum TSH levels are found slightly decreased among depressed patients, and our statement of inappropriately elevated serum TSH levels in comparison to daily production of T_4 and T_3 (3), can be compared with the situation of chronic TRH stimulation (by oral administration) in man. In this experimental setting, TRH induced an initial increase in serum TSH, whereas later, at the time when serum T_4 and T_3 began to increase, serum TSH levels were normalized and were unable to respond to repeated TRH stimulation (18, 19). Thus chronic stimulation with TRH in man can provoke a pattern in serum levels of TSH, T_4 and T_3 similar to those seen in the depressed patient.

Chronic stimulation of the thyrotrophs in the pituitary during depression might be caused by TRH, since TRH levels have been found elevated in CSF in two studies (in total 31 patients) (20, 21), although normal levels were found in a third study including 17 patients (22). Serum TSH levels are also influenced by somatostatin, which inhibits the TSH release from the pituitary (23). Three studies have found that the CSF concentration of somatostatin is reduced during depression (24–26). This might contribute to an increase in serum TSH levels.

Two subgroups of the depression syndrome seem to present with a different pattern in the HPT-axis. One group consists of patients with bipolar depression (27, 28), and another group of patients is characterized by a depression resistant to treatment with tricyclic antidepressants (TCA) (29, 30). Both groups have a tendency towards slightly elevated basal serum TSH levels (approximately 20% have levels above the upper normal reference range), or exaggerated TSH response to TRH stimulation. These changes seem independent of previous or ongoing lithium treatment, and probably reflect some degree of thyroid insufficiency. Immunological mechanisms seem to play a role, since patients with bipolar depression have increased frequency of sera positive for antithyroid peroxidase (anti-TPO) antibodies (28). Whether this is an epiphenomenon to bipolar

depression, or might be considered as one of several pathogenic factors, is at present unknown. Interestingly, postpartum changes in mood seem to be associated with the presence of anti-TPO antibodies and thyroid dysfunction (31). Thyroid hormone substitution might therefore be of potential benefit in alleviating depressive symptoms in the puerperal period. A preliminary study, in which L - T_4 was administered immediately after delivery, in fact resulted in reduced depression score among the L - T_4 -treated women compared with placebo treatment (32).

Predictive value of changes of Δ TSH

Another interesting aspect of the TSH response to TRH is the possible predictive value of the changes before and after antidepressive treatment. It is a typical clinical finding that depressed patients, after recovery following ECT, demonstrate a high frequency of early relapse, typically within 6 months, if not treated prophylactically with antidepressive medication. We have repeatedly demonstrated that those patients with early relapse of endogenous depression have not normalized their reduced TSH response to TRH, despite clinical recovery. This has been demonstrated after treatment with ECT as well as sleep deprivation (i.e. nonpharmacological treatment modalities) (1, 33, 34). More than 80% of those patients with early relapse did not change their maximal TSH response to TRH (Δ max TSH). An unchanged Δ max TSH was defined as post-treatment Δ max TSH minus pre-treatment Δ max TSH ($\Delta\Delta$ max TSH) less than 2 mU/l. By contrast more than 80% of those patients who remained clinically cured for more than 6 months had a $\Delta\Delta$ max TSH exceeding 2 mU/l (1). Serum T_4 as well as free T_4 levels remained elevated in those patients with early relapse and continued reduced Δ max TSH (9). Thus a continued disturbance of the HPT-axis despite clinical improvement of endogenous depression suggests that the patient is not clinically cured. However, other studies in patients treated with antidepressive medication, rather than nonpharmacological treatment, have been unable to confirm these results (2).

In a small group of patients treated with amitriptyline we confirmed this lack of predictive value of the TRH test (33). Thus it is possible that the use of antidepressive medication might interfere with the HPT-axis on multiple sites, thus blurring the finding seen in the unmedicated, clinically cured patient. Alternatively, in the studies using TCA, the TRH tests have been performed with intervals too long to maintain a stable Δ TSH in patients not cured of depression.

Relationships between depression and thyroid hormones – a hypothesis

The pathogenesis of endogenous depression is not known, and is most probably multifactorial. The

Figure 1 Decreased brain serotonin levels (serotonin (-)) activate (indicated by (+)) the HPT-axis, and increased T_3 (by endogenous as well as exogenous sources) increases brain serotonin levels (feed-back) – a hypothesis. D-II and D-III refer to iodothyronine deiodinases type II and III, respectively. 3,3'-T2: 3,3'-diiodothyronine.

currently favored hypothesis is that a lack of serotonin in the brain has a central role (35). This hypothesis has generated the development of effective antidepressive drugs belonging to the group of serotonin re-uptake inhibitors. TRH seems under a constant inhibition by serotonin, and reduced intracerebral serotonin concentration will lead to increased TRH concentrations in brain tissue (reviewed in 23) (Fig. 1). As a consequence, TSH secretion will be stimulated. In addition CSF levels of somatostatin are reduced in depression. Since TSH secretion is under a constant inhibition of somatostatin (23), the consequence might be a further stimulated TSH secretion. Increased TSH levels result in enhanced thyroïdal production of T_4 and T_3 . Due to the thyroid-pituitary feedback, increased T_4 and T_3 levels tend to reduce serum TSH levels, reaching a new steady state usually within the normal range. The new TSH levels are, however, inappropriately elevated in relation to the increased T_4 production (3).

Acute as well as chronic T_3 treatment has been shown to increase the serotonin levels in the cerebral cortex of rats (36). In man, plasma serotonin levels correlate positively with T_3 concentrations (37, 38), and treatment of hyperthyroidism results in a reduction in serum serotonin levels (37). Brain serotonin levels in rats seem similarly affected, i.e. synthesis is reduced in

hypothyroidism (39) and increased in hyperthyroidism (40, 41).

Thus the findings that reduced intracerebral serotonin concentrations lead to increased TRH and thereby to increased thyroid hormone levels, and that increased T_3 levels lead to an increase in brain serotonin, seem to constitute a classical feedback mechanism relevant to alleviation of depression.

Intracerebral T_3 seems mainly a result of local production by deiodination of T_4 . In general, at least three deiodinating enzymes seem capable of affecting the turnover of T_4 and T_3 (reviewed in 42). The type-I deiodinase (D-I) results in both inner (5-) and outer (5'-) ring deiodination, mainly of T_4 , which thereby provides a circulating source of T_3 to the peripheral tissues. The type-II deiodinase (D-II) results in outer (5'-) ring deiodination, mainly of T_4 . This enzyme functions to regulate intracellular T_3 levels in those tissues where T_3 is most critical, such as brain and pituitary. In line with this, the enzyme activity is increased in hypothyroidism and reduced in hyperthyroidism. The type-III (D-III) enzyme results in inner (5-) ring deiodination, of both T_4 into rT_3 , and T_3 into 3,3'-T2, and, in contrast to the D-II, but similar to D-I, its activity is increased in hyperthyroidism and decreased in hypothyroidism. Whereas all three enzymes are present in rat brain

(42), human brain tissues seem to contain only D-II and D-III enzymes (43). The properties of the D-II and D-III enzymes in rat and human tissues seem quite similar (43). Thus D-II activity increases T_3 production in brain and pituitary, and consequently also the local production of serotonin. In addition, D-III activity might be expected to decrease the local concentration of T_3 and indirectly of serotonin in brain tissues.

Studies of the influence of several known antidepressive drugs on rat D-II and D-III enzyme activities have revealed a similar pattern: lithium (44), des-imipramine (45), carbamazepine (46), and fluoxetine (47) have been shown to enhance the activity of the D-II enzyme resulting in increased local T_3 concentrations in brain tissues. Also sleep deprivation of rats, which is a known treatment modality for depression, has been shown to increase the D-II activity (48). In contrast, lithium, carbamazepine, and fluoxetine have been shown to decrease the activity of the D-III enzyme (44, 46, 47), also resulting in increased local T_3 concentration (for overview, see Fig. 1). Although this is an attractive pattern, a contradictory study has demonstrated that lithium decreases the activity of D-II in mouse neural and pituitary tissues (49). One must also be aware that the above mentioned changes are not necessarily found in the same regions of the brain, and furthermore it is not known whether these regions have any relevance to depression.

The consequence of the effect of the antidepressive drugs on the D-II and D-III enzyme activities is an increased local T_3 concentration in brain tissues and, as discussed above, this might increase local serotonin concentration, and thus might form an additional biological basis for the positive effect of these drugs in alleviating the depressive state.

Thus the hypothesis that a major pathogenic factor for depression is serotonin deficiency is sufficient to explain the changes seen in the HPT-axis. However, another hypothesis, put forward many years ago, has been that the depressed patient suffers a state of local hypothyroidism in the brain. This hypothesis, originally based on the similarities of symptoms seen in hypothyroidism and depression, seems supported by the finding that T_3 treatment alleviated the depressive symptoms (50). Reduced T_3 content in brain tissues leads to serotonin deficiency in brain tissues, and thus is sufficient to explain the findings in the HPT-axis seen in depression (Fig. 1).

Both hypotheses can explain that (i) patients with thyroid insufficiency are especially sensitive to develop depression (51, 52), (ii) T_3 treatment is beneficial also in euthyroid, depressed patients, (iii) pharmacological stimulation of the D-II and inhibition of the D-III brain enzyme could be expected to have a beneficial effect on the depression, and (iv) T_3 treatment might be at least as effective as T_4 treatment. Only the serotonin deficiency hypothesis might explain the low normal serum TSH and low Δ max TSH seen during depression, and that patients with continued low Δ max TSH after

apparent recovery from depression are subject to early relapse. This parameter can thus be regarded as a marker of continued active disease.

Another fruitful hypothesis developed for the biological explanation of depression suggested that depression was due to a state of relative brain deficiency of catecholamines, especially norepinephrine (53). This has formed the basis of the development of another group of useful antidepressive drugs, norepinephrine re-uptake inhibitors. In 1981 Whybrow & Prange Jr (54) hypothesized that thyroid hormones, by enhancing beta-adrenergic receptor function, promote transmission in central noradrenergic pathways and accelerate recovery. Later, the same group (55) demonstrated that T_3 levels in rat brain synaptosomes are much higher than whole brain levels, and that T_3 but not T_4 can be released from depolarized synaptosomes. Recently Rozanov & Dratman (56) found increased T_3 concentrations in nuclei and projection sites of rat central noradrenergic systems. These studies suggest that T_3 may play a neuromodulatory or neurotransmitter role in the noradrenergic central nervous system. Norepinephrine stimulates the release of both TRH and TSH (23), and therefore norepinephrine deficiency cannot explain the changes in the HPT-axis seen in depression, especially the increased CSF TRH levels. However the hypothesis is fully compatible with the beneficial effect of T_3 treatment in depression.

Treatment of depression with hormones of the HPT-axis

The similarities between symptoms seen in depression and untreated hypothyroidism have lead to several clinical trials in which thyroid hormones have been given to depressed patients, either alone or in combination with other antidepressive treatment modalities.

Treatment without other treatment modalities

TRH In the 1970s a number of studies demonstrated a mild but transient improvement in mood in depressed patients treated with TRH for a few days or at most 3 weeks. Later other studies including controlled double-blind designs, failed to demonstrate any beneficial effect of TRH (reviewed in 57).

T_3 T_3 given as the only drug to depressed patients has been evaluated in only two studies. Feldmesser-Reiss (50) treated 24 patients with depression or melancholia with 10–15 μ g T_3 daily, and observed considerable improvement in 10 patients. Wilson *et al.* (58) gave 25 μ g, increasing to 62.5 μ g, of T_3 daily for 9 days to nine patients, which apparently was as effective on depression as imipramine, as evaluated by the Hamilton rating scale. Imipramine plus T_3 had no additional effect. Unfortunately this otherwise randomized, double-blind study had no placebo group. Thus in

these two studies T_3 given alone seems to have an antidepressive effect, but placebo-controlled studies are lacking.

TSH and T_4 We are not aware of any studies using TSH or T_4 as the only treatment of depressed patients.

Treatment in combination with other treatment modalities

Several studies have evaluated the effect of the hormones of the HPT-axis. These studies can be separated into two typical designs: (i) either combination therapy with thyroid hormone and some kind of antidepressive treatment from the beginning, or (ii) adjunction of thyroid hormones to patients refractory to conventional antidepressive medication (nonresponders).

Initial combination therapy with T_3 or TSH in depression Concerning the initial combination therapy, only the effect of T_3 has been evaluated. Repeatedly in randomized, double-blind placebo-controlled studies, 20 to 50 μg T_3 per day in combination with TCA has been shown to shorten the period of depression as compared with TCA alone (59–62), but the final number of recovered patients was not influenced (59, 60, 62). However, in two other well-conducted trials, but with fewer patients, no additional effect of T_3 with TCA could be demonstrated (63, 64). The studies demonstrating a beneficial effect of addition of T_3 to TCA included in total 112 patients, whereas the two negative studies included only 29 patients. Furthermore the majority of patients studied were females.

The effect of T_3 does not seem to be due to changed metabolism of TCA, and might thus be regarded as an independent factor (65).

Unfortunately no studies are available concerning the effect of the addition of T_3 to the modern serotonin re-uptake inhibitory drugs.

In one study T_3 or placebo was given to patients receiving ECT (66). Patients treated with T_3 needed significantly fewer number of ECTs, and demonstrated less damage to memory functions, in comparison to the placebo group.

One study demonstrated a beneficial effect of TSH over placebo (in a randomized, double-blind fashion) when given to nine patients as intramuscular injections (10 units of TSH on day 1 and day 8) in adjunction to TCA (67). This study has unfortunately not been repeated, but it seems likely that the effect of TSH is due to an increased thyroidal secretion of T_4 and T_3 .

T_3 and T_4 in refractory depression (nonresponders)

T_3 treatment of the depressed patient resistant to TCA was first reported beneficial in an open study without a placebo group (68). Since then the positive effect of 25–50 μg T_3 daily as an adjunctive therapy has been

confirmed in many studies, including a recent meta-analysis (69), which included four randomized double-blind trials (in total 69 patients) and three unblinded studies using historical controls (in total 185 patients). This analysis also discussed in detail those studies which did not find any beneficial effect of T_3 . Overall the addition of T_3 to TCA increased the response rate significantly from 24 to 57%.

In another randomized double-blind study Joffe *et al.* (70) compared the ability of T_3 and lithium to convert nonresponders to TCA into responders, i.e. reduce the depressive symptoms on a Hamilton Rating Scale. T_3 was equally effective as lithium, and both drugs were superior to placebo.

No formal studies (only case reports) concerning the effect of the addition of T_3 in depressed patients resistant to therapy with monoamine oxidase inhibitors or serotonin re-uptake inhibitors exist.

One study has compared the effect of physiological doses of T_4 (150 μg daily) and T_3 (37.5 μg daily) as adjunctive therapy to nonresponders after 4 weeks on TCA. They found that T_3 was more effective than T_4 (71). Although randomized and double-blind, no placebo group was included, thus it is uncertain whether T_4 is superior to placebo.

Interestingly, an open, uncontrolled trial on nine L - T_4 -substituted hypothyroid patients (median 150 μg daily) with depression and no response to TCA treatment, reported that seven of these patients recovered from depression after the addition of T_3 (72), suggesting a decreased deiodination of T_4 to T_3 in brain.

In another open trial on six nonresponding patients (to TCA) with nonrapid cycling bipolar affective disorders, Baumgartner (73) gave supraphysiological doses of T_4 (250–500 μg daily) resulting in a reduced number of relapses. The results of this preliminary study are also compatible with a reduced deiodination of T_4 to T_3 .

In the rapid cycling form of bipolar affective disorders, T_4 adjunctive therapy has, in an open design without placebo controls (in total eight patients), been found to reduce both manic and depressive symptoms as well as number of relapses when given in supraphysiological doses (up to 500 μg daily) (74, 75).

These studies are all compatible with an inhibition of the D-II deiodinase or a stimulation of the D-III deiodinase in brain tissues (resulting in a reduced local T_3 concentration) in affective disorders.

Antidepressive drugs – effect on thyroid hormones

Changes in thyroid hormone and TSH levels during treatment of depressed patients with TCA and serotonin re-uptake inhibitors seem a result of the underlying psychiatric disorder, or its epiphenomena, i.e. increased

serum cortisol and starvation, and not due to a direct effect of the drug on hormone secretion, plasma protein-binding and turnover (1, 17, 76).

In contrast lithium inhibits the secretion of thyroid hormones from the thyroid gland, resulting in an increase in serum TSH (77, 78) and thyroid auto-antibodies (79), which in susceptible patients might lead to overt hypothyroidism (79). This effect of lithium might result in a decreased supply of thyroid hormones to the brain, which according to the presented hypothesis is inconvenient. However the antidepressive effect of lithium has been demonstrated to be positively correlated to serum levels of T_3 (80, 81) and negatively to serum TSH levels (82). The effects of lithium on the deiodinases D-II and D-III with the result of local increase of T_3 (see above) might counteract the inappropriate effect of lithium on the thyroid hormone secretion.

Carbamazepine, an anti-epileptic drug, which also has some antidepressive effect, influences the HPT-axis in a complex manner. In addition to the earlier mentioned effect on the brain deiodinases (Fig. 1), carbamazepine seems to enhance the hepatic (D-I) deiodination of both T_4 and T_3 (83). Thus these patients often present with low to normal serum levels of T_4 and T_3 , but normal TSH levels (83), suggestive of some inhibitory effect on the thyrotrophs as well.

Conclusions

It is hypothesized that the changes seen in the HPT-axis in untreated depression might be explained partly by a cerebral serotonin deficiency. These changes tend to increase serum levels of T_3 , which subsequently might increase serotonin content in the brain. Although this hypothesis is attractive, it is based on few and small studies, which clearly need confirmation.

Subjects with thyroid insufficiency seem especially susceptible to the development of depression.

Well established antidepressive treatment modalities influence the brain deiodinases D-II and D-III, which are responsible for the local T_3 production in brain tissues, in a manner leading to increased T_3 concentrations. Defects of the brain deiodinases might be a pathogenic factor in depression.

While the effect of the addition of physiological doses of T_3 to TCA is well documented in depression, confirmatory studies are needed concerning the effect of T_3 given alone, and T_3 given in addition to ECT.

In refractory depressions, the addition of T_3 to TCA has been shown to be equal to the addition of lithium, and superior to the addition of placebo or T_4 , to TCA in alleviating depressive symptoms. The superiority of T_3 compared with T_4 as adjunctive therapy must be further evaluated.

No studies exist on the effect of T_3 as adjunctive therapy to the serotonin re-uptake inhibitors, nor on the effect of T_3 in rapid cycling depression.

The effects of thyroid hormones, TSH and TRH, on the depressive syndrome have been extensively studied during the past three decades. Despite the fact that many of these studies are more than 20 years old, many fulfill modern criteria of good clinical science, in terms of using the randomized, double-blind placebo-controlled design. Throughout this review, we have made efforts to describe the design of the studies cited. However, the potential weakness of these studies is the relatively small number of patients enrolled in the different treatment modalities. This is especially relevant, since the depressive syndrome is a heterogenous group of disorders affecting a large number of subjects; and furthermore the diagnostic criteria have changed over the years. Therefore previous findings should be re-evaluated, using modern scientific principles, and including a large number of patients. Furthermore, the effect of especially T_3 in combination with various antidepressive treatment modalities (TCA, serotonin re-uptake inhibitors, ECT, sleep deprivation) should be studied.

The depressive syndrome is most likely a pathogenically heterogenous group. Patients are characterized phenomenologically, not by biological markers. Even phenomenologically identical groups might harbor different pathogenic subgroups of depression. Therefore future studies should also focus on identification of biological abnormalities in those patients who benefit from T_3 treatment, in order to identify in which subgroups T_3 treatment should be used.

In the light of the substantial amounts of data on the beneficial effect of T_3 in depression, the restricted usage of T_3 in this setting is surprising. T_3 given in physiological doses is cheap and the monitoring is easy due to the new sensitive TSH assays. However, precautions should be taken in patients with concurrent heart disease, since T_3 may sensitize the heart to the arrhythmogenic effects of TCA. Suicidal patients especially seem to be exposed if they take overdoses of T_3 and TCA in combination.

References

- 1 Kirkegaard C. The thyrotropin response to thyrotropin-releasing hormone in endogenous depression. *Psychoneuroendocrinology* 1981 6 189–212.
- 2 Baumgartner A, Gräf K-J, Kürten I & Meinhold H. The hypothalamic–pituitary–thyroid axis in psychiatric patients and healthy subjects: Parts 1–4. *Psychiatry Research* 1988 24 271–332.
- 3 Kirkegaard C, Körner A & Faber J. Increased production of thyroxine and inappropriately elevated serum thyrotropin levels in endogenous depression. *Biological Psychiatry* 1990 27 472–476.
- 4 Faber J. The metabolism of iodothyronines in health and disease with special reference to diiodothyronines. *Danish Medical Bulletin* 1984 31 257–270.
- 5 Kirkegaard C & Faber J. Free thyroxine and 3,3',5'-triiodothyronine levels in cerebrospinal fluid in patients with endogenous depression. *Acta Endocrinologica* 1991 124 166–172.
- 6 Docter R, Krenning EP, de Jong M & Hennemann G. The sick euthyroid syndrome: Changes in thyroid hormone serum

- parameters and hormone metabolism. *Clinical Endocrinology* 1993 **39** 499–518.
- 7 Ekins R. The free hormone hypothesis and measurement of free hormones. *Clinical Chemistry* 1992 **38** 1289–1293.
 - 8 Faber J, Rogowski P, Kirkegaard C, Siersbæk-Nielsen K & Friis T. Serum free T_4 , T_3 , rT_3 , 3,3'-diiodothyronine and 3',5'-diiodothyronine measured by ultrafiltration. *Acta Endocrinologica* 1984 **107** 357–365.
 - 9 Kirkegaard C & Faber J. Influence of free thyroid hormone levels on the TSH response to TRH in endogenous depression. *Psychoneuroendocrinology* 1986 **11** 491–497.
 - 10 Faber J, Kirkegaard C, Jørgensen B & Kludt J. The hidden, nonexchangeable pool of 3,5,3'-triiodothyronine and 3',5'-triiodothyronine in man: does it exist? *Acta Endocrinologica* 1989 **120** 667–671.
 - 11 Kirkegaard C & Faber J. Altered serum levels of thyroxine, triiodothyronines and diiodothyronines in endogenous depression. *Acta Endocrinologica* 1981 **96** 199–207.
 - 12 Linnoila M, Cowdry R, Lamberg B-A, Makinen T & Rubinow D. CSF triiodothyronine (rT_3) levels in patients with affective disorders. *Biological Psychiatry* 1983 **18** 1489–1492.
 - 13 Maes M, Vandewoude M, Maes L, Schotte C & Cosyns P. A revised interpretation of the TRH test results in female depressed patients. *Journal of Affective Disorders* 1989 **16** 203–213.
 - 14 Duval F, Macher J-P & Mokrani M-C. Difference between evening and morning thyrotropin response to protirelin in major depressive episode. *Archives of General Psychiatry* 1990 **47** 443–448.
 - 15 Maes M, Meltzer HY, Cosyns P, Suy E & Schotte C. An evaluation of basal hypothalamic-pituitary-thyroid axis function in depression: Results of a large-scaled and controlled study. *Psychoneuroendocrinology* 1993 **18** 607–620.
 - 16 Weeke A & Weeke J. Disturbed circadian variation of serum thyrotropin in patients with endogenous depression. *Acta Psychiatrica Scandinavica* 1978 **57** 281–289.
 - 17 Duval F, Mokrani M-C, Crocq M-A, Jautz M, Bailly P, Diep T-S & Macher J-P. Effect of antidepressant medication on morning and evening thyroid function tests during a major depressive episode. *Archives of General Psychiatry* 1996 **53** 833–840.
 - 18 Kirkegaard C, Ølgaard K, Bliddal H, Faber J, Friis T & Siersbæk-Nielsen K. Different responses of serum TSH and T_3 to oral TRH in various types of non-toxic goitre. *Journal of Molecular Medicine* 1980 **4** 67–73.
 - 19 Ylikorkala O, Kivinen S & Kauppila A. Oral administration of TRH in puerperal women: Effect on insufficient lactation, thyroid hormones and on the responses of TSH and prolactin to intravenous TRH. *Acta Endocrinologica* 1980 **93** 413–418.
 - 20 Kirkegaard C, Faber J, Hummer L & Rogowski P. Increased levels of TRH in cerebrospinal fluid from patients with endogenous depression. *Psychoneuroendocrinology* 1979 **4** 227–235.
 - 21 Banki CM, Bissette G, Arato M & Nemeroff CB. Elevation of immunoreactive CSF TRH in depressed patients. *American Journal of Psychiatry* 1988 **145** 1526–1531.
 - 22 Roy A, Wolkowitz OM, Bissette G & Nemeroff CB. Differences in CSF concentrations of thyrotropin-releasing hormone in depressed patients and normal subjects: Negative findings. *American Journal of Psychiatry* 1994 **151** 600–602.
 - 23 Morley JE. Neuroendocrine control of thyrotropin secretion. *Endocrine Review* 1981 **2** 396–436.
 - 24 Gerner RH & Yamada T. Altered neuropeptide concentrations in cerebrospinal fluid of psychiatric patients. *Brain Research* 1982 **238** 298–302.
 - 25 Post RM, Gold P, Rubinow DR, Ballenger JC, Bunney WE & Goodwin FK. Peptides in the cerebrospinal fluid of neuropsychiatric patients: An approach to central nervous system peptide function. *Life Sciences* 1982 **31** 1–15.
 - 26 Rubinow DR, Gold PW, Post RM, Ballenger JC, Cowdry R, Bollinger J & Reichlin S. CSF somatostatin in affective illness. *Archives of General Psychiatry* 1983 **40** 377–386.
 - 27 Gold MS, Pottash ALC & Extein I. Hypothyroidism and depression: Evidence from complete thyroid evaluation. *Journal of the American Medical Association* 1981 **245** 1919–1922.
 - 28 Oomen HAPC, Schipperijn AJM & Drexhage HA. The prevalence of affective disorder and in particular of rapid cycling of bipolar disorder in patients with abnormal thyroid function tests. *Clinical Endocrinology* 1996 **45** 215–223.
 - 29 Howland RH. Thyroid dysfunction in refractory depression: Implications for pathophysiology and treatment. *Journal of Clinical Psychiatry* 1993 **54** 47–54.
 - 30 Hickie I, Bennett B, Mitchell P, Wilhelm K & Orlay W. Clinical and subclinical hypothyroidism in patients with chronic and treatment-resistant depression. *Australian and New Zealand Journal of Physiology* 1996 **30** 246–252.
 - 31 Harris B, Othman S, Davies JA, Weppner GJ, Richards CJ, Newcombe RG, Lazarus JH, Parkes AB, Hall R & Phillips DIW. Association between postpartum thyroid dysfunction and thyroid antibodies and depression. *British Medical Journal* 1992 **305** 152–156.
 - 32 Lazarus J, Harris B, Parkes A, Richards C & Hall R. Postpartum thyroid disease (PPT) and depression – preliminary results of a randomized trial of thyroxine in prevention of psychiatric symptomatology. *Journal of Endocrinological Investigation* 1994 **17** (Suppl 1) 26.
 - 33 Krog-Meyer I, Kirkegaard C, Kijne B, Lumholtz B, Smith E, Lykke-Olsen L & Bjørum N. Prediction of relapse with the TRH test and prophylactic amitriptyline in 39 patients with endogenous depression. *American Journal of Psychiatry* 1984 **141** 945–948.
 - 34 Kvist J & Kirkegaard C. Effect of repeated sleep deprivation on clinical symptoms and the TRH test in endogenous depression. *Acta Psychiatrica Scandinavica* 1980 **62** 494–502.
 - 35 Prange Jr AJ, Wilson IC & Lynn CW. L-Tryptophan in mania: Contribution to a permissive hypothesis of affective disorders. *Archives of General Psychiatry* 1974 **30** 56–62.
 - 36 Sandrini M, Vitale G, Vergoni AV, Ottani A & Bertolini A. Effect of acute and chronic treatment with triiodothyronine on serotonin levels and serotonergic receptor subtypes in the rat brain. *Life Sciences* 1996 **58** 1551–1559.
 - 37 Upadhyaya L, Agrawal JK, Dubey GP & Udupa KN. Biogenic amines and thyrotoxicosis. *Acta Endocrinologica* 1992 **126** 315–318.
 - 38 Cleare AJ, McGregor A & O'Keane V. Neuroendocrine evidence for an association between hypothyroidism, reduced central 5-HT activity and depression. *Clinical Endocrinology* 1995 **43** 713–719.
 - 39 Singhal RL, Rastogi RB & Hrdina PD. Brain biogenic amines and altered thyroid function. *Life Sciences* 1975 **17** 1617–1626.
 - 40 Rastogi RB & Singhal RL. Influence of neonatal and adult hyperthyroidism on behavior and biosynthetic capacity for norepinephrine, dopamine and 5-hydroxytryptamine in rat brain. *Journal of Pharmacology and Experimental Therapeutics* 1976 **198** 609–618.
 - 41 Atterwill CK. Effect of acute and chronic tri-iodothyronine (T_3) administration to rats on central 5-HT and dopamine-mediated behavioural responses and related brain biochemistry. *Neuropharmacology* 1981 **20** 131–144.
 - 42 Köhrle J, Hesch RD & Leonard JL. Intracellular pathways of iodothyronines metabolism. In *Werner and Ingbar's The Thyroid: A Fundamental and Clinical Text*, edn 6, pp 144–189. Eds LE Braverman & RD Utiger. Philadelphia: Lippincott, 1991.
 - 43 Campos-Barros A, Hoell T, Musa A, Sampaolo S, Stoltzenburg G, Pinna G, Eravci M, Meinhold H & Baumgartner A. Phenolic and tyrosyl ring iodothyronine deiodination and thyroid hormone concentrations in the human central system. *Journal of Clinical Endocrinology and Metabolism* 1996 **81** 2179–2185.
 - 44 Baumgartner A, Campos-Barros A, Gaio U, Hensenius C, Frege I & Meinhold H. Effects of lithium on thyroid hormone metabolism in rat frontal cortex. *Biological Psychiatry* 1994 **36** 771–774.
 - 45 Campos-Barros A, Meinhold H, Stula M, Müller F, Köhler R, Eravci M, Putzien O & Baumgartner A. The influence of desimipramine on thyroid hormone metabolism in rat brain. *Journal of Pharmacology and Experimental Therapeutics* 1994 **268** 1143–1152.

- 46 Baumgartner A, Campos-Barros A, Gaio U, Hessenius C, Flechner A & Meinhold H. Carbamazepine affects triiodothyronine production and metabolism in rat hippocampus. *Life Sciences* 1994 **54** 401–407.
- 47 Baumgartner A, Dubeyko M, Campos-Barros A, Eravci M & Meinhold H. Subchronic administration on fluoxetine to rats affects triiodothyronine production and deiodination in regions of the cortex and in the limbic forebrain. *Brain Research* 1994 **635** 68–74.
- 48 Campos-Barros A, Köhler R, Müller F, Eravci M, Meinhold H, Wesemann W & Baumgartner A. The influence of sleep deprivation on thyroid hormone metabolism in rat frontal cortex. *Neuroscience Letters* 1993 **162** 145–148.
- 49 St Germain D. Regulatory effect of lithium on thyroxine metabolism in murine neural and anterior pituitary tissues. *Endocrinology* 1987 **120** 1430–1436.
- 50 Feldmesser-Reiss EE. The application of triiodothyronine in the treatment of mental disorders. *Journal of Neurological and Mental Disorders* 1958 **127** 540–545.
- 51 Cowdry RW, Wehr TA, Zis AP & Goodwin FK. Thyroid abnormalities associated with rapid-cycling bipolar illness. *Archives of General Psychiatry* 1983 **40** 414–420.
- 52 Bauer MS, Whybrow PC & Winokur A. Rapid cycling bipolar affective disorder. *Archives of General Psychiatry* 1990 **47** 427–432.
- 53 Schildkraut JJ. The catecholamine hypothesis of affective disorders: A review of supporting evidence. *American Journal of Psychiatry* 1965 **122** 509–522.
- 54 Whybrow PC & Prange Jr AJ. A hypothesis of thyroid–catecholamine-receptor interaction. *Archives of General Psychiatry* 1981 **38** 106–113.
- 55 Mason GA, Walker CH & Prange Jr AJ. L-Triiodothyronine: Is this peripheral hormone a central neurotransmitter? *Neuropsychopharmacology* 1993 **8** 253–258.
- 56 Rozanov CB & Dratman MB. Immunohistochemical mapping of brain triiodothyronine reveals prominent localization in central noradrenergic systems. *Neuroscience* 1996 **74** 897–915.
- 57 Stein D & Avni J. Thyroid hormones in the treatment of affective disorders. *Acta Psychiatrica Scandinavica* 1988 **77** 623–636.
- 58 Wilson IC, Prange Jr AJ & Lara PP. L-Triiodothyronine alone and with imipramine in the treatment of depressed women. In *The Thyroid Axis, Drugs, and Behavior*, pp 49–62. Ed AJ Prange Jr. New York: Raven Press, 1974.
- 59 Prange Jr AJ, Wilson IC, Rabon AM & Lipton MA. Enhancement of imipramine antidepressant activity by thyroid hormone. *American Journal of Psychiatry* 1969 **126** 457–469.
- 60 Wilson IC, Prange Jr AJ, McClane TK, Rabon AM & Lipton MA. Thyroid hormone enhancement of imipramine in nonretarded depressions. *New England Journal of Medicine* 1970 **282** 1063–1067.
- 61 Coppen A, Whybrow PC, Noguera R, Maggs R & Prange Jr AJ. The comparative antidepressant value of L-tryptophan and imipramine with and without attempted potentiation by lithium. *Archives of General Psychiatry* 1972 **26** 234–241.
- 62 Wheatley D. Potentiation of amitriptyline by thyroid hormone. *Archives of General Psychiatry* 1972 **26** 229–233.
- 63 Feighner JP, King LJ, Schuckit MA, Croughan J & Briscoe W. Hormonal potentiation of imipramine and ECT in primary depression. *American Journal of Psychiatry* 1972 **128** 1230–1238.
- 64 Steiner M, Radwan M, Elizur A, Blum I, Atsmon A & Davidson S. Failure of L-triiodothyronine (T₃) to potentiate tricyclic antidepressant response. *Current Therapeutic Research* 1978 **23** 655–659.
- 65 Garbutt J, Malekpour B, Brunswik D, Jonnalagadda MR, Jolliff L, Podolak R, Wilson I & Prange Jr AJ. Effects of triiodothyronine on drug levels and cardiac function in depressed patients treated with imipramine. *American Journal of Psychiatry* 1979 **136** 980–983.
- 66 Stern RA, Nevels CT, Shelhorse ME, Prohaska ML, Mason GA & Prange Jr AJ. Antidepressant and memory effects of combined thyroid hormone treatment and electroconvulsive therapy: Preliminary findings. *Biological Psychiatry* 1991 **30** 623–627.
- 67 Prange Jr AJ, Wilson IC, Knox A, McClane TK & Lipton MA. Enhancement of imipramine by thyroid stimulating hormone: Clinical and theoretical implications. *American Journal of Psychiatry* 1970 **127** 191–199.
- 68 Earle B. Thyroid hormone and tricyclic antidepressants in resistant depressions. *American Journal of Psychiatry* 1970 **126** 1667–1669.
- 69 Aronsen R, Offman HJ, Joffe RT & Naylor CD. Triiodothyronine augmentation in the treatment of refractory depression. *Archives of General Psychiatry* 1996 **53** 842–848.
- 70 Joffe RT, Singer W, Levitt AJ & MacDonald C. A placebo-controlled comparison of lithium and triiodothyronine augmentation of tricyclic antidepressants in unipolar refractory depression. *Archives of General Psychiatry* 1993 **50** 387–393.
- 71 Joffe RT & Singer W. A comparison of triiodothyronine and thyroxine in the potentiation of tricyclic antidepressants. *Psychiatry Research* 1990 **32** 241–251.
- 72 Cooke RG, Joffe RT & Levitt AJ. T₃ augmentation of antidepressant treatment in T₄-replaced thyroid patients. *Journal of Clinical Psychiatry* 1992 **53** 16–18.
- 73 Baumgartner A, Bauer M & Hellweg R. Treatment of intractable non-rapid cycling bipolar affective disorder with high-dose thyroxine: An open trial. *Neuropsychopharmacology* 1994 **10** 183–189.
- 74 Stancer HC & Persad E. Treatment of intractable rapid-cycling manic-depressive disorder with levothyroxine. *Archives of General Psychiatry* 1982 **39** 311–312.
- 75 Bauer MS & Whybrow PC. The effect of changing thyroid function on cyclic affective illness in a human subject. *American Journal of Psychiatry* 1986 **143** 633–636.
- 76 Linnolia M, Gold P, Potter WZ & Wehr TA. Tricyclic antidepressants do not alter thyroid hormones levels in patients suffering from a major affective disorder. *Psychiatry Research* 1981 **4** 357–360.
- 77 Emerson CH, Dyson WL & Utiger RD. Serum thyrotropin and thyroxine concentrations in patients receiving lithium carbonate. *Journal of Clinical Endocrinology and Metabolism* 1973 **36** 338–346.
- 78 Lauridsen UB, Kirkegaard C & Nerup J. Lithium and the pituitary–thyroid axis in normal subjects. *Journal of Clinical Endocrinology and Metabolism* 1974 **39** 383–385.
- 79 Myers DH, Carter RA, Burns BH, Armond A, Hussain SB & Chengapa VK. A prospective study of the effects of lithium on thyroid function and on the prevalence of antithyroid antibodies. *Psychological Medicine* 1985 **15** 55–61.
- 80 Hatterer JA, Kocsis JH & Stokes PE. Thyroid function in patients maintained on lithium. *Psychiatry Research* 1988 **26** 249–257.
- 81 Baumgartner A, von Stuckrad M, Müller-Oerlinghausen B, Gräf K-J & Kürten I. The hypothalamic–pituitary–thyroid axis in patients maintained on lithium prophylaxis for years: High triiodothyronine serum concentrations are correlated to the prophylactic efficacy. *Journal of Affective Disorders* 1995 **34** 211–218.
- 82 Prohaska ML, Stern RA, Nevels CT, Mason GA & Prange Jr AJ. The relationship between thyroid status and neuropsychological performance in psychiatric outpatients maintained on lithium. *Neuropsychiatry Neuropsychology and Behavioral Neurology* 1996 **9** 30–34.
- 83 Wenzel KW. Disturbances of thyroid function tests by drugs. *Acta Medica Austriaca* 1996 **23** 57–60.

Received 27 February 1997

Accepted 14 July 1997