Somatostatin receptors in the haematopoietic system

Sigrid P M A Oomen¹, Leo J Hofland², P Martin van Hagen^{2,3}, Steven W J Lamberts² and Ivo P Touw¹

¹Institute of Haematology, Dr Daniel den Hoed Cancer Centre, ²Department of Internal Medicine and ³Department of Immunology, Erasmus University, Rotterdam, The Netherlands

(Correspondence should be addressed to Ivo P Touw, Institute of Haematology, Erasmus University (Room H Ee 1314), PO Box 1738, 3000 DR Rotterdam, The Netherlands; Email: touw@hema.fgg.eur.nl)

Introduction

Multiple interactions exist between the immune, haematopoietic, endocrine and nervous systems (1,2). The bi-directional communication between the immune/haematopoietic and nervous systems is mediated by complex mechanisms involving multiple soluble factors (e.g. neuropeptides, neurotrophic factors, neurotransmitters and cytokines) produced by each system (3–5). Examples of such factors are: the neurotransmitter neuropeptide Y (6) produced by megakaryocytes (7), substance P, which enhances the proliferation of primitive bone marrow cells and progenitors (4), and nerve growth factor (NGF), which contributes to differentiation of human basophils (8) and stimulates the release of inflammatory mediators from these cells (9).

The bone marrow is the major site of haematopoiesis in adults. All blood cells descend from pluripotent haematopoietic stem cells, which develop in lineage-committed progenitors. These committed progenitors expand and differentiate towards functional end cells. Haematopoiesis is controlled by a complex cytokine network in combination with cellular signals provided by cell-to-cell contact with stromal elements within the bone marrow. A number of studies have demonstrated that somatostatin inhibits proliferation of lymphoid and haematopoietic cells (10). However, little is known of the effects of somatostatin on primary haematopoietic stem cells and progenitor cells. This brief overview will focus on recent insights into the expression and functional significance of somatostatin receptors (SST) in the haematopoietic system.

Haematopoiesis

Development

During mammalian embryogenesis, the haematopoietic system is formed from mesodermally derived cells localised in the aorta–gonad–mesonephros region and the yolk-sac and are predominantly pluripotent haematopoietic stem cells. At a later stage in the development of the foetus, haematopoiesis takes place in the liver and subsequently haematopoiesis shifts to the spleen and the bone marrow. The spleen then gradually becomes a less

important haematopoietic organ and, at birth, haematopoiesis in humans is almost exclusively situated in the bone marrow (11).

Haematopoiesis is a strictly regulated process. All blood cells are derived from a small population of pluripotent stem cells that are capable of self-renewal and differentiation towards distinct lineage-committed progenitor cells. These committed progenitor cells can undergo proliferation followed by terminal differentiation into the different mature blood cell types. Blood cells have a finite lifespan and must be replaced constantly throughout life. In addition to this requirement to maintain circulating cell numbers, it is also necessary to respond to host challenges with appropriately increased output of the specific cell types required. Finally, it is necessary to down regulate output when the response is no longer required. This continuous production is tightly balanced and regulated essentially by two mechanisms. Stromal cells and extracellular matrix in the bone marrow provide a suitable microenvironment required for haematopoietic cell development. In addition, a network of cytokines and haematopoietic growth factors (HGF) specifically controls the proliferation, differentiation, survival, and function of different haematopoietic cells. This network is particularly important under stress conditions, such as infection or bleeding, when rapid increases in specific blood cell types are needed.

Migration and homing of cells

During foetal development, the multipotential and self-renewing haematopoietic stem cells migrate from the foetal liver to the bone marrow. This phenomenon of targeted migration *via* the circulation to a specific tissue is referred to as ‘homing’ (12). Regulation of progenitor cell mobilisation and homing is a complex process, involving adhesion molecules, paracrine cytokines, and chemokines (Fig. 1). Two types of migration can be distinguished. Chemoattraction is the unidirectional movement of cells towards a positive gradient of a compound, whereas chemokinesis reflects activation of cell motility and an induction of cell migration in a random direction. Chemokines comprise a large number of structurally related proteins that regulate migration and activation of leukocytes through G-protein-coupled cell-surface receptors (14). The first chemoattractant reported for human CD34⁺ progenitor

Figure 1 Trafficking of haematopoietic progenitor cells. Mobilisation and homing of haematopoietic progenitor cells are multifactorial processes that involve interactions via adhesion molecules, chemokines, and paracrine cytokines. Transendothelial migration most probably has a role in haematopoietic progenitor cell trafficking. Adhesion molecules expressed on progenitor and bone marrow endothelial cells may regulate transition from resting to the circulating progenitor cell compartment and *vice versa*. Chemokines produced in the bone marrow stroma build up transendothelial gradients that may either support or inhibit migration of progenitor cells across the endothelial layer. In addition, endothelial cells can produce cytokines that influence proliferation and motility of progenitors, and haematopoietic progenitor cells may also produce cytokines that act on endothelial cells. PB: peripheral blood, BM: bone marrow. Redrafted with permission from (13).

cells is stromal-cell-derived factor-1 (SDF-1) (15). This chemokine, produced by bone marrow stromal cells, is a ligand for the chemokine receptor CXCR4. *In vitro*, SDF-1 elicits maximal transendothelial migration of ~25% of the CD34⁺ population 3 h after exposure. SDF-1 is also a chemoattractant for human lymphocytes and monocytes (16). Furthermore, SDF-1 and its receptor CXCR4 were found to be critical for murine bone marrow engraftment by human severe combined immunodeficient (SCID) repopulating stem cells (17). Besides SDF-1, stem cell factor (SCF) has been shown to elicit some chemotaxis/chemokinesis on mouse progenitor cells (18). However, the percentage of cells migrating in response to SCF was lower, and maximal migration occurred much later (8–24 h) than seen in response to SDF-1 (2–4 h), suggesting that SCF-induced migration might be attributable to indirect effects.

Somatostatin receptors in the haematopoietic system

The presence of SSTs has been demonstrated in human lymphoid tissues, lymphoid cell lines and peripheral blood cells (19–26). Both SST₂ and SST₃ transcripts have been detected in freshly isolated human thymocytes (27). In contrast, human peripheral blood B- and T-lymphocytes express only SST₃. Monocytes express SST₂ upon activation, for example by lipopolysaccharides (28). Until now, no data have been available on SST expression on haematopoietic precursors. We therefore examined the expression of SST subtypes on human bone marrow cells by RT-PCR and by flow cytometric analysis using fluorescent somatostatin. Of the five SST subtypes, SST₂ is exclusively expressed in human bone marrow cells. Interestingly, SST₂ are present on a small subset (<1%) of cells. Immunophenotypic analysis showed that this subpopulation represents the CD34⁺ fraction, with the CD117⁺ (c-kit⁺) subset of CD34⁺ cells showing the greatest expression (29). This fraction comprises the most primitive stages of haematopoietic differentiation, including the pluripotent stem cells and early lineage-committed progenitor cells.

Haematopoietic malignancies: lymphomas and leukaemias

SSTs are also present on cells derived from several haematological malignancies (20, 30–32). SSTs have been detected *in vivo* by scintigraphy using radiolabelled somatostatin analogues in both T and B non-Hodgkin's lymphoma and Hodgkin's disease (20, 30, 33, 34). Somatostatin receptor autoradiography, in which tissue sections are incubated with isotope-labelled somatostatin or somatostatin analogues, has been applied to demonstrate the presence of somatostatin binding sites in biopsy specimens from malignant lymphomas (20). Using similar methods, somatostatin binding sites have been detected on acute lymphoblastic leukaemia and acute myeloid leukaemia (AML) (31). Receptors for somatostatin have also been detected on lymphoblastic leukaemia by using a fluorescent somatostatin (32). We have recently demonstrated that, as in normal bone marrow, only SST subtype 2 is expressed on AML cells. Importantly, in contrast to normal human bone marrow, the SST₂ expression is not restricted to CD34⁺/CD117⁺ in AML cells (S P M A Oomen, unpublished observations), which might suggest that SST₂ expression in AML is not as tightly regulated as in normal cells.

Function of SST in the haematopoietic system

Inhibition of proliferation

Somatostatin is best known for its inhibitory actions on hormone secretion and cell proliferation. Moderate

Figure 2 Octreotide induces migration of human CD34⁺ cells. (A) Analysis of Fluo-somatostatin (SS) binding to CD34⁺/CD117⁺ cells. A window was set to select cells on the basis of CD34/CD117 expression. Cells were incubated in the absence (solid line) or presence (broken line) of a 100-fold excess of (D-Trp⁸)-somatostatin-14. (B) Migration assay of CD34⁺ cells in response to indicated concentrations of octreotide: migrated cells as a percentage of input cells.

inhibitory effects of somatostatin and octreotide on the *in vitro* proliferation of AML cells have been reported earlier (31). The antiproliferative effect of somatostatin on AML cells depended on the type of haematopoietic growth factor used to induce proliferation. The inhibitory effects were most prominent when AML cells had been stimulated with granulocyte colony-stimulating factor (G-CSF). The growth inhibitory effects of somatostatin have been attributed to binding and activation of a subclass of protein tyrosine phosphatase (PTPase) enzymes (35–37). Src homology domain-containing protein-tyrosine phosphatase-1 (SHP-1), previously referred to as haematopoietic cell phosphatase, has been postulated as the PTPase responsible for SST₂-mediated inhibitory growth signalling (35). We have recently studied the mechanisms underlying somatostatin responses in a myeloid cell line (mouse 32D cells) stably expressing SST₂ and G-CSF receptors. In this model, somatostatin and octreotide reduced G-CSF-induced proliferation by approximately 50%. Incubation with octreotide significantly increased the activity of SHP-1 in these cells (S P M A Oomen, unpublished observations). Because SHP-1 does not bind directly to the G-CSF receptor complex (38,39), these data fit into a hypothetical model in which SST₂ recruits SHP-1 to the plasma membrane,

where it can down modulate proliferative signals from the G-CSF receptor.

Migration

Only a few studies have dealt with effects of somatostatin on cell migration (chemotaxis) of monocytes and the results have been contradictory (38, 40, 41). In view of the observation that SST₂ is expressed on CD34⁺ bone marrow cells and because other G_{αi}-coupled receptors such as CXCR4 and IL-8 receptor have been implicated in the control of haematopoietic cell migration (15, 42, 43), we determined whether somatostatin induces migration of primitive haematopoietic cells. We found that octreotide indeed acts as a potent pro-migratory stimulus for CD34⁺ bone marrow cells (29) (Fig. 2). Using the 32D cell line model referred to above, we were able to show that octreotide acts predominantly as a chemoattractant, but has also some chemokinetic activity (Fig. 3). Finally, we observed that somatostatin also induced migration of AML cells (S P M A Oomen, unpublished observations). These data suggest that somatostatin may have an effect on the homing and migration of normal and AML cells *in vivo*, with possible implications for the clinical application of somatostatin and its analogues.

Figure 3 Chemotactic and chemokinetic activity of octreotide. Octreotide 10^{-9} mol/l (oct) was added to upper or lower chambers, or both, of a transwell as indicated. Data are expressed as the percent migration to a positive gradient (100%). These experiments were performed with a model for haematopoietic precursor cells.

Somatostatin in the haematopoietic system

Because somatostatin is rapidly degraded in the circulation, it is anticipated that the peptide acts very locally. This implies that somatostatin-producing cells must be in the vicinity of the target cells.

The bone marrow contains nerve terminals that produce multiple neuropeptides, including somatostatin (44–46). It has been suggested that nerve fibres contribute to the regulation of blood cell production and the release of blood cells from the marrow into the circulation (47, 48), although there is some controversy over these findings (49, 50). Nevertheless, it is possible that localized production of somatostatin by neural cells in the bone marrow contributes to homing of haematopoietic progenitors. An alternative, not mutually exclusive, possibility is that somatostatin is produced by the bone marrow stroma cells. Indeed, it is already known that stromal cells produce SDF-1 and SCF (16, 51). Somatostatin-producing cells were detected at the interface between bone and bone marrow, in close contact with vessels (52). At present we do not know what is the relevance of local production of somatostatin for the homing of haematopoietic progenitors *in vivo*. It also remains to be determined whether bone marrow stroma cells are capable of producing somatostatin. To date, we have been unable to detect somatostatin mRNA transcripts in a murine stromal cell line (FBMD-1). However, stroma is composed of several cell types; to determine whether bone marrow stroma cells can produce somatostatin, more stroma-derived cell lines and primary bone marrow stroma cultures have to be examined. Finally, we have examined whether somatostatin might act as an autocrine or paracrine regulator (or both) of migratory responses of bone marrow cells. We have consistently been unable to detect somatostatin mRNA in normal bone marrow and AML cells by RT-PCR, indicating that autocrine/paracrine activation of SST₂ does not have a role in migration of haematopoietic cells (S P M A Oomen, unpublished observations).

Conclusion and future perspectives

Our *in vitro* data have established that somatostatin, apart from inhibiting proliferation, exerts unique migration-inducing effects on normal and leukaemic haematopoietic progenitors. Somatostatin-induced migration may play a part in the homing and trafficking of these cells to different organs, and specific niches herein, during normal development and in pathological conditions. Therefore, a major challenge is now to establish the significance of these findings for haematopoietic stem cell migration *in vivo*.

Acknowledgements

This work was supported by funds from the Netherlands Organization for Scientific Research (NWO), and grants from the Dutch Cancer Society.

References

- Oksenberg JR, Mor-Yosef S, Ezra Y & Brauthbar C. Antigen presenting cells in human decidua tissue. II: Accessory cells for the development of anti-trinitrophenyl cytotoxic T lymphocytes. *Journal of Reproductive Immunology* 1987 **10** 309–318.
- Renold FK, Dazin P, Goetzl EJ & Payan DG. Interleukin-3 modulation of mouse bone marrow derived mast cell receptors for somatostatin. *Journal of Neuroscience Research* 1987 **18** 195–202.
- Savino W & Dardenne M. Immune-neuroendocrine interactions. *Immunology Today* 1995 **16** 318–322.
- Rameshwar P, Poddar A & Gascon P. Hematopoietic regulation mediated by interactions among the neurokinins and cytokines. *Leukemia and Lymphoma* 1997 **28** 1–10.
- Maestroni GJ. Adrenergic regulation of haematopoiesis. *Pharmacological Research* 1995 **32** 249–253.
- Lundberg JM, Terenius L, Hokfelt T, Martling CR, Tatemoto K, Mutt V *et al.* Neuropeptide Y(NPY)-like immunoreactivity in peripheral noradrenergic neurons and effects of NPY on sympathetic function. *Acta Physiologica Scandinavica* 1982 **116** 477–480.
- Ericsson A, Schalling M, McIntyre KR, Lundberg JM, Larhammar D, Seroogy K *et al.* Detection of neuropeptide Y and its mRNA in megakaryocytes: enhanced levels in certain autoimmune mice. *PNAS* 1987 **84** 5585–5589.
- Tsuda T, Wong D, Dolovich J, Bienenstock J, Marshall J & Denburg JA. Synergistic effects of nerve growth factor and granulocyte-

- macrophage colony-stimulating factor on human basophilic cell differentiation. *Blood* 1991 **77** 971–979.
- 9 Bischoff SC & Dahinden CA. Effect of nerve growth factor on the release of inflammatory mediators by mature human basophils. *Blood* 1992 **79** 2662–2669.
 - 10 van Hagen PM, Krenning EP, Kwekkeboom DJ, Reubi JC, Anker-Lugtenburg PJ, Lowenberg B *et al.* Somatostatin and the immune and haematopoietic system; a review. *European Journal of Clinical Investigation* 1994 **24** 91–99.
 - 11 Dzierzak E, Medvinsky A & de Bruijn M. Qualitative and quantitative aspects of haematopoietic cell development in the mammalian embryo. *Immunology Today* 1998 **19** 228–236.
 - 12 Gesner BM, Woodruff JJ & McCluskey RT. An autoradiographic study of the effect of neuraminidase or trypsin on transfused lymphocytes. *American Journal of Pathology* 1969 **57** 215–230.
 - 13 Mohle R, Bautz F, Rafii S, Moore MA, Brugger W & Kanz L. Regulation of transendothelial migration of hematopoietic progenitor cells. *Annals of the New York Academy of Sciences* 1999 **872** 176–185 discussion 185–186.
 - 14 Luster AD. Chemokines – chemotactic cytokines that mediate inflammation. *New England Journal of Medicine* 1998 **338** 436–445.
 - 15 Aiuti A, Webb IJ, Bleul C, Springer T & Gutierrez-Ramos JC. The chemokine SDF-1 is a chemoattractant for human CD34+ hematopoietic progenitor cells and provides a new mechanism to explain the mobilization of CD34+ progenitors to peripheral blood. *Journal of Experimental Medicine* 1997 **185** 111–120.
 - 16 Bleul CC, Fuhlbrigge RC, Casasnovas JM, Aiuti A & Springer TA. A highly efficacious lymphocyte chemoattractant, stromal cell-derived factor 1 (SDF-1) [see comments]. *Journal of Experimental Medicine* 1996 **184** 1101–1109.
 - 17 Peled A, Petit I, Kollet O, Magid M, Ponomarev T, Byk T *et al.* Dependence of human stem cell engraftment and repopulation of NOD/SCID mice on CXCR4. *Science* 1999 **283** 845–848.
 - 18 Okumura N, Tsuji K, Ebihara Y, Tanaka I, Sawai N, Koike K *et al.* Chemotactic and chemokinetic activities of stem cell factor on murine hematopoietic progenitor cells. *Blood* 1996 **87** 4100–4108.
 - 19 ten Bokum AM, Melief MJ, Schonbrunn A, van der Ham F, Lindeman J, Hoffland LJ *et al.* Immunohistochemical localization of somatostatin receptor sst2A in human rheumatoid synovium. *Journal of Rheumatology* 1999 **26** 532–535.
 - 20 van Hagen PM, Krenning EP, Reubi JC, Mulder AH, Bakker WH, Oei HY *et al.* Somatostatin analogue scintigraphy of malignant lymphomas. *British Journal of Haematology* 1993 **83** 75–79.
 - 21 van Hagen PM, Krenning EP, Reubi JC, Kwekkeboom DJ, Bakker WH, Mulder AH *et al.* Somatostatin analogue scintigraphy in granulomatous diseases. *European Journal of Nuclear Medicine* 1994 **21** 497–502.
 - 22 van Hagen PM, Markuse HM, Lamberts SW, Kwekkeboom DJ, Reubi JC & Krenning EP. Somatostatin receptor imaging. The presence of somatostatin receptors in rheumatoid arthritis. *Arthritis and Rheumatism* 1994 **37** 1521–1527.
 - 23 ten Bokum AM, Hoffland LJ, de Jong G, Bouma J, Melief MJ, Kwekkeboom DJ *et al.* Immunohistochemical localization of somatostatin receptor sst2A in sarcoid granulomas. *European Journal of Clinical Investigation* 1999 **29** 630–636.
 - 24 Reubi JC, Horisberger U, Kappeler A & Laissue JA. Localization of receptors for vasoactive intestinal peptide, somatostatin, and substance P in distinct compartments of human lymphoid organs. *Blood* 1998 **92** 191–197.
 - 25 Reubi JC, Horisberger U, Waser B, Gebbers JO & Laissue J. Preferential location of somatostatin receptors in germinal centers of human gut lymphoid tissue. *Gastroenterology* 1992 **103** 1207–1214.
 - 26 Reubi JC, Waser B, Horisberger U, Krenning E, Lamberts SW, Gebbers JO *et al.* *In vitro* autoradiographic and *in vivo* scintigraphic localization of somatostatin receptors in human lymphatic tissue. *Blood* 1993 **82** 2143–2151.
 - 27 Ferone D, van Hagen PM, Pivonello R, Colgo A, Lamberts SWJ & Hoffland LJ. Physiological and pathophysiological role of somatostatin receptors in the human thymus. *European Journal of Endocrinology* 2000 **143** (Suppl 1) S27–S34.
 - 28 Lichtenauer-Kaligis EGR, van Hagen PM, Lamberts SWJ & Hoffland LJ. Somatostatin receptor subtypes in human immune cells. *European Journal of Endocrinology* 2000 **143** (Suppl 1) S21–S25.
 - 29 Oomen SPMA, Antonissen C, van Hennik PB, Lichtenauer-Kaligis EGR, Hermans MHA, Ploemacher RE *et al.* American Society of Hematology 41st annual meeting. New Orleans, Louisiana, USA. December 3–7 1999. Abstract No. 1111. Part 1 of 2. *Blood* 1999 **94** 251A.
 - 30 Reubi JC, Waser B, van Hagen M, Lamberts SW, Krenning EP, Gebbers JO *et al.* *In vitro* and *in vivo* detection of somatostatin receptors in human malignant lymphomas. *International Journal of Cancer* 1992 **50** 895–900.
 - 31 Santini V, Lamberts SW, Krenning EP, Backx B & Lowenberg B. Somatostatin and its cyclic octapeptide analog SMS 201-995 as inhibitors of proliferation of human acute lymphoblastic and acute myeloid leukemia. *Leukemia Research* 1995 **19** 707–712.
 - 32 Hiruma K, Koike T, Nakamura H, Sumida T, Maeda T, Tomioka H *et al.* Somatostatin receptors on human lymphocytes and leukaemia cells. *Immunology* 1990 **71** 480–485.
 - 33 Lipp RW, Silly H, Ranner G, Dobnig H, Passath A, Leb G *et al.* Radiolabeled octreotide for the demonstration of somatostatin receptors in malignant lymphoma and lymphadenopathy [see comments]. *Journal of Nuclear Medicine* 1995 **36** 13–18.
 - 34 van den Anker-Lugtenburg PJ, Lowenberg B, Lamberts SW & Krenning EP. The relevance of somatostatin receptor expression in malignant lymphomas. *Metabolism* 1996 **45** 96–97.
 - 35 Lopez F, Esteve JP, Buscail L, Delesque N, Saint-Laurent N, Vaysse N *et al.* Molecular mechanisms of antiproliferative effect of somatostatin: involvement of a tyrosine phosphatase. *Metabolism* 1996 **45** 14–16.
 - 36 Reardon DB, Dent P, Wood SL, Kong T & Sturgill TW. Activation *in vitro* of somatostatin receptor subtypes 2, 3, or 4 stimulates protein tyrosine phosphatase activity in membranes from transfected Ras-transformed NIH 3T3 cells: coexpression with catalytically inactive SHP-2 blocks responsiveness. *Molecular Endocrinology* 1997 **11** 1062–1069.
 - 37 Bousquet C, Delesque N, Lopez F, Saint-Laurent N, Esteve JP, Bedecs K *et al.* sst2 somatostatin receptor mediates negative regulation of insulin receptor signaling through the tyrosine phosphatase SHP-1. *Journal of Biological Chemistry* 1998 **273** 7099–7106.
 - 38 Tapley P, Shevde NK, Schweitzer PA, Gallina M, Christianson SW, Lin IL *et al.* Increased G-CSF responsiveness of bone marrow cells from hematopoietic cell phosphatase deficient viable motheaten mice. *Experimental Hematology* 1997 **25** 122–131.
 - 39 Ward AC, Oomen SPMA, Smith L, Gits J, van Leeuwen D, Soede-Bobok AA *et al.* The SH2 domain-containing protein tyrosine phosphatase SHP-1 is induced by granulocyte colony-stimulating factor (G-CSF) and modulates signaling from the G-CSF receptor. *Leukemia* 2000 **14** 1284–1291.
 - 40 Sacerdote P, Bianchi M & Panerai AE. Human monocyte chemotactic activity of calcitonin and somatostatin related peptides: modulation by chronic peptide treatment. *Journal of Clinical Endocrinology and Metabolism* 1990 **70** 141–148.
 - 41 Wiedermann CJ, Reinisch N & Braunsteiner H. Stimulation of monocyte chemotaxis by human growth hormone and its deactivation by somatostatin. *Blood* 1993 **82** 954–960.
 - 42 Moser B, Schumacher C, von Tscharnar V, Clark-Lewis I & Baggiolini M. Neutrophil-activating peptide 2 and gro/melanoma growth-stimulatory activity interact with neutrophil-activating peptide 1/interleukin 8 receptors on human neutrophils. *Journal of Biological Chemistry* 1991 **266** 10666–10671.
 - 43 Lee J, Horuk R, Rice GC, Bennett GL, Camerato T & Wood WI. Characterization of two high affinity human interleukin-8 receptors. *Journal of Biological Chemistry* 1992 **267** 16283–16287.

- 44 Weihe E, Nohr D, Michel S, Muller S, Zentel HJ, Fink T *et al.* Molecular anatomy of the neuro-immune connection. *International Journal of Neuroscience* 1991 **59** 1–23.
- 45 Calvo W. The innervation of the bone marrow in laboratory animals. *American Journal of Anatomy* 1968 **123** 315–328.
- 46 Yamazaki K & Allen TD. Ultrastructural morphometric study of efferent nerve terminals on murine bone marrow stromal cells, and the recognition of a novel anatomical unit: the 'neuro-reticular complex'. *American Journal of Anatomy* 1990 **187** 261–276.
- 47 Afan AM, Broome CS, Nicholls SE, Whetton AD & Miyan JA. Bone marrow innervation regulates cellular retention in the murine haemopoietic system. *British Journal of Haematology* 1997 **98** 569–577.
- 48 Miyan JA, Broome CS & Afan AM. Coordinated host defense through an integration of the neural, immune and haemopoietic systems. *Domestic Animal Endocrinology* 1998 **15** 297–304.
- 49 Benestad HB, Strom-Gundersen I, Ole Iversen P, Haug E & Nja A. No neuronal regulation of murine bone marrow function [see comments]. *Blood* 1998 **91** 1280–1287.
- 50 Miyan JA, Broome CS & Whetton AD. Neural regulation of bone marrow [letter; comment]. *Blood* 1998 **92** 2971–2973.
- 51 Heinrich MC, Dooley DC, Freed AC, Band L, Hoatlin ME, Keeble WW *et al.* Constitutive expression of steel factor gene by human stromal cells. *Blood* 1993 **82** 771–783.
- 52 Godlewski A. Calcitonin and somatostatin immunoreactive cells are present in human bone marrow and bone marrow cells are responsive to calcitonin and somatostatin. *Experimental and Clinical Endocrinology* 1990 **96** 219–233.

Received 1 July 2000

Accepted 19 July 2000