EXPERIMENTAL STUDY

Interleukin-18, a proinflammatory cytokine, contributes to the pathogenesis of non-thyroidal illness mainly via the central part of the hypothalamus-pituitary-thyroid axis

Anita Boelen, Joan Kwakkel, Marianne Platvoet-ter Schiphorst, Birgit Mentrup^{1,2}, Astrid Baur¹, Josef Koehle^{1,2} and Wilmar M Wiersinga

Department of Endocrinology, Academic Medical Center, University of Amsterdam, The Netherlands, ¹Klinische Forschergruppe, Medizinische Poliklinik der Universität Würzburg, Würzburg, Germany and ²Institut für Experimentelle Endokrinologie, Charité Universitätsmedizin, Humboldt-University, Berlin, Germany

(Correspondence should be addressed to A Boelen; Email: a.boelen@amc.uva.nl)

Abstract

Objective: Proinflammatory cytokines are involved in the pathogenesis of non-thyroidal illness (NTI), as shown by studies with IL-6^{-/-} and IL-12^{-/-} mice. Interleukin (IL)-6 changes peripheral thyroid hormone metabolism, and IL-12 seems to be involved in the regulation of the central part of the hypothalamic-pituitary-thyroid (HPT) axis during illness. IL-18 is a proinflammatory cytokine which shares important biological properties with IL-12, such as interferon (IFN)- γ -inducing activity.

Design: By studying the changes in the HPT-axis during bacterial lipopolysaccharide (LPS)-induced illness in IL-18^{-/-}, IFN γ R^{-/-} and wild-type (WT) mice, we wanted to unravel the putative role of IL-18 and IFN γ in the pathogenesis of NTI.

Results: LPS induced a decrease in pituitary type 1 deiodinase (D1) activity ($P < 0.05$, ANOVA) in WT mice, but not in IL-18^{-/-} mice, while the decrease in D2 activity was similar in both strains. LPS decreased serum thyroid hormone levels and liver D1 mRNA within 24 h similarly in IL-18^{-/-}, and WT mice. The expression of IL-1, IL-6 and IFN γ mRNA expression was significantly lower in IL-18^{-/-} mice than in WT, while IL-12 mRNA expression was similar. IFN γ R^{-/-} mice had higher basal D1 activity in the pituitary than WT mice ($P < 0.05$); LPS induced a decrease of D2, but not of D1, activity in the pituitary which was similar in both strains. In the liver, the LPS-induced increase in cytokine expression was not different between IFN γ R^{-/-} mice and WT mice, and the decrease in serum T₃ and T₄ levels and hepatic D1 mRNA was also similar.

Conclusions: The relative decrease in serum T₃ and T₄ and liver D1 mRNA in response to LPS is similar in IL-18^{-/-}, IFN γ R^{-/-} and WT mice despite significant changes in hepatic cytokine induction. However, the LPS-induced decrease in D1 activity in the pituitary of WT mice is absent in IL-18^{-/-} mice; in contrast, LPS did not decrease pituitary D1 activity in the IFN γ R^{-/-} mice or their WT, which might be due to the genetic background of the mice. Our results suggest that IL-18 is also involved in the regulation of the central part of the HPT axis during illness.

European Journal of Endocrinology 151 497–502

Introduction

Changes in thyroid hormone metabolism are frequently observed in ill patients without primary thyroid diseases. These changes, also known as non-thyroidal illness (NTI) or sick euthyroid syndrome (SES), are characterized by a decrease of serum T₃ and an increase in serum rT₃ (mediated by inhibition of 5'-deiodinase (D1) and an increase in type 3 5-deiodinase (D3)), and a decrease of serum T₄ and fT₄ in severe cases. Despite these changes, serum TSH does not increase and can even be decreased. Recently, we have shown that both the peripheral and central parts of the hypothalamic-pituitary-thyroid (HPT) axis are downregulated during the acute phase of illness (1).

Proinflammatory cytokines, especially interleukin (IL)-6, IL-1 β , tumour necrosis factor (TNF)- α and interferon (IFN)- γ , are important mediators of the acute phase of illness and have inhibiting effects on peripheral thyroid hormone metabolism (2). Furthermore, we have shown in an *in vivo* NTI animal model (mice receiving a sublethal dosis of bacterial endotoxin lipopolysaccharide (LPS), resulting within 24 h in decreased serum T₃ and T₄ levels and reduced liver D1 mRNA expression (3), that IL-12 might be involved in changes in the central part (pituitary D1 and D2 activity) of the HPT axis during illness (4).

IL-12 is an important proinflammatory cytokine, synergizing in its action with IL-18, another proinflammatory cytokine. IL-18 was originally identified as

IFN γ -inducing factor (IGIF). IL-18 mRNA is expressed in a wide variety of cells, including hepatic Kupffer cells, macrophages, T cells, B cells, dendritic cells, astrocytes and microglia cells. IL-18 shares biological properties with IL-12 but is not structurally related to this cytokine. IL-18 has a primary and three-dimensional structure which resembles that of IL-1 and the IL-18 receptor system, and its signal-transduction pathway is analogous to those of the IL-1 receptor, making IL-18 part of the IL-1 family (5).

We showed that the LPS-induced decrease of D1 activity in the pituitary of female mice was not present in IL-12^{-/-} mice, indicating a role of IL-12 in alterations of the central part of the HPT axis during illness. However, it is unknown which IL-12-dependent factor is responsible for the observed alterations in the pituitary. IL-12 is required for IFN γ expression, which exerts inhibitory effects on different aspects of thyroid hormone metabolism (3). IL-12 is known to act synergistically with IL-18 to stimulate cellular IFN γ production, but by different signalling pathways. IL-18 activates NF- κ B and AP-1, while IL-12-dependent gene expression is mediated by signalling through Stat4 and Stat3 (6). Employing IL-18^{-/-} mice enables us to clarify the role of proinflammatory cytokines, especially with respect to signal-transduction pathways, in the pathogenesis of NTI. Therefore, we studied the HPT axis in our NTI model both centrally (pituitary) and peripherally (liver) in IL-18^{-/-} and IFN γ R^{-/-} mice compared to wild-type (WT) animals. By studying the HPT axis during systemic illness in IFN γ R^{-/-} mice, we are able to exclude the effects of IL-18-induced IFN γ on altered thyroid hormone metabolism.

Materials and methods

Animals

IL-18^{-/-} mice (from a C57/BL6 background) were generated by Takeda and colleagues (7). Normal C57BL/6 WT mice, used as controls for IL-18^{-/-} mice, were obtained from Harlan Sprague-Dawley (Horst, The Netherlands). IFN γ R^{-/-} mice (from a 129/Sv/Ev genetic background (8)) and 129/Sv/Ev WT mice were donated by M Kopf (Basel Institute for Immunology). Female IL-18^{-/-}, IFN γ R^{-/-} and matching WT mice were used at 6–12 weeks of age. The mice were kept in 12-h light/dark cycles in a temperature-controlled room (22 °C) with food and water *ad libitum*. A week before the experiment, the mice were housed in groups according to the experimental set-up.

Mice were injected intraperitoneally with 150 g LPS (endotoxin), *E. coli* 127:B8, Sigma), diluted in 0.5 ml sterile 0.9% NaCl. At different time points after LPS injection (t = 0, 2, 4, 8 and 24 h), six mice were anaesthetized with isoflurane and killed. Blood was taken by cardiac puncture, and serum was stored at -20 °C until analysed. The liver and pituitary were obtained

and stored immediately in liquid nitrogen. The study was approved by the local animal welfare committee.

Thyroid hormones

Serum T₃ and T₄ were measured with in-house RIA (9). To prevent interassay variation, all samples of one experiment were measured within the same assay.

RNA isolation and RT-PCR

mRNA was isolated from 10 mg liver tissue with the Magna Pure apparatus and the Magna Pure LC mRNA isolation kit II (tissue) (Roche Biochemicals, Mannheim, Germany), according to the manufacturer's protocol, and cDNA synthesis was performed with the First Strand cDNA synthesis kit for RT-PCR (AMV) (Roche Molecular Biochemicals, Mannheim, Germany). Real-time PCR (using the LightCycler, Roche Molecular Biochemicals, Mannheim, Germany) was performed for the quantitative estimation of D1, IL-1 β , IL-6, IL-12, IL-18 and IFN γ , as described before (4). All results were corrected as to their mRNA content, using HPRT mRNA as a housekeeping gene.

D1 and D2 activity

Pituitary tissues were homogenized in ice-cold homogenization buffer (250 mM sucrose, 20 mM HEPES, 1 mM ethylenediamine tetraacetic acid and 1 mM dithiothreitol (DTT), pH 7.0) by sonication. The protein contents of the homogenates were determined by a modified Bradford protein assay (BioRad, Munich, Germany), using β -globulin as protein standard (10). Total protein of 20–60 g was used for the assays.

Specific activities of D1 and D2 in pituitary homogenates were determined in parallel in the absence or presence of 1 mM PTU (6-*n*-propyl-2-thio-uracil), using 10 nM non-radioactive rT₃. The fraction of iodide release blocked by 1 mM PTU was assigned to D1. 5'-D activities of each sample were determined in triplicates and expressed as fmol ¹²⁵I released per min/mg protein for pituitary (11).

Statistics

Differences between cytokine knockout and WT mice after LPS administration were evaluated by analysis of variance with two grouping factors (time and treatment), using Excel (Microsoft). The differences between cytokine knockout and WT mice at different time points were analysed by Student's *t*-test (Excel).

Results

IL-18^{-/-} mice

Pituitary D1 and D2 activities were determined at several time points after LPS administration. The activity is

presented (Fig. 1) as a percentage of the activity (in fmol/mg per min) determined before LPS administration ($D1 = 17.4 \pm 9.9$, $D2 = 5.4 \pm 0.7$; WT (C57/Bl6): $D1 = 18.8 \pm 16.7$, $D2 = 7.6 \pm 1.8$). LPS administration induced a significant decrease of D1 activity in the pituitary of WT mice. However, no decrease of D1 activity was observed in IL-18^{-/-} mice ($P < 0.05$). The LPS-induced decrease of pituitary D2 activity was not significantly different in IL-18^{-/-} mice from WT mice.

LPS administration resulted also in markedly increased expression of liver IL-1 β , IL-6, IL-12 and IFN γ mRNA in both IL-18^{-/-} and WT mice, peaking within 4 h. IL-12 mRNA expression was similar in IL-18^{-/-} and WT mice, while IL-1 β , IL-6 and IFN γ mRNA expression was significantly impaired in IL-18^{-/-} mice ($P < 0.001$) compared to WT mice (Fig. 2).

Serum T₃ and T₄ and liver D1 mRNA expression decreased similarly within 24 h in both IL-18^{-/-} and WT mice (Fig. 3) after LPS administration. This decrease in D1 transcript levels was not due to any diurnal variation, because liver D1 mRNA expression in saline-injected WT mice increased after 8 h (data not shown). However, the serum levels of T₃ and T₄ at the beginning of the experiment (basal levels) were significantly higher in IL-18^{-/-} mice, while basal liver D1 mRNA expression seemed to be lower in IL-18^{-/-} mice (Table 1).

Figure 1 Relative changes of type 1 and type 2 5'-DI activity in the pituitary of IL-18^{-/-} (○), IFN γ R^{-/-} (○) and WT (●) mice after LPS administration. Mean values $\pm$ S.E.M. are depicted; P values indicate differences between groups by ANOVA.

IFN γ R^{-/-} mice

Pituitary D1 and D2 activity in IFN γ R^{-/-} mice is presented (Fig. 1) as a percentage of the activity (in fmol/mg per ml) determined before LPS administration (IFN γ R^{-/-}: $D1 = 45.2 \pm 10.6$, $D2 = 3.8 \pm 1.4$; WT (129/Sv/Ev): $D1 = 21.5 \pm 5.6$, $D2 = 6.0 \pm 1.7$). LPS administration resulted in an increase in pituitary D1 activity in 129/Sv/Ev WT mice, while no significant changes were observed in IFN γ R^{-/-} mice after LPS

Figure 2 Relative expression of liver IL-1 β , IL-6, IL-12, IFN γ (only in IL-18^{-/-} mice) and IL-18 (only in IFN γ R^{-/-} mice) mRNA in IL-18^{-/-} (○), IFN γ R^{-/-} (○) and WT (●) mice within 24 h after LPS administration. Mean values $\pm$ S.E.M. are depicted; P values indicate differences between groups by ANOVA.

Figure 3 Serum T_3 and T_4 levels and relative expression of liver 5'-deiodinase mRNA in $IL-18^{-/-}$ (○), $IFN\gamma R^{-/-}$ (○) and WT (●) mice within 24 h after LPS administration. Mean values $\pm$ S.E.M. are depicted; P values indicate differences between groups by ANOVA.

administration. However, the basal D1 activity was significantly higher in $IFN\gamma R^{-/-}$ than WT mice. The LPS-induced decrease of pituitary D2 activity was not different in $IFN\gamma R^{-/-}$ mice from WT mice.

Table 1 Basal thyroid hormone levels and liver D1 mRNA expression in $IL-18^{-/-}$, $IFN\gamma R^{-/-}$ and matching strains of WT mice.

	T_3 (nmol/l)	T_4 (nmol/l)	D1 mRNA (relative expression)
$IL-18^{-/-}$	$1.80 \pm 0.12^{**}$	$72.5 \pm 14.1\#$	33.0 ± 14.9
WT (C57B16)	1.48 ± 0.08	58.0 ± 5.7	43.7 ± 15.3
$IFN\gamma R^{-/-}$	$2.45 \pm 0.19^{**}$	$47.0 \pm 7.3^*$	0.38 ± 0.28
WT (129SvEv)	1.57 ± 0.47	36.6 ± 6.3	0.66 ± 0.24

Values were given as mean $\pm$ S.D.

$P = 0.06$, $IL-18^{-/-}$ vs matching WT; * $P < 0.05$, $IFN\gamma R^{-/-}$ vs matching WT; ** $P < 0.01$, $IL-18^{-/-}$ or $IFN\gamma R^{-/-}$ vs matching WT.

The LPS-induced cytokine response ($IL-1\beta$, $IL-6$, $IL-12$ and $IL-18$) in the liver did not differ in $IFN\gamma R^{-/-}$ mice and WT mice (Fig. 2).

LPS induced a similar decrease of serum T_3 , T_4 and liver D1 mRNA within 24 h in both $IFN\gamma R^{-/-}$ and WT mice (Fig. 3), although serum levels of T_3 and T_4 at the beginning of the experiment (basal levels) were significantly higher in $IFN\gamma R^{-/-}$ mice, while basal liver D1 mRNA expression seemed to be lower in $IFN\gamma R^{-/-}$ mice (Table 1). LPS-induced alterations in liver D1 mRNA expression were not due to any diurnal variation, because liver D1 mRNA expression in saline-injected WT mice increased after 8 h (data not shown).

Discussion

NTI might be viewed as part of the acute-phase response, the response of an organism to severe pathological conditions, and it is predominantly mediated by $IL-1$ -type and $IL-6$ -type cytokines. Previously, we have shown that $IL-6$, an important proinflammatory cytokine, is involved in alterations of the peripheral part of the HPT axis (12). Furthermore, $IL-12$, another proinflammatory cytokine, might be involved in changes of the central part of the HPT axis during illness, as indicated by the absence of a LPS-induced decrease in D1 activity in the pituitary in $IL-12^{-/-}$ mice (4). $IL-12$ is a potent $IFN\gamma$ -inducing cytokine and shares biological properties with $IL-18$, another proinflammatory cytokine. In the present study, we therefore evaluated the role of $IL-18$ and $IFN\gamma$ in the pathogenesis of NTI by comparing $IL-18^{-/-}$ and $IFN\gamma R^{-/-}$ mice with matching strains of WT mice.

We observed significant differences in basal thyroid hormone metabolism between $IL-18^{-/-}$ and $IFN\gamma R^{-/-}$ mice and WT mice. Serum thyroid hormone levels were higher while basal liver D1 mRNA expression was lower in $IL-18^{-/-}$ and $IFN\gamma R^{-/-}$ mice. These differences might be due to unknown genetic differences induced by genetic manipulation or to unknown inflammation-independent effects of $IL-18$ on basal $IFN\gamma$ expression ($IFN\gamma$ could interfere with several mechanisms in thyrocytes (3)), because the observed differences were present in both $IL-18^{-/-}$ and $IFN\gamma R^{-/-}$ mice. For this reason, however, we have presented our data as percentage of decrease of basal levels (relative decrease) in order to compare the LPS-induced effects on thyroid hormone metabolism.

LPS induced peripheral and central alterations of the HPT axis. The peripheral changes, characterized by a decrease of serum T_3 , T_4 and D1 mRNA expression, were similar in both $IL-18^{-/-}$ and $IFN\gamma R^{-/-}$ mice and WT mice. We have previously shown that serum TSH remained unchanged after LPS administration compared to control animals (3). The changes in liver 5'-DI mRNA expression and serum thyroid hormones, however, were similar despite different hepatic cytokine

profiles. It is known that especially IL-1 β and IFN γ might be capable of influencing 5'-DI activity or mRNA expression *in vitro* by several mechanisms (13–15), and our results suggest that the cytokines activated by LPS (relatively low levels of IL-1 β , IL-6 and IFN γ in IL-18 $^{-/-}$ mice) might be sufficient for the LPS-induced decrease of liver D1 mRNA.

The absence of a specific proinflammatory cytokine seems more important in the illness-induced alterations of the central part of the HPT axis. The anterior pituitary contains D1 and D2, which play a role in T₃ production and pituitary feedback (16, 17). The effects of IL-1 β , IL-6 and TNF α on pituitary D1 and D2 activity in reaggregates of rat anterior pituitaries were studied, and in these reaggregates, IL-1 β (dose-dependently) and TNF α increased D2 activity (18). We observed decreased pituitary D2 activity after LPS administration, which was, however, similar in IL-18 $^{-/-}$, IFN γ R $^{-/-}$ and WT mice.

In contrast, LPS administration did not result in decreased D1 activity in the pituitary of IL-18 $^{-/-}$ mice, as was observed in matching WT mice, indicating a dominant role of IL-18 in the LPS-induced decrease of pituitary D1 activity. IL-18 exerts its action via NF- κ B and AP-1 (6), and these pathways might be essential for decreasing D1 activity in the pituitary. It is known that both AP-1 and NF- κ B interact with SRC-1 and CREB-binding protein (CBP), which are needed for transcription activation by NF- κ B and AP-1 (19, 20). Yu and Koenig (13) recently showed that inhibition of T₃-dependent induction of D1 mRNA and activity by IL-1 and IL-6 (*in vitro*) can be partially overcome by exogenous SRC-1. It was therefore hypothesized that cytokine-induced competition for limiting amounts of coactivators decreases hepatic D1 gene expression during illness. In addition, activation of NF- κ B by TNF α interferes with thyroid hormone action, as demonstrated by impairment of T₃-dependent induction of D1 gene expression in HepG2 cells (21). As mentioned before, the absence of IL-18 does not have significant effects on the LPS-induced decrease in liver D1 mRNA expression. Probably the marked increase in several hepatic cytokines, which signal also via AP-1 and NF- κ B (IL-1, IL-6 and TNF α), will result in a prominent deficit of SRC-1. In the pituitary however, LPS-induced cytokine expression might be much lower; therefore, the lack of IL-18 could be critical in maintaining the balance between SRC-1 deficit (decreased D1 activity) and SRC-1 surplus (D1 activity not affected).

Furthermore, pituitary D1 regulation by T₃ occurs via TR β 2, a product of the TR β gene and the abundant isoform in the pituitary (22), while TR β 1 is the major receptor form in liver. Recently, we have shown that LPS administration results in decreased TR β 2 mRNA expression in the pituitary (1), which might contribute to decreased D1 activity in the pituitary. Competition for limiting amounts of coactivators might also play a

role in decreased pituitary TR β 2 mRNA expression, because in the pituitary a specific transcription factor, Pit-1, is present; this is essential for the expression of TR β 2 mRNA, and it interacts, among others, with CBP (23).

In view of our recent study on IL-12 $^{-/-}$ mice and the results of the IL-18 $^{-/-}$ mice, an alternative explanation could be that the lack of IL-18 or IL-12 resulted in the absence of a dominant cytokine that is essential for decreasing D1 activity in the pituitary. Both IL-12 and IL-18 induce IFN γ , which could interfere with many events in thyroid hormone metabolism, especially in the thyroid (3). It is known that LPS administration results in increased IL-12 and IL-18 expression, which subsequently synergistically enhances IFN γ mRNA transcription by activating Stat4 and AP-1. The enhanced IFN γ mRNA expression might be responsible for decreased D1 activity in the pituitary after LPS administration, as confirmed by *in vitro* studies (14). Therefore, we also studied the effects of LPS on the pituitary of IFN γ R $^{-/-}$ mice compared to their WT. However, LPS administration did not decrease pituitary D1 activity in either IFN γ R $^{-/-}$ or WT mice; in contrast, an increase in D1 activity was observed in WT mice. This could be due to the genetic background of these mice (129/SvEv), which is different from IL-12 $^{-/-}$ (Balb/c) and IL-18 $^{-/-}$ (C57/BL6) mice. Subtle differences in tissue-specific factors might prevent LPS-induced effects in the pituitary, although the absence of IFN γ signalling results in higher pituitary D1 activity (absolute values) during 24 h compared to WT mice.

In conclusion, LPS administration induced in IL-18 $^{-/-}$, IFN γ R $^{-/-}$ and WT mice a sustained fall in liver D1 mRNA accompanied by a decrease in serum T₃ and T₄, indicating that neither IL-18 nor IFN γ alone is essential in LPS-induced alterations of peripheral thyroid hormone metabolism. However, the LPS-induced decrease of D1 activity in the pituitary is absent in IL-18 $^{-/-}$ mice. The data suggest that IL-18 is involved in the regulation of the central part of the HPT axis during illness, although we cannot rule out the influence of other IL-18-related factors. Whether local production of cytokines by folliculostellate cells in the pituitary (24) or by Kupffer cells in the liver (25) contributes to tissue-specific differences in responses of deiodinase expression to LPS stimulation in these knockout mouse models remains to be studied.

Acknowledgements

We thank J Daalhuisen and I Kop (Department of Experimental Internal Medicine, Academic Medical Center) for expert biotechnical assistance, Metin Yenilmez (Humboldt University, Berlin) for measuring D1 and D2 activity, and the staff of the Laboratory of Endocrinology (Academic Medical Center) for measuring thyroid hormones. This study was supported in part by Deutsche Forschungsgemeinschaft (German

Research Foundation) grants (Ko 922/3-3 and Ko 922/7-2) to J K.

References

- Boelen A, Kwakkel J, Thijssen-Timmer DC, Alkemade A, Fliers E & Wiersinga WM. Simultaneous changes in central and peripheral components of the hypothalamus-pituitary-thyroid axis in lipopolysaccharide-induced acute illness in mice. *Journal of Endocrinology* 2004 **182** 315–323.
- Wiersinga WM. Nonthyroidal illness. In *The Thyroid*, pp 281–295. Eds LE Braverman & RD Utiger. Philadelphia: Lippincott, 2000.
- Boelen A, Platvoet-ter Schiphorst MC, Bakker O & Wiersinga WM. The role of cytokines in the lipopolysaccharide-induced sick euthyroid syndrome in mice. *Journal of Endocrinology* 1995 **146** 475–483.
- Boelen A, Kwakkel J, Platvoet-ter Schiphorst MC, Baur A, Kohrle J & Wiersinga WM. Contribution of interleukin-12 to the pathogenesis of nonthyroidal illness. *Hormone and Metabolic Research* 2004 **36** 101–106.
- Akira S. The role of IL-18 in innate immunity. *Current Opinion in Immunology* 2000 **12** 59–63.
- Nakahira M, Ahn HJ, Park WR, Gao P, Tomura M, Park CS, Hamaoka T, Ohta T, Kurimoto M & Fujiwara H. Synergy of IL-12 and IL-18 for IFN-gamma gene expression: IL-12-induced STAT4 contributes to IFN-gamma promoter activation by up-regulating the binding activity of IL-18-induced activator protein 1. *Journal of Immunology* 2002 **168** 1146–1153.
- Takeda K, Tsutsui H, Yoshimoto T, Adachi O, Yoshida N, Kishimoto T, Okamura H, Nakanishi K & Akira S. Defective NK cell activity and Th1 response in IL-18-deficient mice. *Immunity* 1998 **8** 383–390.
- Huang S, Hendriks W, Althage A, Hemmi S, Bluethmann H, Kamijo R, Vilcek J, Zinkernagel RM & Aguet M. Immune response in mice that lack the interferon-gamma receptor. *Science* 1993 **259** 1742–1745.
- Wiersinga WM & Chopra IJ. Radioimmunoassay of thyroxine (T₄), 3,5,3'-triiodothyronine (T₃), 3,3',5'-triiodothyronine (reverse T₃), and 3,3'-diiodothyronine (T₂). *Methods in Enzymology* 1982 **84** 272–303.
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Annals of Biochemistry* 1976 **72** 248–254.
- Kohrle J, Rasmussen UB, Rokos H, Leonard JL & Hesch RD. Selective affinity labeling of a 27-kDa integral membrane protein in rat liver and kidney with N-bromoacetyl derivatives of L-thyroxine and 3,5,3'-triiodo-L-thyronine. *Journal of Biological Chemistry* 1990 **265** 6146–6154.
- Boelen A, Maas MA, Lowik CW, Platvoet MC & Wiersinga WM. Induced illness in interleukin-6 (IL-6) knock-out mice: a causal role of IL-6 in the development of the low 3,5,3'-triiodothyronine syndrome. *Endocrinology* 1996 **137** 5250–5254.
- Yu J & Koenig RJ. Regulation of hepatocyte thyroxine 5'-deiodinase by T₃ and nuclear receptor coactivators as a model of the sick euthyroid syndrome. *Journal of Biological Chemistry* 2000 **275** 38296–38301.
- Jakobs TC, Mentrup B, Schmutzler C, Dreher I & Kohrle J. Proinflammatory cytokines inhibit the expression and function of human type I 5'-deiodinase in HepG2 hepatocarcinoma cells. *European Journal of Endocrinology* 2002 **146** 559–566.
- Davies PH, Sheppard MC & Franklyn JA. Inflammatory cytokines and type I 5'-deiodinase expression in phil rat liver cells. *Molecular and Cellular Endocrinology* 1997 **129** 191–198.
- Kohrle J. The selenoenzyme family of deiodinase isozymes controls local thyroid hormone availability. *Reviews of Endocrine Metabolic Disorders* 2000 **1** 49–58.
- Baur A, Buchfelder M & Kohrle J. Expression of 5'-deiodinase enzymes in normal pituitaries and in various human pituitary adenomas. *European Journal of Endocrinology* 2002 **147** 263–268.
- Baur A, Bauer K, Jarry H & Kohrle J. Effects of proinflammatory cytokines on anterior pituitary 5'-deiodinase type I and type II. *Journal of Endocrinology* 2000 **167** 505–515.
- Lee SK, Kim HJ, Na SY, Kim TS, Choi HS, Im SY & Lee JW. Steroid receptor coactivator-1 coactivates activating protein-1-mediated transactivations through interaction with the c-Jun and c-Fos subunits. *Journal of Biological Chemistry* 1998 **273** 16651–16654.
- Na SY, Lee SK, Han SJ, Choi HS, Im SY & Lee JW. Steroid receptor coactivator-1 interacts with the p50 subunit and coactivates nuclear factor kappaB-mediated transactivations. *Journal of Biological Chemistry* 1998 **273** 10831–10834.
- Nagaya T, Fujieda M, Otsuka G, Yang JP, Okamoto T & Seo H. A potential role of activated NF-kappa B in the pathogenesis of euthyroid sick syndrome. *Journal of Clinical Investigations* 2000 **106** 393–402.
- Forrest D, Reh TA & Rusch A. Neurodevelopmental control by thyroid hormone receptors. *Current Opinion in Neurobiology* 2002 **12** 49–56.
- Andersen B & Rosenfeld MG. POU domain factors in the neuroendocrine system: lessons from developmental biology provide insights into human disease. *Endocrine Reviews* 2001 **22** 2–35.
- Fauquier T, Lacampagne A, Travo P, Bauer K & Mollard P. Hidden face of the anterior pituitary. *Trends in Endocrinology and Metabolism* 2002 **13** 304–309.
- Tsukamoto HR. Redox regulation of cytokine expression in Kupffer cells. *Antioxidant Redox Signals* 2002 **4** 741–748.

Received 23 April 2004

Accepted 12 July 2004