CLINICAL STUDY

Association study of AMP-activated protein kinase subunit genes in polycystic ovary syndrome

Kari Sproul^{1,2}, Michelle R Jones³, Ricardo Azziz^{1,2,4} and Mark O Goodarzi^{1,3,4,5}

¹Department of Obstetrics and Gynecology, Cedars-Sinai Medical Center, Los Angeles, California 90048, USA, ²Department of Obstetrics and Gynecology, the David Geffen School of Medicine at UCLA, Los Angeles, California 90095, USA, ³Division of Endocrinology, Diabetes and Metabolism, Department of Medicine, Cedars-Sinai Medical Center, 8700 Beverly Boulevard, Room B-131, Los Angeles, California 90048, USA, ⁴Department of Medicine, the David Geffen School of Medicine at UCLA, Los Angeles, California 90095, USA and ⁵Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, California 90048, USA

(Correspondence should be addressed to M O Goodarzi at Division of Endocrinology, Diabetes and Metabolism, Department of Medicine, Cedars-Sinai Medical Center; Email: mark.goodarzi@cshs.org)

Abstract

Objective: To examine the genes for AMP-activated protein kinase (AMPK) subunits $\alpha 2$ (*PRKAA2*) and $\gamma 3$ (*PRKAG3*) as candidates for polycystic ovary syndrome (PCOS) and its component traits.

Design and methods: A total of 287 white PCOS women were recruited from the reproductive endocrinology clinic at the University of Alabama at Birmingham and 187 white control subjects were recruited from the surrounding community. Seven *PRKAA2* single nucleotide polymorphisms (SNPs) and four *PRKAG3* SNPs were genotyped in PCOS cases and controls. Genotyping and association analysis were performed at Cedars-Sinai Medical Center.

Results: Nominal associations of *PRKAA2* variants with insulin-related traits and the *PRKAG3* Pro71Ala variant with PCOS were not statistically significant after multiple testing correction. Among PCOS patients, there were no associations between variants in AMPK subunit genes and androgenic or reproductive traits.

Conclusions: Variants in genes for AMPK $\alpha 2$ and AMPK $\gamma 3$ were not associated with PCOS or its component traits. Our evidence does not demonstrate that AMPK is a major genetic risk factor for PCOS.

European Journal of Endocrinology 161 405–409

Introduction

Polycystic ovary syndrome (PCOS) is a common complex genetic disorder, affecting ~6–8% of reproductive age women (1, 2). PCOS is characterized by hyperandrogenism, ovulatory dysfunction, polycystic ovarian morphology, and frequently insulin resistance (IR).

AMP-activated protein kinase (AMPK) is a cellular energy sensor, acting to maintain ATP levels in the cell by sensing the AMP/ATP ratio and then phosphorylating metabolic enzymes and regulating gene expression with the net result of activating processes that generate ATP and inhibiting processes that consume ATP. AMPK influences food intake, fatty acid and glucose uptake in skeletal and cardiac muscle, insulin secretion, and hepatic fatty acid synthesis and gluconeogenesis (3). Consequently, AMPK subunit genes have been examined as candidates for type 2 diabetes susceptibility, with conflicting results (4, 5).

As PCOS is a metabolic disorder associated with an increased risk of diabetes, AMPK may play a role in this association. Metformin, commonly prescribed for PCOS patients, activates AMPK activity in skeletal muscle (6). In addition, AMPK activation appears to be decreased in

PCOS ovarian theca cells, suggesting that metformin stimulation of AMPK in the ovary itself may promote ovulation (7). We therefore examined the genes for AMPK $\alpha 2$ (*PRKAA2*) and AMPK $\gamma 3$ (*PRKAG3*) as candidate genes for PCOS and its component traits.

Subjects and methods

Subjects and phenotyping

A total of 287 consecutive unrelated white patients with PCOS, aged 13–47 years, and 187 healthy unrelated white control women, aged 14–60, were recruited from the Birmingham, AL, USA area. PCOS cases were recruited from the reproductive endocrine practice of one of the investigators (RA) at the University of Alabama at Birmingham (UAB). All subjects were non-Hispanic Caucasians. Participation in research studies was offered to patients meeting inclusion criteria (premenopausal, not pregnant, on no hormonal therapy for at least 3 months, and meeting diagnostic criteria for PCOS). PCOS was diagnosed following the 1990 NIH consensus criteria (8).

Controls were healthy women with a history of regular menstrual cycles and no family history of hirsutism. These women had no evidence of hirsutism, acne or alopecia, or endocrine dysfunction and had not taken hormonal therapy (including oral contraceptives) for at least 3 months prior to testing. Controls were recruited by word of mouth and advertisements in the Birmingham, Alabama area, through a call for 'healthy women' without detailing further the nature of the studies.

Subjects underwent a physical examination and serum androgen measurement, as per a previously described protocol (1). Fasting glucose and insulin were also obtained in a subset of the cohort (~70%). This subset did not differ demographically or hormonally from the study subjects overall. The computer-based homeostasis model assessment (HOMA, www.dtu.ox.ac.uk/homa) utilizes fasting glucose and insulin to calculate indices of IR (HOMA-IR) and insulin secretion (HOMA-%B) (9). For the insulin-related traits only, six PCOS subjects with diabetes were excluded because the hyperglycemia of diabetes may induce secondary changes in insulin-related traits that reduce their utility for genetic analysis. In addition, in PCOS cases only, we evaluated the following reproductive traits: severe oligomenorrhea (defined as >3 months between menses), infertility (yes/no), and parity (yes/no). Clinical characteristics of the subjects are given in supplementary Table 1, which can be viewed online at <http://www.eje-online.org/supplemental/>.

All subjects gave written informed consent, and the study was performed according to the guidelines of the Institutional Review Boards of UAB and Cedars-Sinai Medical Center, Los Angeles, CA, USA.

Genetic analysis

We selected six *PRKAA2* single nucleotide polymorphisms (SNPs; rs11206887, rs2143749, rs2746349, rs12749128, rs857155, and rs3738568) using genotype data of the Caucasian population of the HapMap database (release 24, phase II) (10). These SNPs capture 28 out of 35 (80%) of the Caucasian HapMap SNP alleles at $r^2 > 0.8$. An additional *PRKAA2* SNP (rs2051040) was selected because it was associated with type 2 diabetes in a Japanese cohort (4). Four validated *PRKAG3* SNPs (rs6436094, rs16859382, rs692243, and rs650898) were genotyped, selected based on frequency >5% because only two SNPs in HapMap (rs6436094, rs692243) are polymorphic in Caucasians. SNP rs692243 is a nonsynonymous variant (proline to alanine at codon 71). The eleven SNPs were genotyped in the 474 subjects using the 5'-exonuclease assay (TaqMan MGB, Applied Biosystems, Foster City, CA, USA). Supplementary Table 2, which can be viewed online at <http://www.eje-online.org/supplemental/> lists PCR primers and TaqMan MGB probes.

Statistical analysis

Unpaired *t*-tests and χ^2 tests were used to compare clinical characteristics between women with and without PCOS; quantitative trait values were log- or square root-transformed as appropriate to reduce non-normality.

Association with the presence/absence of PCOS and qualitative traits was evaluated using logistic regression, adjusting for age and body mass index (BMI) by including both as independent variables. Association with quantitative traits was evaluated using analysis of covariance, again adjusting for age and BMI. Recognizing that insulin secretion responds to ambient IR, HOMA-%B associations were also calculated with HOMA-IR as a covariate in addition to age and BMI. Quantitative trait associations were conducted in the PCOS and control groups separately. To handle multiple testing, significance was taken as $P < 0.006$, considering that we analyzed two genes against four families of traits (PCOS diagnosis, androgens, metabolic traits and reproductive endpoints), yielding a Bonferroni correction factor of 8 (i.e. eight independent comparisons). Association analyses excluding 26 post-menopausal subjects from the control group yielded the same results as analyses conducted in the entire cohort; therefore, the latter results are reported.

Results

PRKAA2

We genotyped seven SNPs spanning the *PRKAA2* gene (Fig. 1A). SNP frequencies are shown in Table 1. All markers were in Hardy–Weinberg equilibrium. Linkage disequilibrium among these markers (D') ranged from 0.45 to 1.0 (average pairwise D' of 0.90). The r^2 values ranged from 0 to 0.98 (average 0.23).

Among the women with PCOS, carriers of the minor alleles of SNPs rs2143749, rs2746349, or rs857155 had higher fasting insulin levels than noncarriers ($P = 0.044$, 0.021, and 0.027 respectively). Carriers of the minor allele of SNP rs2143749 also had increased HOMA-%B ($P = 0.030$), which was not significant when HOMA-IR was also included as a covariate ($P = 0.26$). Lastly, PCOS women who were carriers of the minor allele of SNP rs2746349 had increased HOMA-IR compared with noncarriers ($P = 0.039$). Trait values for PCOS women are shown in the electronic supplementary Table 3, which can be viewed online at <http://www.eje-online.org/supplemental/>. None of these *PRKAA2* associations were significant after considering multiple testing. We did not find significant associations between *PRKAA2* variants and PCOS itself, BMI, androgens, or reproductive traits. No significant associations were observed with quantitative traits in control subjects.

Figure 1 Gene structure and linkage disequilibrium plots for *PRKAA2* (A) and *PRKAG3* (B). Gene structure is shown at the top. *PRKAA2* has nine exons and is located on chromosome 1 (1p31). *PRKAG3* has 14 exons and is located on the reverse (–) strand of chromosome 2 (2q35). The locations of the genotyped SNPs relative to the exons are indicated. The linkage disequilibrium plots at the bottom display D' values (%) for each pair of SNPs in the box at the intersection of the diagonals from each SNP. The dark solid blocks indicate $D' = 1$ (100%) for the corresponding pair of variants. The lighter solid boxes also indicate $D' = 1$, but with a low confidence score. 3' UTR indicates SNP located in the 3' UTR.

PRKAG3

We genotyped four SNPs spanning the *PRKAG3* gene (Fig. 1B). The genotype frequencies (Table 1) for all markers were in Hardy–Weinberg equilibrium. Linkage disequilibrium among the four markers (D') ranged from 0.61 to 1.0 (average 0.91); r^2 values ranged from 0.001 to 0.55 (average 0.10).

Carriers of the minor allele of SNP rs692243 (Pro71Ala) had an increased likelihood of PCOS, with an age- and BMI-adjusted odds ratio of 1.82 (95% confidence interval, CI, 1.07–3.10, $P=0.028$), which did not meet the multiple testing-corrected P value cut-off of 0.006. We found no associations in PCOS or control women between the *PRKAG3* variants and any of the quantitative traits, including BMI.

Discussion

This is the first study evaluating association of variants in the *PRKAA2* and *PRKAG3* genes with PCOS. Nominal associations of *PRKAA2* variants with

insulin-related traits and of the *PRKAG3* Pro71Ala variant with PCOS were not statistically significant after correcting for multiple testing.

AMPK is a heterotrimeric enzyme composed of a catalytic alpha subunit (α) and two noncatalytic regulatory (β and γ) subunits. Seven genes code for the possible components of AMPK ($\alpha 1$, $\alpha 2$, $\beta 1$, $\beta 2$, $\gamma 1$, $\gamma 2$, and $\gamma 3$). We considered *PRKAA2* and *PRKAG3*, which code for the $\alpha 2$ and $\gamma 3$ subunits, as the highest priority candidate genes, given prior physiologic evidence implicating their proteins in metformin action (6) and insulin sensitivity (11, 12).

No other human genetic association studies of AMPK in PCOS have been published. A few human genetic studies examined AMPK in nonPCOS cohorts with conflicting results. Some demonstrated associations between *PRKAA2* variants and type 2 diabetes, HOMA-IR, and cholesterol fractions (4, 13, 14). One study found no association of variation in *PRKAA2* with type 2 diabetes, fasting glucose, IR, insulin secretion, or body fat distribution (5). Our *PRKAA2* results are consistent with the latter study.

Table 1 Frequency and position information on *PRKAA2* and *PRKAG3* single nucleotide polymorphisms (SNPs).

SNP designation	Alleles	Location in gene	Overall	PCOS	Control
<i>PRKAA2</i>					
rs11206887	A/G	Intron 1	0.166	0.154	0.185
rs2143749	C/G	Intron 1	0.345	0.352	0.333
rs2746349	A/G	Intron 1	0.486	0.478	0.497
rs2051040	G/A	Intron 2	0.346	0.348	0.343
rs12749128	A/G	Intron 2	0.024	0.016	0.038
rs857155	A/C	Intron 6	0.475	0.467	0.488
rs3738568	T/C	Exon 9 (3' UTR)	0.030	0.029	0.032
<i>PRKAG3</i>					
rs6436094	A/G	Exon 14 (3' UTR)	0.271	0.291	0.235
rs16859382	C/T	Intron 4	0.011	0.012	0.009
rs692243	C/G	Exon 3 (Pro71Ala)	0.195	0.233	0.135
rs650898	T/C	Intron 2	0.087	0.093	0.076

PRKAG3 variants were not associated with insulin sensitivity or secretion in prior studies (15), similar to our current results in PCOS women. Previously, the minor allele of SNP rs692243, which is a nonsynonymous variant (proline-to-alanine at codon 71), was shown to be significantly associated with higher serum LDL-cholesterol and apolipoprotein B-100 levels, supporting the functional significance of this SNP (15). We do not have cholesterol levels for our patients, but we did observe that this same SNP, rs692243, was marginally associated with PCOS itself.

Studies conducted primarily in animals documented AMPK expression in the ovary, suggesting a role for this fuel sensor in fertility (16). Recent work has found expression of AMPK in human ovaries with AMPK activation decreased in PCOS thecal cells, and that metformin treatment of these cells increased AMPK activity (7). Thus, activation of AMPK in the ovary itself could be a reason why patients with PCOS who are treated with metformin experience amelioration of ovulatory dysfunction. However, variants in the AMPK subunit genes we analyzed were not associated with ovulatory dysfunction in PCOS patients as assessed by severe oligomenorrhea, infertility, and/or parity. As reproductive traits are interrelated with metabolic traits in PCOS, the association of reproductive traits with AMPK SNPs was predictably lacking in the absence of any association of metabolic traits with these SNPs.

The value of our study is that it reports on genes that have not yet been evaluated in PCOS patients, in a homogeneous and well-characterized case/control cohort. We have performed a number of genetic association studies in this cohort from Alabama. While this may seem duplicative, it is critical to completely assess a well-defined population, minimizing the variability brought on by analyzing different genes in different populations. Thus, it is common in genetic epidemiology to use a single cohort for ongoing investigation and to report each investigation as a separate experiment, when that is how the work was performed. For example, many genetic association publications have resulted from the Insulin Resistance Atherosclerosis Family Study and the Multi-ethnic Study of Atherosclerosis.

A limitation of the current study is that IR and insulin secretion were measured by HOMA indices, which may be insufficiently reflective of these traits to allow detection of association with genetic variants, particularly in the setting of PCOS (17). This may have hindered our ability to discover robust associations of AMPK subunit genetic variants with insulin-related traits. Analysis in a cohort characterized by physiologic studies (e.g. euglycemic clamp) would be necessary to definitively rule out an association of *PRKAA2* or *PRKAG3* variants with IR.

In conclusion, variants in the genes for AMPK α 2 and AMPK γ 3 were not associated with PCOS susceptibility or the component metabolic, reproductive, or androgenic traits in PCOS patients, suggesting that *PRKAA2*

and *PRKAG3* are not major genetic risk factors for PCOS. Whether genetic variants in other AMPK subunit genes play a role in PCOS susceptibility remains to be determined.

Declaration of interest

There are no conflicts of interest related to the content of this work.

Funding

This study was supported in part by NIH grants R01-HD29364 and K24-HD01346 (to R Azziz), R01-DK79888 (to M O Goodarzi), and M01-RR00425 (General Clinical Research Center Grant from the NCCR), and an endowment from the Helping Hand of Los Angeles, Inc.

References

- 1 Azziz R, Woods KS, Reyna R, Key TJ, Knochenhauer ES & Yildiz BO. The prevalence and features of the polycystic ovary syndrome in an unselected population. *Journal of Clinical Endocrinology and Metabolism* 2004 **89** 2745–2749.
- 2 Diamanti-Kandarakis E, Kouli CR, Bergiele AT, Filandra FA, Tsianateli TC, Spina GG, Zapanti ED & Bartzis MI. A survey of the polycystic ovary syndrome in the Greek island of Lesbos: hormonal and metabolic profile. *Journal of Clinical Endocrinology and Metabolism* 1999 **84** 4006–4011.
- 3 Kahn BB, Alquier T, Carling D & Hardie DG. AMP-activated protein kinase: ancient energy gauge provides clues to modern understanding of metabolism. *Cell Metabolism* 2005 **1** 15–25.
- 4 Horikoshi M, Hara K, Ohashi J, Miyake K, Tokunaga K, Ito C, Kasuga M, Nagai R & Kadowaki T. A polymorphism in the AMPK α 2 subunit gene is associated with insulin resistance and type 2 diabetes in the Japanese population. *Diabetes* 2006 **55** 919–923.
- 5 Sun MW, Lee JY, de Bakker PI, Burt NP, Almgren P, Rastam L, Tuomi T, Gaudet D, Daly MJ, Hirschhorn JN, Altshuler D, Groop L & Florez JC. Haplotype structures and large-scale association testing of the 5' AMP-activated protein kinase genes *PRKAA2*, *PRKAB1*, and *PRKAB2* with type 2 diabetes. *Diabetes* 2006 **55** 849–855.
- 6 Musi N, Hirshman ME, Nygren J, Svanfeldt M, Bavenholm P, Rooyackers O, Zhou G, Williamson JM, Ljunqvist O, Efendic S, Moller DE, Thorell A & Goodyear LJ. Metformin increases AMP-activated protein kinase activity in skeletal muscle of subjects with type 2 diabetes. *Diabetes* 2002 **51** 2074–2081.
- 7 McAllister J. Stress signaling in the PCOS ovary. In *Proceedings of the 4th Special Scientific Meeting of the Androgen Excess and PCOS Society*, p 9. San Francisco, CA, 2008.
- 8 Zawadzki JK & Dunaif A. Diagnostic criteria for polycystic ovary syndrome: towards a rational approach. In *Polycystic Ovary Syndrome*, pp 377–384. Eds A Dunaif, JR Givens, F Haseltine & GR Merriam, Boston: Blackwell Scientific, 1992.
- 9 Levy JC, Matthews DR & Hermans MP. Correct homeostasis model assessment (HOMA) evaluation uses the computer program. *Diabetes Care* 1998 **21** 2191–2192.
- 10 The International HapMap Consortium. The International HapMap Project. *Nature* 2003 **426** 789–796.
- 11 Barnes BR, Marklund S, Steiler TL, Walter M, Hjalmar G, Amarger V, Mahlapuu M, Leng Y, Johansson C, Galuska D, Lindgren K, Abrikk M, Stapleton D, Zierath JR & Andersson L. The 5'-AMP-activated protein kinase gamma3 isoform has a key role in carbohydrate and lipid metabolism in glycolytic skeletal muscle. *Journal of Biological Chemistry* 2004 **279** 38441–38447.
- 12 Viollet B, Andreelli F, Jorgensen SB, Perrin C, Geloan A, Flamez D, Mu J, Lenzner C, Baud O, Bennoun M, Gomas E, Nicolas G,

- Wojtaszewski JF, Kahn A, Carling D, Schuit FC, Birnbaum MJ, Richter EA, Burcelin R & Vaulont S. The AMP-activated protein kinase alpha2 catalytic subunit controls whole-body insulin sensitivity. *Journal of Clinical Investigation* 2003 **111** 91–98.
- 13 Keshavarz P, Inoue H, Nakamura N, Yoshikawa T, Tanahashi T & Itakura M. Single nucleotide polymorphisms in genes encoding LKB1 (STK11), TORC2 (CRTC2) and AMPK alpha2-subunit (PRKAA2) and risk of type 2 diabetes. *Molecular Genetics and Metabolism* 2008 **93** 200–209.
- 14 Spencer-Jones NJ, Ge D, Snieder H, Perks U, Swaminathan R, Spector TD, Carter ND & O'Dell SD. AMP-kinase alpha2 subunit gene PRKAA2 variants are associated with total cholesterol, low-density lipoprotein-cholesterol and high-density lipoprotein-cholesterol in normal women. *Journal of Medical Genetics* 2006 **43** 936–942.
- 15 Weyrich P, Machicao F, Staiger H, Simon P, Thamer C, Machann J, Schick F, Guirguis A, Fritsche A, Stefan N & Haring HU. Role of AMP-activated protein kinase gamma 3 genetic variability in glucose and lipid metabolism in non-diabetic whites. *Diabetologia* 2007 **50** 2097–2106.
- 16 Dupont J, Chabrolle C, Rame C, Tosca L & Coyral-Castel S. Role of the peroxisome proliferator-activated receptors, adenosine monophosphate-activated kinase, and adiponectin in the ovary. *PPAR Research* 2008 **2008** 1–9.
- 17 Diamanti-Kandarakis E, Kouli C, Alexandraki K & Spina G. Failure of mathematical indices to accurately assess insulin resistance in lean, overweight, or obese women with polycystic ovary syndrome. *Journal of Clinical Endocrinology and Metabolism* 2004 **89** 1273–1276.

Received 17 June 2009

Accepted 29 June 2009