CLINICAL STUDY

Impaired quality of life and sexual function in overweight and obese men: the European Male Ageing Study

Thang S Han, Abdelouahid Tajar¹, Terence W O'Neill¹, Min Jiang², György Bartfai³, Steven Boonen^{4,5}, Felipe Casanueva⁶, Joseph D Finn⁷, Gianni Forti⁸, Aleksander Giwercman⁹, Ilpo T Huhtaniemi¹⁰, Krzysztof Kula¹¹, Neil Pendleton¹², Margus Punab¹³, Alan J Silman¹⁴, Dirk Vanderschueren^{5,15}, Michael E J Lean¹⁶, Frederick C W Wu⁷ and the EMAS group

Department of Endocrinology, University College London Hospital Medical School, London, UK, ¹Arthritis Research UK, Epidemiology Unit, Manchester Academic Health Science Centre, The University of Manchester, Manchester, UK, ²Department of Physiology, University of Turku, Turku, Finland, ³Department of Obstetrics, Gynaecology and Andrology, Albert Szent-Györgyi Medical University, Szeged, Hungary, ⁴Leuven University Division of Geriatric Medicine, Katholieke Universiteit Leuven, Leuven, Belgium, ⁵Leuven University Center for Metabolic Bone Diseases, Katholieke Universiteit Leuven, Leuven, Belgium, ⁶Department of Medicine, Santiago de Compostela University, Complejo Hospitalario Universitario de Santiago (CHUS), CIBER de Fisiopatología Obesidad y Nutrición (CB06/O3), Instituto Salud Carlos III, Santiago de Compostela, Spain, ⁷Department of Endocrinology, Manchester Academic Health Science Centre, Manchester Royal Infirmary, The University of Manchester, Manchester, UK, ⁸Andrology Unit, Department of Clinical Physiopathology, University of Florence, Florence, Italy, ⁹Department of Urology, Scania Andrology Centre, Malmö University Hospital, University of Lund, Lund, Sweden, ¹⁰Department of Reproductive Biology, Imperial College London, Hammersmith Campus, London, UK, ¹¹Department of Andrology and Reproductive Endocrinology, Medical University of Lodz, Lodz, Poland, ¹²Clinical Gerontology, Manchester Academic Health Science Centre, Hope Hospital, The University of Manchester, Salford, UK, ¹³Andrology Unit, United Laboratories of Tartu University Clinics, Tartu, Estonia, ¹⁴Arthritis Research UK, Chesterfield, UK, ¹⁵Leuven University Division of Endocrinology, Katholieke Universiteit Leuven, Leuven, Belgium and ¹⁶Department of Human Nutrition, University of Glasgow, Glasgow, Scotland, UK

(Correspondence should be addressed to T S Han; Email: t.s.han@doctors.org.uk)

Abstract

Background: Few published data link overweight and obesity with measures of quality of life (QoL) including sexual health in men.

Objective: To assess the association of overweight/obesity with impairment of physical and psychological QoL and sexual functions in men.

Design and setting: Cross-sectional, multicentre survey of 3369 community-dwelling men aged 40–79 (mean $\pm$ s.d., 60 ± 11) years randomly selected from eight European centres.

Outcomes: Adiposity was assessed by body mass index (BMI) and waist circumference (WC), QoL and functional impairments by physical and psychological function domains of the Short Form-36 questionnaire, Beck's Depression Inventory and the European Male Ageing Study sexual function questionnaire.

Results: Complete data on sexual activities and erectile function were available in 2734 (92%) and 3193 (95%) of the participants respectively. From the population studied, 814 men were obese (BMI ≥ 30 kg/m²) and 1171 had WC ≥ 102 cm, 25% of all men were unable to do vigorous activity and 2–13% reported depressive symptoms. Symptoms of sexual dysfunction ranged between 22% (low sexual desire) and 40% (infrequent morning erections) of the participants. Among obese men with both BMI ≥ 30 kg/m² and WC ≥ 102 cm, at least one symptom of impaired physical, psychological and sexual function was reported by 41, 43 and 73% of the participants respectively. Compared with the reference group of non-obese men (BMI < 30 kg/m² and WC < 102 cm), men with BMI ≥ 30 kg/m² and WC ≥ 102 cm more frequently reported at least one symptom of impaired physical function (odds ratio (OR)=2.67; confidence interval (CI): 2.07–3.45, $P<0.001$), impaired psychological function (OR=1.48; CI: 1.14–1.90, $P<0.01$) and impaired sexual function (OR=1.45; CI: 1.14–1.85, $P<0.01$). These functional impairments were also more prevalent in men who had WC ≥ 102 cm even with BMI < 30 kg/m², but those with BMI ≥ 30 kg/m² and WC < 102 cm generally did not suffer from increased impaired physical or sexual health. Men with high BMI and WC were at even greater likelihood of having a composite of two or more or three or more symptoms compared with those with normal BMI and WC.

Conclusions: Men with high WC, including those who are 'non-obese' with BMI < 30 kg/m², have poor QoL with symptoms of impaired physical, psychological and sexual functions. Health promotion to improve QoL should focus on prevention of obesity and central fat accumulation.

European Journal of Endocrinology 164 1003–1011

Introduction

Links between adverse adiposity and chronic illnesses are well documented (1), but surprisingly, little is published on its association with quality of life (QoL) including sexual health, especially in men. Overweight/obesity, indicated by high body mass index (BMI) or waist circumference (WC), has reached an epidemic proportion worldwide. Between the 1980s and the mid-2000s, the prevalence of obesity (BMI ≥ 30 kg/m²) has approximately doubled in Europe (2), Australia (3) and the United States (4, 5) with no sign of abating (6), and highest rates occur among South Asians, African Americans and Hispanics as well as the socioeconomically disadvantaged (7–9). In 2002, the World Health Organization reported between 25 and 75% of European adults in both non-transitional and transitional countries to be overweight and up to 20% of men and 30% of women to be obese (10). By 2004, 66% of American adults were overweight and 32% obese (8). Over the 10 years to 1999–2000, morbid obesity (BMI ≥ 40 kg/m²) increased from 0.8 to 2.2% (11). In Scotland, 1.4% of men and 3.5% of women reach BMI >40 kg/m² (12). Longitudinal studies have shown that the fastest weight increases were in young adults with little increase beyond the 5th decade (8, 13). This pattern may change as the obesity epidemic progresses.

Obesity exerts major burdens on health care in many ways, from obviously obesity-related conditions (14–16) to the hidden costs of personal care and indirect costs associated with absenteeism, disability, premature mortality and workers' compensation (17). The health care costs of obesity extend well beyond the conventional costs of obesity-related diseases, with increased prescribing costs in almost all drug classes (18). In the United States, health care spending in 2002 for obese adults was 56% higher than that for normal-weight adults, an excess of US \$1244 per person per annum (19). Expenditure on obesity-related conditions reached US \$147 billion in 2008, representing 9.1% of all medical spending compared with 6.5% in 1998 (20), reflecting disproportionate of the most expensive drug group for a variety of conditions (18).

The present analysis of the European Male Ageing Study (EMAS) database of European Union men aged 40–79 years aims to define the extent to which a range of common 'QoL' symptoms usually attributed to ageing, including impaired physical, psychological and sexual functions, are associated with overweight/obesity, as indicated by a high BMI and WC.

Methods

Subjects and study design

A total of 3369 men aged 40–79 (mean $\pm$ s.d.: 60 $\pm$ 11) years were recruited from population registers in eight

European centres, including centres from five non-transitional (Florence, Italy; Leuven, Belgium; Malmö, Sweden; Manchester, UK; Santiago de Compostela, Spain) and three transitional countries (Łódź, Poland; Szeged, Hungary; Tartu, Estonia). For the baseline survey, stratified random sampling was used with the aim of recruiting equal numbers of men in each centre and into each of four age bands (40–49, 50–59, 60–69 and 70–79 years). After completing a postal questionnaire including information about smoking habits, alcohol consumption and the presence of morbidities (hypertension, cardiovascular disease (CVD), stroke, cancer, bronchitis, asthma, peptic ulcer, epilepsy, diabetes mellitus, hepatic, renal and prostate diseases), subjects attended research clinics for a health screen as described previously (21). Each participant completed interviewer-assisted questionnaires and underwent clinical assessments including anthropometry (weight, height, WC). The questionnaires included the Short Form-36 (SF-36) health survey (22) and Beck's Depression Inventory (BDI) (23). Subjects then privately completed the EMAS sexual function questionnaire (EMAS-SFQ) (24) (Supplementary Table 1, see section on supplementary data given at the end of this article). The two physical function and psychological function domains comprising 27 dimensions from SF-36 and BDI questionnaires and the one sexual function domain comprising five dimensions were used for analysis in this study. Ethics approval for the study was obtained in accordance with local institutional requirements in each centre and participants gave informed consent.

Statistical analysis

Individuals were categorised according to their BMI (<25 , 25–30 and BMI ≥ 30 kg/m²) or the previously defined WC 'action levels' (<94 , 94–102 and ≥ 102 cm) (25, 26). For the purpose of this analysis, smoking status was categorised as either current or ever smoker, alcohol intake as frequent (5 days a week or more) and infrequent (<5 days/week) drinkers and co-morbidity as none or at least one of the health problems (hypertension, heart conditions, stroke, cancer, bronchitis, asthma, peptic ulcer, epilepsy, diabetes, liver, kidney and prostate diseases).

The χ^2 test of independence was used to assess the associations between items of impaired physical, psychological or sexual function and BMI or WC. Multivariable logistic regression analysis, with or without adjustments for potential confounding factors (age, smoking status, alcohol intake and centre), was used to estimate the likelihood of having symptoms of impaired physical, psychological and sexual functions (dependent variables) in obese men with BMI ≥ 30 kg/m² and/or WC ≥ 102 cm (independent variables). We decided that the results should be presented without adjusting for 'co-morbidity' because the 'co-morbidities' are in part secondary to adiposity, and we are interested in the

overall effect of adiposity. Finally, the same regression approach was used to assess the likelihood of having at least one symptom within each of the three domains (physical, psychological or sexual function) in different groups of men classified according to their various combinations of adiposity (BMI <30 or ≥ 30 kg/m² and WC <102 or ≥ 102 cm). To assess the burden of symptoms met by individuals, composites of two or more or three or more symptoms compared with none were also analysed in men across different categories of adiposity.

Robust standard errors were used to adjust for recruitment centre, which takes into account the hierarchy of the study design (individuals nested within centres). Results from the logistic regression models are expressed as odds ratios (ORs) and 95% confidence intervals (CIs).

Receiver operating characteristic (ROC) analysis was used to evaluate the predictive power of BMI and WC on the outcomes (impaired symptoms) in the regression models. The higher the area under the curve, i.e. the greater the curvature bowing away from the 50% line (zero prediction), the greater the predictive power. All statistical analyses were conducted using Intercooled STATA version 9.2 (Stata Corp, College Station, TX, USA).

Results

A total of 3369 (mean $\pm$ s.d. age: 60 $\pm$ 11 years) men participated in the EMAS study. Complete data concerning frequency of sexual activities (excluding men without a sexual partner) and erectile function were available in 2734 (92%) and 3193 (95%) of the participants respectively. Demographics of the EMAS sample are shown in Table 1. With regard to physical and psychological functions (Supplementary Table 2, see section on supplementary data given at the end of this article), 25% of men were unable to do vigorous activity and 2–13% reported depressive symptoms. Symptoms of impaired sexual function affected between 22% (low sexual desire), 30% (erectile dysfunction) and 40% (infrequent morning erections) of the participants. Functional impairment in these domains was consistently more prevalent in men who had either high BMI (Table 2) or WC (Table 3). Among those who had both high BMI and high WC, 41% had at least one impaired physical function (Fig. 1a), 43% had at least one symptom of impaired psychological function (Fig. 1b) and 73% had at least one of the symptoms of impaired sexual function (Fig. 1c).

Multivariable logistic regression models (Table 2) show that compared with the reference group of lean men (based on BMI <25 kg/m²), obese men (BMI ≥ 30 kg/m²) were more likely to report impaired physical function (ORs range 2.2–3.3), impaired psychological function (ORs up to 2.7) and erectile dysfunction

Table 1 Characteristics of 3369 European Union men.

Characteristics	Values
Age (years; mean (s.d.))	60.0 (11.0)
Height (cm; mean (s.d.))	173.6 (7.3)
Weight (kg; mean (s.d.))	83.5 (14.0)
BMI (kg/m ² ; mean (s.d.))	27.7 (4.1)
Waist circumference (cm; mean (s.d.))	98.5 (11.1)
Quality of life (SF-36) ^a ; mean (s.d.)	
Physical function domain	27.1 (3.8)
Physical role domain	17.2 (3.8)
Body pain domain	9.3 (2.4)
General health domain	17.9 (4.0)
Vitality domain	14.9 (3.0)
Social function domain	8.9 (1.5)
Emotion domain	13.4 (2.5)
Mental domain	20.3 (3.5)
Depressive symptoms (BDI) ^b ; mean (s.d.)	7.0 (6.5)
BMI <i>n</i> (%)	
≥ 25 and <30 kg/m ^{2c}	1629 (49.2)
≥ 30 kg/m ²	814 (24.6)
Waist circumference <i>n</i> (%)	
≥ 94 and <102 cm ^c	1006 (30.3)
≥ 102 cm	1171 (35.3)
Current smokers (<i>n</i> (%))	705 (21.1)
Alcohol intake ≥ 5 days/week (<i>n</i> (%))	762 (22.8)

^aHigher scores indicate better quality of life.

^bLower scores indicate greater depressive symptoms.

^c56 men had missing BMI data and 50 men had missing waist circumference data. The remaining men with BMI <25 kg/m² were 870 and waist circumference <94 were 1142.

(OR 1.8, 95% CI: 1.4–2.3). Similar patterns emerged in men with high WC (≥ 102 cm) (Table 3).

The symptoms from the three domains found to be associated with BMI and/or WC were further analysed in different categories of obese men and non-obese men to explore potential interaction effects of BMI and WC (Table 4). Compared with the reference group of non-obese men (BMI <30 kg/m² and WC <102 cm), men with BMI ≥ 30 kg/m² and WC ≥ 102 cm were more likely to have symptoms of impaired physical and psychological function and more likely to have erectile dysfunction. These relationships persisted, only slightly less strongly, even in men with WC ≥ 102 cm but BMI <30 kg/m². By contrast, apart from tiredness, generally there was no increased risk in these health impairments in men with high BMI but with WC <102 cm (Table 4). All the above relationships were independent of age and lifestyle factors.

Finally multivariable logistic regression was used to test the association between the presence of at least one symptom or composites of two or more and three or more symptoms within each of the three domains in men with adverse adiposity (Tables 2–4). Different combinations of groups of men with varying degrees of obesity (BMI <30 or ≥ 30 kg/m² and WC <102 or ≥ 102 cm) were used as independent variable. Table 4 shows that compared with the reference group of non-obese men (BMI <30 kg/m² and WC <102 cm), obese men with BMI ≥ 30 kg/m² and WC ≥ 102 cm had higher likelihood of reporting at least one symptom of

Table 2 Prevalence and odds ratios to estimate the risk of symptoms from three domains of physical function (Short Form-36, SF-36), psychological function (SF-36 and Beck's Depression Inventory (BDI)) and sexual symptoms (European Male Ageing Study sexual function questionnaire, EMAS-SFQ) in different categories of body mass index (BMI).

	BMI (kg/m ² ; n (%))			χ^2 test ^b	BMI (kg/m ² ; OR (95% CI)) ^a	
	<25	≥25 and <30	≥30		≥25 and <30	≥30
Physical function						
Unable to do vigorous activity	164 (19.0)	376 (23.3)	275 (34.0)	<0.001	1.18 (0.92, 1.53)	2.23 (1.59, 3.13) [‡]
Unable to do moderate activity	24 (2.8)	49 (3.0)	30 (3.7)	0.51	0.89 (0.39, 2.00)	1.20 (0.79, 1.80)
Unable to carry heavy weights	23 (2.7)	38 (2.4)	34 (4.2)	0.03	0.78 (0.39, 1.54)	1.53 (0.68, 3.42)
Unable to climb one flight of stairs	15 (1.7)	25 (1.6)	19 (2.4)	0.37	0.81 (0.47, 1.42)	1.29 (0.59, 2.80)
Unable to climb several flights of stairs	41 (4.8)	109 (6.7)	98 (12.1)	<0.001	1.38 (0.87, 2.17)	2.85 (1.76, 4.63) [‡]
Unable to bend	32 (3.7)	87 (5.4)	93 (11.5)	<0.001	1.34 (0.74, 2.40)	3.27 (1.80, 5.95) [‡]
Unable to feeding yourself	13 (1.5)	16 (1.0)	17 (2.1)	0.08	0.59 (0.25, 1.37)	1.30 (0.55, 3.07)
Unable to rise from a chair	10 (1.2)	14 (0.9)	13 (1.6)	0.26	0.67 (0.21, 2.16)	1.20 (0.59, 2.44)
Unable to walk in your home	10 (1.2)	10 (0.6)	7 (0.9)	0.36	0.49 (0.22, 1.09)	0.59 (0.27, 1.32)
Unable to bath yourself	10 (1.2)	21 (1.3)	9 (1.1)	0.91	0.98 (0.51, 1.85)	0.90 (0.38, 2.17)
Unable to walk more than 1 km	37 (4.3)	112 (6.9)	91 (11.3)	<0.001	1.62 (0.93, 2.84)	2.99 (2.05, 4.36) [‡]
Reduced work	37 (4.3)	73 (4.5)	71 (8.8)	<0.001	1.08 (0.68, 1.72)	2.30 (1.22, 4.34) [‡]
≥ 1 impaired physical function	182 (21.3)	410 (26.0)	311 (39.4)	<0.001	1.22 (0.93, 1.58)	2.56 (1.87, 3.50) [‡]
≥2 impaired physical function	68 (8.0)	188 (11.9)	153 (19.4)	<0.001	1.55 (1.00, 2.39) [*]	3.57 (2.36, 5.40) [‡]
≥3 impaired physical function	49 (5.7)	95 (6.0)	96 (12.2)	<0.001	1.05 (0.64, 1.73)	3.13 (1.92, 5.09) [‡]
Psychological function						
Down in the dumps	25 (2.9)	40 (2.5)	29 (3.6)	0.30	0.86 (0.61, 1.22)	1.32 (0.90, 1.93)
Full of life	94 (11.0)	156 (9.7)	94 (11.7)	0.27	0.83 (0.67, 1.04)	1.08 (0.84, 1.37)
Down hearted	35 (4.1)	83 (5.1)	41 (5.1)	0.47	1.35 (0.92, 1.96)	1.38 (0.75, 2.51)
Tiredness	63 (7.7)	138 (9.0)	112 (14.9)	<0.001	1.22 (0.90, 1.65)	2.20 (1.67, 2.90) [‡]
Worn out	44 (5.2)	79 (4.9)	65 (8.1)	<0.001	0.99 (0.73, 1.34)	1.73 (1.25, 2.40) [†]
Nervousness	55 (6.4)	96 (5.9)	50 (6.2)	0.88	0.98 (0.74, 1.31)	1.03 (0.90, 1.18)
Energetic	100 (11.7)	148 (9.2)	130 (16.2)	<0.001	0.75 (0.50, 1.13)	1.49 (1.09, 2.04) [*]
Irritability	27 (3.1)	38 (2.3)	30 (3.7)	0.15	0.77 (0.50, 1.19)	1.22 (0.70, 2.11)
Loss of interest	12 (1.4)	39 (2.4)	26 (3.2)	0.05	1.89 (0.56, 6.36)	2.68 (1.11, 6.46) [*]
Loss of concentration	41 (4.7)	92 (5.7)	43 (5.3)	0.61	1.13 (0.74, 1.72)	1.07 (0.67, 1.73)
Change in sleep	97 (11.2)	214 (13.2)	124 (15.4)	0.04	1.21 (0.91, 1.59)	1.45 (1.03, 2.05) [*]
Loss of energy	43 (5.0)	82 (5.1)	64 (7.9)	0.01	1.05 (0.74, 1.51)	1.80 (1.21, 2.66) [†]
Fatigue	42 (4.8)	83 (5.1)	60 (7.4)	0.04	1.05 (0.74, 1.51)	1.80 (1.21, 2.66) [†]
Sadness	36 (4.2)	70 (4.3)	33 (4.1)	0.96	1.10 (0.68, 1.79)	1.07 (0.62, 1.83)
Worthless	20 (2.3)	33 (2.0)	25 (3.1)	0.27	0.88 (0.49, 1.60)	1.44 (0.92, 2.26)
≥ 1 impaired psychological function	263 (33.1)	547 (36.1)	314 (42.8)	<0.001	1.15 (0.92, 1.44)	1.56 (1.20, 2.03) [†]
≥2 impaired psychological function	131 (16.5)	262 (17.3)	175 (23.9)	<0.001	1.11 (0.88, 1.40)	1.77 (1.30, 2.42) [‡]
≥3 impaired psychological function	82 (10.3)	156 (10.3)	108 (14.7)	0.004	1.06 (0.79, 1.42)	1.76 (1.13, 2.76) [*]
Sexual function						
Low morning erections	299 (35.6)	614 (38.7)	350 (44.5)	<0.001	0.99 (0.75, 1.30)	1.32 (0.99, 1.77)
Erectile dysfunction	206 (24.8)	470 (30.2)	286 (36.7)	<0.001	1.15 (0.84, 1.57)	1.77 (1.39, 2.25) [‡]
Low frequency of sexual thoughts	228 (26.8)	435 (27.4)	227 (28.8)	0.66	0.84 (0.70, 1.01)	0.92 (0.76, 1.12)
Low sexual desire	165 (19.4)	351 (22.1)	195 (24.7)	0.04	1.03 (0.83, 1.28)	1.24 (0.92, 1.67)
Overall sexual satisfaction	238 (28.7)	405 (26.0)	224 (28.6)	0.24	0.86 (0.77, 0.96) [†]	0.98 (0.86, 1.11)
≥ 1 impaired sexual function	512 (63.7)	955 (63.6)	530 (71.0)	0.001	0.84 (0.72, 0.99) [*]	1.20 (0.98, 1.46)
≥2 impaired sexual function	278 (34.6)	557 (37.1)	300 (40.2)	0.075	0.85 (0.64, 1.14)	1.21 (0.93, 1.56)
≥3 impaired sexual function	153 (19.0)	335 (22.3)	201 (26.9)	0.001	0.90 (0.61, 1.34)	1.59 (1.10, 2.29) [*]

* $P < 0.05$, [†] $P < 0.01$, [‡] $P < 0.001$.^aMultivariable logistic regression adjusted for age, smoking status, alcohol intake and centre (reference group: BMI <25 kg/m²).^b P values for χ^2 test of independence.

impaired physical function (OR 2.67; CI: 2.07–3.45, $P < 0.001$), at least one symptom of impaired psychological function (OR 1.48; CI: 1.14–1.90, $P < 0.01$) and at least one symptom of impaired sexual function (OR 1.45; CI: 1.14–1.85, $P < 0.01$). These health impairments persisted in men with BMI <30 kg/m² and WC ≥102 cm. By contrast, men with BMI ≥30 kg/m² and WC <102 cm generally did not show significant association with the above health impairments except for impaired psychological function (and low sexual satisfaction). Further analysis revealed that men with

BMI ≥30 kg/m² and WC ≥102 cm were at even greater likelihood of having a composite of two or more and three or more symptoms compared with those with BMI <30 kg/m² and WC <102 cm (Table 4).

The areas under the ROC curves in the prediction of three or more symptoms of physical, psychological and sexual dysfunction respectively, were 0.60, 0.55 and 0.55 by BMI and were 0.64, 0.58 and 0.60 by WC. Adding BMI to WC in the regression model did not change the power of prediction (areas under the ROC curves 0.65, 0.58 and 0.60 respectively).

Table 3 Prevalence and odds ratios to estimate the risk of symptoms from three domains of physical function (Short Form-36, SF-36), psychological function (SF-36 and Beck's Depression Inventory (BDI)) and sexual symptoms (European Male Ageing Study sexual function questionnaire, EMAS-SFQ) in different categories of waist circumference (WC).

	WC (cm; n (%))			χ^2 test ^b	WC (cm; OR (95% CI)) ^a	
	<94	≥94 and <102	≥102		≥94 and <102	≥102
Physical function						
Unable to do vigorous activity	198 (17.5)	213 (21.3)	412 (35.5)	<0.001	1.14 (0.91, 1.42)	2.26 (1.96, 2.62) [†]
Unable to do moderate activity	26 (2.3)	23 (2.3)	59 (5.1)	<0.001	0.82 (0.41, 1.65)	1.71 (1.14, 2.57) [†]
Unable to carry heavy weights	26 (2.3)	23 (2.3)	51 (4.4)	<0.001	0.85 (0.59, 1.23)	1.59 (1.05, 2.41) [*]
Unable to climb one flight of stairs	16 (1.4)	10 (1.0)	37 (3.2)	<0.001	0.64 (0.30, 1.38)	1.84 (1.27, 2.64) [†]
Unable to climb several flights of stairs	50 (4.4)	56 (5.6)	148 (12.7)	<0.001	1.15 (0.74, 1.79)	2.71 (2.02, 3.64) [†]
Unable to bend	41 (3.6)	46 (4.6)	127 (10.9)	<0.001	1.10 (0.70, 1.73)	2.71 (1.79, 4.11) [†]
Unable to feeding yourself	12 (1.1)	10 (1.0)	23 (2.0)	0.08	0.89 (0.43, 1.85)	1.54 (0.89, 2.67)
Unable to rise from a chair	11 (1.0)	9 (0.9)	20 (1.7)	0.15	0.73 (0.23, 2.34)	1.44 (0.87, 2.37)
Unable to walk in your home	11 (1.0)	7 (0.7)	12 (1.0)	0.69	0.56 (0.26, 1.21)	0.90 (0.45, 1.80)
Unable to bath yourself	11 (1.0)	10 (1.0)	19 (1.6)	0.26	0.83 (0.40, 1.72)	1.38 (0.71, 2.68)
Unable to walk more than 1 km	49 (4.3)	53 (5.3)	145 (12.5)	<0.001	1.11 (0.80, 1.52)	2.73 (2.10, 3.55) [†]
Reduced work	49 (4.3)	38 (3.8)	97 (8.3)	<0.001	0.83 (0.60, 1.15)	1.87 (1.40, 2.50) [†]
≥1 impaired physical function	224 (20.0)	226 (23.3)	461 (40.6)	<0.001	1.10 (0.89, 1.36)	2.45 (2.15, 2.78) [†]
≥2 impaired physical function	90 (8.0)	93 (9.6)	234 (20.6)	<0.001	1.11 (0.86, 1.42)	3.13 (2.58, 3.79) [†]
≥3 impaired physical function	58 (5.2)	48 (5.0)	141 (12.4)	<0.001	0.87 (0.62, 1.22)	2.91 (2.06, 4.12) [†]
Psychological function						
Down in the dumps	31 (2.8)	24 (2.4)	41 (3.5)	0.28	0.81 (0.49, 1.33)	1.24 (0.74, 2.10)
Full of life	108 (9.6)	97 (9.7)	142 (12.3)	0.07	0.93 (0.83, 1.06)	1.20 (0.97, 1.49)
Down hearted	53 (4.7)	45 (4.5)	63 (5.5)	0.56	0.95 (0.62, 1.45)	1.12 (0.91, 1.37)
Tiredness	83 (7.7)	78 (8.2)	155 (14.2)	<0.001	1.08 (0.66, 1.76)	1.96 (1.55, 2.47) [†]
Worn out	54 (4.8)	46 (4.6)	90 (7.8)	<0.001	0.96 (0.59, 1.58)	1.63 (1.27, 2.09) [†]
Nervousness	66 (5.9)	58 (5.8)	79 (6.8)	0.55	1.00 (0.62, 1.62)	1.21 (0.86, 1.70)
Energetic	104 (9.3)	98 (9.8)	178 (15.4)	<0.001	1.04 (0.72, 1.49)	1.69 (1.20, 2.37) [†]
Irritability	30 (2.6)	28 (2.8)	37 (3.2)	0.74	1.07 (0.63, 1.81)	1.22 (0.80, 1.86)
Loss of interest	13 (1.1)	20 (2.0)	43 (3.7)	<0.001	1.74 (0.73, 4.17)	3.04 (1.66, 5.59) [†]
Loss of concentration	50 (4.4)	45 (4.5)	82 (7.0)	0.01	0.94 (0.58, 1.55)	1.44 (0.97, 2.16)
Change in sleep	126 (11.1)	131 (13.1)	182 (15.6)	0.01	1.19 (0.93, 1.54)	1.42 (0.99, 2.03)
Loss of energy	54 (4.8)	33 (3.3)	104 (8.9)	<0.001	0.68 (0.35, 1.31)	1.87 (1.35, 2.59) [†]
Fatigue	56 (4.9)	46 (4.6)	86 (7.4)	0.01	0.68 (0.35, 1.31)	1.87 (1.35, 2.59) [†]
Sadness	41 (3.6)	46 (4.6)	53 (4.6)	0.43	1.29 (0.78, 2.12)	1.28 (0.78, 2.10)
Worthless	27 (2.4)	18 (1.8)	33 (2.8)	0.28	0.77 (0.46, 1.28)	1.13 (0.73, 1.75)
≥1 impaired psychological function	348 (33.0)	324 (34.9)	458 (42.8)	<0.001	1.05 (0.83, 1.32)	1.44 (1.22, 1.71) [†]
≥2 impaired psychological function	157 (14.9)	155 (16.7)	261 (24.4)	<0.001	1.11 (0.88, 1.39)	1.78 (1.42, 2.24) [†]
≥3 impaired psychological function	98 (9.3)	89 (9.6)	162 (15.1)	<0.001	1.03 (0.76, 1.39)	1.78 (1.28, 2.48) [†]
Sexual function						
Low morning erections	371 (33.5)	389 (39.4)	509 (45.2)	<0.001	1.09 (0.88, 1.35)	1.33 (1.06, 1.67) [*]
Erectile dysfunction	258 (23.5)	280 (29.0)	426 (38.3)	<0.001	1.32 (0.99, 1.75)	1.73 (1.50, 1.98) [†]
Low frequency of sexual thoughts	264 (23.6)	272 (27.7)	355 (31.3)	<0.001	1.04 (0.80, 1.34)	1.06 (0.84, 1.35)
Low sexual desire	208 (18.6)	194 (19.7)	310 (27.4)	<0.001	1.41 (1.10, 1.81) [†]	1.31 (0.96, 1.79)
Overall sexual satisfaction	290 (26.7)	257 (26.4)	323 (28.9)	0.37	1.02 (0.83, 1.27)	1.14 (0.97, 1.33)
≥1 impaired sexual function	635 (59.9)	590 (63.5)	777 (72.8)	<0.001	1.28 (1.12, 1.46) [†]	1.45 (1.14, 1.85) [†]
≥2 impaired sexual function	339 (32.0)	338 (36.4)	460 (43.1)	<0.001	1.37 (1.00, 1.88) [*]	1.48 (1.13, 1.94) [†]
≥3 impaired sexual function	188 (17.7)	200 (21.5)	303 (28.4)	<0.001	1.40 (0.90, 2.19)	1.87 (1.28, 2.72) [†]

* $P < 0.05$, $^{\dagger}P < 0.01$, $^{\ddagger}P < 0.001$.^aMultivariable logistic regression adjusted for age, smoking status, alcohol intake and centre (reference group: WC <94 cm).^b P values for χ^2 test of independence.

Discussion

This study has shown that obese men, with high BMI and/or high WC, are at greater risk for a range of symptoms encompassing physical, psychological and sexual domains that can diminish QoL independent of age and lifestyle factors. A large WC plays a key role as a predictor of poor QoL, as evident by the multidomain symptomatic functional impairments even in men who are not considered to be 'obese' based on BMI ≥ 30 kg/m². This is likely to be because in subjects

with similar BMI, variations in WC also reflect differences in intra-abdominal fat mass. High BMI has been shown to associate with erectile dysfunction in previous cross-sectional and prospective studies (27, 28), and WC has been shown to be a better predictor of erectile dysfunction than BMI (29, 30). These findings are particularly important because the prevalence of obesity indicated by high BMI, which is equally matched by that of large WC, continues to rise worldwide, consuming some 10% of health care resources (3, 9, 31–34). This further strengthens the justification for

public health preventative measures in combating the rising prevalence of obesity, providing additional incentivisation for individuals to improve QoL. Weight management should target younger age groups when fat accumulation is most rapid (13), particularly those

who are approaching WC action level 1 (WC=94 cm) (25) and high-risk groups such as those with strong family history of obesity and related complications more commonly observed in ethnic groups. The public is now increasingly informed of obesity-related health risks through education and health promotion programmes but the people at highest risk are those in lower socioeconomic/education levels (8).

In epidemiological studies, BMI is widely used as a proxy for body fatness, and it is also influenced by muscle mass. WC also correlates highly with total body fat and with intra-abdominal fat when the relationship is not overwhelmed by variations in total body fat (35). WC is associated with a range of health consequences including hypertension, diabetes mellitus, coronary heart disease, respiratory disease, musculoskeletal disorders, cancers and premature death (36–39). The initial observations in cross-sectional studies have all been confirmed in longitudinal cohort studies (40–42). WC generally shows less stronger associations with health outcomes than BMI, possibly because it is not influenced by variations in muscle mass. Greater muscle mass is likely to reflect better health and greater physical activity but tends to elevate BMI in certain individuals. The confounding effect of variations in muscle mass on BMI is likely to be particularly important in studies involving ageing where sarcopenia may increasingly influence BMI relationships.

The main strengths of our study are that it is based on a large population-based sample that used uniform methods to assess depressed mood, physical activity and physical function. Methodological limitations inherent to the EMAS study have previously been described in detail (21), although a number of specific factors need to be considered in interpreting the results presented here. The overall response rate for participation in the study was 43%. Those who participated may have differed with respect to symptoms and anthropometric measures from those who did not participate, thus some caution is needed in interpreting the data. The main findings, however, were based on an internal comparison of responders, and therefore any selection factors were unlikely to have had any important effect on these data. The results were obtained from predominantly Caucasian European men and should be extrapolated beyond this setting with care. The findings from this cross-sectional study cannot be assumed to be causal because poor health *per se* can lead to obesity and central fat accumulation, but a causal relationship is at least very plausible, as supported by the longitudinal studies cited earlier (37, 38). In this study, categorisation of BMI and WC was based on conventional cut-offs used for clinical and health promotion purposes worldwide, as defined by the World Health Organization (43) for BMI (25 and 30 kg/m²) and by the National Cholesterol Education Program (26), as originally proposed by Lean *et al.* (25) for WC action levels (94 and 102 cm) respectively.

Figure 1 Proportions (numbers in brackets) of men with at least one symptom of impaired physical function (a), and at least one symptom of impaired psychological function (b) and at least one symptom of impaired sexual function (c) in different categories according to the degrees of their adiposity assessed by BMI and by waist circumference. Full colour version of this figure available via <http://dx.doi.org/10.1530/EJE-10-1129>.

Table 4 Odds ratios to estimate the risk of impaired physical function (Short Form-36, SF-36), psychological dysfunction (SF-36 and Beck's Depression Inventory (BDI)) and sexual dysfunction (European Male Ageing Study sexual function questionnaire, EMAS-SFQ) in men with either high body mass index (BMI) or high waist circumference (WC) or both of these indices of adverse adiposity.

	OR (95% CI) ^a		
	BMI ≥ 30 kg/m ² and WC ≥ 102 cm	BMI < 30 kg/m ² and WC ≥ 102 cm	BMI ≥ 30 kg/m ² and WC < 102 cm
Physical function			
Unable to do vigorous activity	2.33 (1.73, 3.14) [†]	1.78 (1.42, 2.22) [†]	1.41 (0.61, 3.22)
Unable to do moderate activity	1.61 (1.25, 2.08) [†]	1.78 (1.06, 2.99) [*]	NA
Unable to carry heavy weights	1.91 (1.18, 3.10) [†]	0.95 (0.58, 1.56)	NA
Unable to climb one flight of stairs	1.86 (0.89, 3.90)	2.26 (1.27, 4.01) [†]	1.44 (0.27, 7.65)
Unable to climb several flights of stairs	2.81 (2.05, 3.85) [†]	1.94 (1.13, 3.34) [*]	1.14 (0.54, 2.38)
Unable to bend	3.18 (2.35, 4.28) [†]	1.74 (1.08, 2.81) [*]	1.34 (0.53, 3.40)
Unable to feeding yourself	1.97 (1.11, 3.51) [*]	0.98 (0.39, 2.45)	NA
Unable to rise from a chair	1.74 (0.97, 3.12)	1.26 (0.57, 2.78)	NA
Unable to walk in your home	0.98 (0.63, 1.53)	1.02 (0.37, 2.83)	NA
Unable to bath yourself	1.17 (0.59, 2.32)	1.96 (0.83, 4.63)	NA
Unable to walk more than 1 km	2.73 (1.87, 4.00) [†]	1.98 (1.47, 2.69) [†]	NA
Reduced work	2.46 (1.47, 4.10) [†]	1.20 (0.83, 1.72)	0.39 (0.07, 2.11)
≥ 1 impaired physical function	2.67 (2.07, 3.45) [†]	1.80 (1.53, 2.13) [†]	1.30 (0.56, 3.03)
≥ 2 impaired physical function	3.38 (2.36, 4.83) [†]	2.21 (1.73, 2.81) [†]	0.91 (0.51, 1.64)
≥ 3 impaired physical function	3.79 (2.59, 5.56) [†]	1.87 (1.23, 2.84) [†]	0.39 (0.09, 1.64)
Psychological function			
Down in the dumps	1.51 (1.08, 2.10) [*]	0.86 (0.39, 1.90)	0.61 (0.07, 5.73)
Full of life	1.27 (0.93, 1.72)	1.12 (0.82, 1.54)	0.98 (0.48, 1.99)
Down hearted	1.17 (0.80, 1.72)	1.03 (0.68, 1.55)	0.68 (0.17, 2.70)
Tiredness	2.14 (1.52, 3.00) [†]	1.65 (1.15, 2.36) [†]	2.23 (1.24, 4.00) [†]
Worn out	1.89 (1.27, 2.81) [†]	1.45 (1.08, 1.94) [*]	2.07 (0.76, 5.60)
Nervousness	1.12 (0.83, 1.50)	1.25 (0.92, 1.71)	0.75 (0.23, 2.39)
Energetic	1.93 (1.52, 2.44) [†]	1.30 (0.88, 1.91)	1.58 (0.95, 2.64)
Irritability	1.42 (0.82, 2.44)	0.70 (0.42, 1.14)	0.99 (0.22, 4.46)
Loss of interest	2.25 (1.67, 3.03) [†]	2.09 (1.32, 3.32) [†]	NA
Loss of concentration	1.15 (0.86, 1.54)	1.98 (1.40, 2.82) [†]	1.38 (0.84, 2.25)
Change in sleep	1.33 (1.03, 1.73) [*]	1.19 (0.76, 1.87)	1.25 (0.67, 2.35)
Loss of energy	2.24 (1.70, 2.96) [†]	1.94 (1.22, 3.07) [†]	NA
Fatigue	1.65 (1.17, 2.34) [†]	0.98 (0.67, 1.45)	0.30 (0.06, 1.52)
Sadness	1.06 (0.67, 1.66)	1.12 (0.68, 1.84)	0.73 (0.20, 2.68)
Worthless	1.60 (1.03, 2.48) [*]	0.53 (0.18, 1.58)	NA
≥ 1 impaired psychological function	1.48 (1.14, 1.90) [†]	1.35 (1.12, 1.64) [†]	1.82 (1.30, 2.53) [†]
≥ 2 impaired psychological function	1.81 (1.27, 2.57) [†]	1.52 (1.13, 2.05) [†]	1.72 (1.24, 2.38) [†]
≥ 3 impaired psychological function	1.88 (1.21, 2.93) [†]	1.54 (1.03, 2.30) [*]	1.53 (0.73, 3.20)
Sexual function			
Low morning erections	1.33 (1.06, 1.67) [*]	1.09 (0.88, 1.35)	1.63 (0.94, 2.81)
Erectile dysfunction	1.73 (1.50, 1.98) [†]	1.32 (0.99, 1.75)	1.39 (0.74, 2.60)
Low frequency of sexual thoughts	1.06 (0.84, 1.35)	1.04 (0.80, 1.34)	0.70 (0.40, 1.22)
Low sexual desire	1.31 (0.96, 1.79)	1.41 (1.10, 1.81) [†]	1.17 (0.50, 2.71)
Overall sexual satisfaction	1.14 (0.97, 1.33)	1.02 (0.83, 1.27)	0.61 (0.39, 0.96) [*]
≥ 1 impaired sexual function	1.45 (1.14, 1.85) [†]	1.28 (1.12, 1.46) [†]	0.92 (0.57, 1.50)
≥ 2 impaired sexual function	1.48 (1.13, 1.94) [†]	1.37 (1.00, 1.88) [*]	0.94 (0.65, 1.37)
≥ 3 impaired sexual function	1.87 (1.28, 2.72) [†]	1.40 (0.90, 2.19)	1.13 (0.69, 1.86)

* $P < 0.05$, $^{\dagger}P < 0.01$, $^{\ddagger}P < 0.001$. NA, no data in this category.

^aMultivariable logistic regression adjusted for age, smoking status, alcohol intake and centre (reference group: BMI < 30 kg/m² and WC < 102 cm).

Application of other statistical cut-off levels such as tertiles of BMI and WC was examined and did not change their patterns of relationships with symptoms of impaired QoL. ROC analysis showed that the power of prediction by indices of adiposity was similar to previous studies (44). We decided that the results should be presented without adjusting for co-morbidity as related to obesity. Analysis with adjustments for co-morbidities showed no significant change in the relationships

between adiposity and symptoms of ageing, which indicates that the impact of individual co-morbidities is relatively slight. By using the outcome 'at least one symptom of' this analysis has assumed equal importance of all the symptoms. This is unlikely to be true for any individual. It is also possible that certain specific symptoms could dominate the associations described. Men with high indices of adiposity were shown to have exaggerated risk of having composites of two or more

and three or more symptoms of impaired functions. It is of interest that indices of adiposity tend to associate more strongly with the 'physical' items (e.g. tiredness or worn out) of the psychological domain than the 'emotional' items (e.g. loss of interest or change in sleep). The present EMAS data did not show correlations between age and BMI or WC probably because the rates of increases in adiposity are most rapid in younger adults and rather less so up to age 60 years, and also in part, because of reduction in muscle mass from age-related muscle atrophy.

In conclusion, many of the complaints and health problems commonly associated with impaired QoL in older men are accounted for by increased body fat, including those with large waists who are not considered to be 'obese' according to BMI. There is a substantial increase in symptoms as people range from BMI <25 to ≥25–30 categories. WC is a more telling physical measure: lower WC appears to be protective against these health complications even in the presence of high BMI. Health promotion should focus on prevention of excess adult fat accumulation, to reduce burdens of ill health, and weight management should focus on improving the impaired QoL of overweight and obese people.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/EJE-10-1129>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

The European Male Ageing Study (EMAS) is funded by the Commission of the European Communities Fifth Framework Programme 'Quality of Life and Management of Living Resources' Grant QLK6-CT-2001-00258.

Acknowledgements

The authors wish to thank the participants in the eight countries and research/nursing staff in the eight centres: C Pott (Manchester), E Wouters (Leuven), M Nilsson (Malmö), M del Mar Fernandez (Santiago de Compostela), M Jedrzejowska (Lodz), H-M Tabo (Tartu) and A Heredi (Szeged) for their meticulous data collection and C Moseley (Manchester) for data entry and project co-ordination. S Boonen is the senior clinical investigator of the Fund for Scientific research (FWO-Vlaanderen) and holder of the Leuven University Chair in Gerontology and Geriatrics.

EMAS Group

Florence (Gianni Forti, Luisa Petrone, Giovanni Corona); Leuven (Dirk Vanderschueren, Steven Boonen, Herman Borghs); Lodz (Krzysztof Kula, Jolanta Slowikowska-Hilczner, Renata Walczak-Jedrzejowska); London (Ilpo Huhtaniemi); Malmö (Aleksander

Giwerzman); Manchester (Frederick Wu, Alan Silman, Terence O'Neill, Joseph Finn, Philip Steer, Abdelouahid Tajar, David Lee, Stephen Pye); Santiago (Felipe F Casanueva, Mary Lage); Szeged (György Bartfai, Imre Földesi, Imre Fejes); Tartu (Margus Punab, Paul Korrovitz); Turku (Min Jiang).

References

- 1 Visscher TL & Seidell JC. The public health impact of obesity. *Annual Review of Public Health* 2001 **22** 355–375. (doi:10.1146/annurev.publhealth.22.1.355)
- 2 Berghöfer A, Pischon T, Reinhold T, Apovian CM, Sharma AM & Willich SN. Obesity prevalence from a European perspective: a systematic review. *BMC Public Health* 2008 **8** 200. (doi:10.1186/1471-2458-8-200)
- 3 Cameron AJ, Welborn TA, Zimmet PZ, Dunstan DW, Owen N, Salmon J, Dalton M, Jolley D & Shaw JE. Overweight and obesity in Australia: the 1999–2000 Australian diabetes, obesity and lifestyle study (AusDiab). *Medical Journal of Australia* 2003 **178** 427–432.
- 4 Flegal KM, Carroll MD, Ogden CL & Johnson CL. Prevalence and trends in obesity among U.S. adults, 1999–2000. *Journal of the American Medical Association* 2002 **288** 1723–1727. (doi:10.1001/jama.288.14.1723)
- 5 Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ & Flegal KM. Prevalence of overweight and obesity in the United States, 1999–2004. *Journal of the American Medical Association* 2006 **295** 1549–1555. (doi:10.1001/jama.295.13.1549)
- 6 Flegal KM, Carroll MD, Ogden CL & Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *Journal of the American Medical Association* 2010 **303** 235–241. (doi:10.1001/jama.2009.2014)
- 7 Kumar BN, Meyer HE, Wandel M, Dalen I & Holmboe-Ottesen G. Ethnic differences in obesity among immigrants from developing countries, in Oslo, Norway. *International Journal of Obesity* 2006 **30** 684–690. (doi:10.1038/sj.ijo.0803051)
- 8 Wang Y & Beydoun MA. The obesity epidemic in the United States – gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiologic Reviews* 2007 **29** 6–28. (doi:10.1093/epirev/mxm007)
- 9 Ford ES & Mokdad AH. Epidemiology of obesity in the western hemisphere. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** S1–S8. (doi:10.1210/jc.2008-1356)
- 10 Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat. *World Population Prospects: The 2006 Revision and World Urbanization Prospects: The 2005 Revision* (<http://esa.un.org/unpp>).
- 11 Freedman DS, Khan LK, Serdula MK, Galuska DA & Dietz WH. Trends and correlates of class 3 obesity in the united states from 1990 through 2000. *Journal of the American Medical Association* 2002 **288** 1758–1761. (doi:10.1001/jama.288.14.1758)
- 12 Corbett J, Given L, Gray L, Leyland A, MacGregor A, Marryat L, Miller M & Reid S. *The Scottish Health Survey 2008, Volume 1*. Scottish Government: Edinburgh; 2009 (www.scotland.gov.uk/Resource/Doc/286063/0087159.pdf).
- 13 Ebrahimi-Mameghani M, Scott JA, Lean MEJ & Burns CM. Changes in weight and waist circumference over 9 years in a Scottish population. *European Journal of Clinical Nutrition* 2008 **62** 1208–1214. (doi:10.1038/sj.ejcn.1602839)
- 14 Hawn MT, Bian J, Leeth RR, Ritchie G, Allen N, Bland KI & Vickers SM. Impact of obesity on resource utilization for general surgical procedures. *Annals of Surgery* 2005 **241** 821–826. (doi:10.1097/01.sla.0000161044.20857.24)
- 15 Hagenmeyer E-G, von Lengerke T, Gothe H, Schiffhorst G & Happpich M. Impact of concomitant obesity on health care resource consumption in patients with diabetes. Häussler B. *iHEA 2007 6th World Congress: Explorations in Health Economics Paper* 2007.

- 16 Akinnusi ME, Pineda LA & El Solh AA. Effect of obesity on intensive care morbidity and mortality: a meta-analysis. *Critical Care Medicine* 2008 **36** 151–158. (doi:10.1097/01.CCM.0000297885.60037.6E)
- 17 Trogon JG, Finkelstein EA, Hylands T, Dellea PS & Kamal-Bahl SJ. Indirect costs of obesity: a review of the current literature. *Obesity Reviews* 2008 **9** 489–500. (doi:10.1111/j.1467-789X.2008.00472.x)
- 18 The Counterweight Project Team. Impact of obesity on drug prescribing in primary care. *British Journal of General Practice* 2005 **55** 743–749.
- 19 Thorpe KE, Florence CS, Howard DH & Joski P. The impact of obesity on rising medical spending. *Health Affairs* 2004 Suppl Web Exclusives W4-480–W4-486. (10.1377/hlthaff.W4.480)
- 20 Finkelstein EA, Trogon JG, Cohen JW & Dietz W. Annual medical spending attributable to obesity: payer- and service-specific estimates. *Health Affairs* 2009 **28** W822–W831. (doi:10.1377/hlthaff.28.5.w822)
- 21 Lee DM, O'Neill TW, Pye SR, Silman AJ, Finn JD, Pendleton N, Tajar A, Bartfai G, Casanueva F, Forti G, Giwercman A, Huhtaniemi IT, Kula K, Punab M, Boonen S, Vanderschueren D, Wu FC & EMAS study group. The European Male Ageing Study (EMAS): design, methods and recruitment. *International Journal of Andrology* 2009 **32** 11–24. (doi:10.1111/j.1365-2605.2008.00879.x)
- 22 Ware JE, Kosinski M & Keller SD. *SF-36 Physical and Mental Summary Scales: a User's Manual*. Boston, Massachusetts: The Health Institute, New England Medical Center, 1994.
- 23 Beck AT, Steer RA & Brown GK. *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation, 1996.
- 24 O'Connor DB, Corona G, Forti G, Tajar A, Lee DM, Finn JD, Bartfai G, Boonen S, Casanueva FF, Giwercman A, Huhtaniemi IT, Kula K, O'Neill TW, Pendleton N, Punab M, Silman AJ, Vanderschueren D & Wu FC. Assessment of sexual health in aging men in Europe: development and validation of the European Male Ageing Study sexual function questionnaire. *Journal of Sexual Medicine* 2008 **5** 1374–1385. (doi:10.1111/j.1743-6109.2008.00781.x)
- 25 Lean ME, Han TS & Morrison CE. Waist circumference as a measure for indicating need for weight management. *BMJ* 1995 **311** 158–161.
- 26 National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 2002 **106** 3143–3421.
- 27 Larsen SH, Wagner G & Heitmann BL. Sexual function and obesity. *International Journal of Obesity* 2007 **31** 1189–1198. (doi:10.1038/sj.ijo.0803604)
- 28 Bacon CG, Mittleman MA, Kawachi I, Giovannucci E, Glasser DB & Rimm EB. A prospective study of risk factors for erectile dysfunction. *Journal of Urology* 2006 **176** 217–221. (doi:10.1016/S0022-5347(06)00589-1)
- 29 Janiszewski PM, Janssen I & Ross R. Abdominal obesity and physical inactivity are associated with erectile dysfunction independent of body mass index. *Journal of Sexual Medicine* 2009 **6** 1990–1998. (doi:10.1111/j.1743-6109.2009.01302.x)
- 30 Riedner CE, Rhoden EL, Ribeiro EP & Fuchs SC. Central obesity is an independent predictor of erectile dysfunction in older men. *Journal of Urology* 2006 **176** 1519–1523. (doi:10.1016/j.juro.2006.06.049)
- 31 Ford ES, Mokdad AH & Giles WH. Trends in waist circumference among US adults. *Obesity Research* 2003 **11** 1223–1231. (doi:10.1038/oby.2003.168)
- 32 Okosun IS, Choi ST, Boltri JM, Parish DC, Chandra KM, Dever GE & Lucas A. Trends of abdominal adiposity in white, black, and Mexican-American adults, 1988 to 2000. *Obesity Research* 2003 **11** 1010–1017. (doi:10.1038/oby.2003.139)
- 33 Okosun IS, Chandra KM, Boev A, Boltri JM, Choi ST, Parish DC & Dever GE. Abdominal adiposity in U.S. adults: prevalence and trends, 1960–2000. *Preventive Medicine* 2004 **39** 197–206. (doi:10.1016/j.ypmed.2004.01.023)
- 34 Li C, Ford ES, McGuire LC & Mokdad AH. Increasing trends in waist circumference and abdominal obesity among US adults. *Obesity* 2007 **15** 216–224. (doi:10.1038/oby.2007.505)
- 35 Han TS, McNeill G, Seidell JC & Lean ME. Predicting intra-abdominal fatness from anthropometric measures: the influence of stature. *International Journal of Obesity* 1997 **21** 587–593. (doi:10.1038/sj.ijo.0800446)
- 36 Pouliot MC, Després JP, Lemieux S, Moorjani S, Bouchard C, Tremblay A, Nadeau A & Lupien PJ. Waist circumference and abdominal sagittal diameter: best simple anthropometric indexes of abdominal visceral adipose tissue accumulation and related cardiovascular risk in men and women. *American Journal of Cardiology* 1994 **73** 460–468. (doi:10.1016/0002-9149(94)90676-9)
- 37 Han TS, van Leer EM, Seidell JC & Lean ME. Waist circumference action levels in the identification of cardiovascular risk factors: prevalence study in a random sample. *BMJ* 1995 **311** 1401–1405.
- 38 Lean ME, Han TS & Seidell JC. Impairment of health and quality of life in people with large waist circumference. *Lancet* 1998 **351** 853–856. (doi:10.1016/S0140-6736(97)10004-6)
- 39 Seidell JC. Waist circumference and waist/hip ratio in relation to all-cause mortality, cancer and sleep apnea. *European Journal of Clinical Nutrition* 2010 **64** 35–41. (doi:10.1038/ejcn.2009.71)
- 40 Balkau B, Picard P, Vol S, Fezeu L, Eschwege E & DESIR Study Group. Consequences of change in waist circumference on cardiometabolic risk factors over 9 years: data from an Epidemiological Study on the Insulin Resistance Syndrome (DESIR). *Diabetes Care* 2007 **30** 1901–1903. (doi:10.2337/dc06-2542)
- 41 Pischon T, Boeing H, Hoffmann K, Bergmann M, Schulze MB, Overvad K, van der Schouw YT, Spencer E, Moons KG, Tjønneland A, Halkjaer J, Jensen MK, Stegger J, Clavel-Chapelon F, Boutron-Ruault MC, Chajes V, Linseisen J, Kaaks R, Trichopoulos A, Trichopoulos D, Bamia C, Sieri S, Palli D, Tumino R, Vineis P, Panico S, Peeters PH, May AM, Bueno-de-Mesquita HB, van Duynhoven FJ, Hallmans G, Weinhall L, Manjer J, Hedblad B, Lund E, Agudo A, Arriola L, Barricarte A, Navarro C, Martinez C, Quirós JR, Key T, Bingham S, Khaw KT, Boffetta P, Jenab M, Ferrari P & Riboli E. General and abdominal adiposity and risk of death in Europe. *New England Journal of Medicine* 2008 **359** 2105–2120. (doi:10.1056/NEJMoa0801891)
- 42 Zhang C, Rexrode KM, van Dam RM, Li TY & Hu FB. Abdominal obesity and the risk of all-cause, cardiovascular, and cancer mortality: sixteen years of follow-up in US women. *Circulation* 2008 **117** 1658–1667. (doi:10.1161/CIRCULATIONAHA.107.739714)
- 43 WHO Regional Office for Europe. Atlas of health in Europe 2nd edition 2008 (<http://www.euro.who.int/pub>).
- 44 Han TS, van Leer EM, Seidell JC & Lean ME. Waist circumference as a screening tool for cardiovascular risk factors: evaluation of receiver operating characteristics (ROC). *Obesity Research* 1996 **4** 533–547.

Received 28 March 2011

Accepted 1 April 2011