CLINICAL STUDY

Circulating vaspin and visfatin are not affected by acute or chronic energy deficiency or leptin administration in humans

Eun Seok Kang^{1,2}, Faidon Magkos¹, Elizabeth Sienkiewicz¹ and Christos S Mantzoros^{1,3,4}

¹Division of Endocrinology, Diabetes, and Metabolism, Department of Medicine, Harvard Medical School, Beth Israel Deaconess Medical Center, 330 Brookline Avenue, SL Room 412, Boston, Massachusetts 02215, USA, ²Division of Endocrinology and Metabolism, Department of Internal Medicine, Severance Hospital Diabetes Center, Yonsei University College of Medicine, 134 Shinchon Seodaemun, Seoul, South Korea, ³Section of Endocrinology, Boston VA Healthcare System, Harvard Medical School, Boston, Massachusetts 02130, USA and ⁴Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts 02215, USA

(Correspondence should be addressed to E S Kang at Division of Endocrinology, Diabetes, and Metabolism, Department of Medicine, Harvard Medical School, Beth Israel Deaconess Medical Center; Email: ekang@bidmc.harvard.edu)

Abstract

Objective: Animal and *in vitro* studies indicate that leptin alleviates starvation-induced reduction in circulating vaspin and stimulates the production of visfatin. We thus examined whether vaspin and visfatin are affected by short- and long-term energy deprivation and leptin administration in human subjects *in vivo*.

Design and methods: We measured circulating levels of vaspin and visfatin i) before and after 72 h of starvation (leading to severe hypoleptinemia) with or without leptin administration in replacement doses in 13 normal-weight subjects, ii) before and after 72 h of starvation with leptin administration in pharmacological doses in 13 lean and obese subjects, iii) during chronic energy deficiency in eight women with hypothalamic amenorrhea on leptin replacement for 3 months, and iv) during chronic energy deficiency in 18 women with hypothalamic amenorrhea on leptin replacement or placebo for 3 months.

Results: Acute starvation decreased serum leptin to 21% of baseline values, ($P=0.002$) but had no significant effect on vaspin and visfatin concentrations ($P>0.05$). Nor did normalization of leptin levels affect the concentrations of these two adipokines ($P>0.9$). Leptin replacement in women with hypothalamic amenorrhea did not significantly alter vaspin and visfatin concentrations, whether relative to baseline or placebo administration ($P>0.25$). Pharmacological doses of leptin did not affect circulating vaspin and visfatin concentrations ($P>0.9$).

Conclusions: Circulating vaspin and visfatin are not affected by acute or chronic energy deficiency leading to hypoleptinemia and are not regulated by leptin in human subjects, indicating that these adipocyte-secreted hormonal regulators of metabolism are independently regulated in humans.

European Journal of Endocrinology 164 911–917

Introduction

Adipose tissue has been increasingly recognized as an important endocrine organ. Approximately 30 biologically active peptides and proteins (adipokines) are produced and secreted by adipose tissue and play key roles in the regulation of food intake and energy balance, insulin action, and glucose and lipid metabolism (1, 2). Vaspin (visceral adipose tissue-derived serine protease inhibitor) and visfatin (pre-B-cell colony-enhancing factor) are two relatively new and less well-described peptides secreted from white adipose tissue. Studies in animal models indicate that vaspin is an endogenous insulin sensitizer (3) and that visfatin has insulin-mimetic effects (4). Both adipokines are elevated in overweight and obese type 2 diabetic patients (5). Dysregulation of adipose tissue secretory function, such

as that occurring in obesity (excess body fat) (6), has been linked with an array of metabolic complications. The lack of normal adipose tissue, e.g. in lipodystrophy and lipoatrophy, is also associated with dysmetabolic sequelae similar to that seen in obesity (7). Hypoleptinemia is prevalent in energy deprivation states in humans, such as food deprivation/starvation, lipoatrophy, and exercise-induced hypothalamic amenorrhea, and exogenous leptin administration corrects many of the metabolic and neuroendocrine abnormalities characteristic of these conditions (8–12).

It remains unknown whether these adipocyte-secreted hormones are acting independently from each other or whether secretion or administration of one influences the other. In rats, fasting decreases serum vaspin levels that are partly normalized by exogenous leptin administration (13). Leptin in physiological doses

has also been reported to stimulate the production and secretion of visfatin from murine and human omental adipose tissue *ex vivo* and in mouse 3T3-L1 adipocytes *in vitro* (14), but animal and *in vitro* data cannot be directly translated into human biology. There is presently no information on whether leptin, which decreases in energy deprivation states, mediates changes in circulating vaspin and visfatin in humans *in vivo*. Furthermore, nothing is known regarding the *in vivo* regulation of vaspin and visfatin levels in humans by energy deficit *per se*, whether acute or chronic. Thus, in this study, we first assessed circulating vaspin and visfatin in healthy men and women in response to acute energy deficit with and without leptin administration in physiological replacement and pharmacological doses, which would alleviate the significant hypoleptinemia induced by acute energy deficit. We then assessed circulating vaspin and visfatin in women with exercise-induced chronic energy deficit and resultant hypoleptinemia, with or without exogenous leptin given at replacement doses.

Materials and methods

The study protocols were approved by the institutional review board of the Beth Israel Deaconess Medical Center (BIDMC) and recombinant methionyl human leptin (metreleptin; formerly r-metHuLeptin) was administered under an FDA-approved, investigator-initiated IND. The same clinical quality leptin was supplied initially by Amgen, Inc. (Thousand Oaks, CA, USA), and later on by Amylin Pharmaceuticals, Inc. (San Diego, CA, USA). Administration regimens were chosen on the basis of leptin pharmacokinetics (15–17) and modified as per protocol demands. All subjects were informed about the risks associated with the study procedures and gave their written informed consent.

Study 1

Short-term energy deprivation (clinical trial registration: n/a). In study 1, six men (age 23.5 ± 1.5 years) and seven women (age 23.7 ± 1.5 years) with a body mass index (BMI) $< 25 \text{ kg/m}^2$ were studied during three separate admissions in the BIDMC General Clinical Research Center (GCRC). These studies were carried out between 2002 and 2006 and were designed to evaluate the role of leptin in the neuroendocrine and immune response to fasting (18, 19). Each subject completed three 3-day trials in random order: a 72 h control trial (fed state), a 72 h total fasting with administration of placebo, and a 72 h total fasting with administration of replacement doses of metreleptin. Briefly, for each trial, subjects were admitted to the GCRC the night before starting the study. Blood samples were obtained at 0800 h on day 1 and at 0800 h on day 3 for measurement of leptin, vaspin, and visfatin concentrations. Each admission was separated by at least 7

weeks to permit recovery of leptin levels and weight to baseline. Metreleptin was administered at a dose of 0.04 (men) or 0.08 (women) mg/kg per day on the first day and 0.1 (men) or 0.2 (women) mg/kg per day on the second and third days, to account for the progressive decrease in leptin concentration with additional days of fasting (on the basis of our pharmacokinetic studies) (15–17). The total daily metreleptin dose for each 24 h period was divided into four equal doses given every 6 h by s.c. injection (administered by nursing staff) to mimic the normal diurnal variation in leptin levels (17). Placebo was administered according to the same schedule as metreleptin.

Study 2

High-dose leptin treatment (clinical trial registration: clinicaltrials.gov, NCT00140205). The effect of high-dose leptin treatment on vaspin and visfatin was evaluated as part of a pharmacokinetic study of metreleptin carried out between 2006 and 2008 (15). In study 2, five healthy men (age: 22.2 ± 0.9 years) and five healthy women (age: 20.4 ± 0.7 years) with a BMI $< 25 \text{ kg/m}^2$ and five obese but otherwise healthy men (age: 23.4 ± 1.5 years) with a BMI $> 30 \text{ kg/m}^2$ were admitted to the GCRC at the BIDMC for a 3-day fasting study with subcutaneous administration of metreleptin at a pharmacological dose of 0.3 mg/kg, once daily at 0800 h for three consecutive days. Blood samples were obtained at 0800 h on days 1 and 3 for measurement of leptin, vaspin, and visfatin concentrations. Details of this protocol have previously been described (15).

Study 3

Chronic energy deprivation – open label (clinical trial registration: n/a). In study 3, eight women (age 24.8 ± 1.9 years) with relative hypoleptinemia ($< 5 \text{ ng/ml}$) and hypothalamic amenorrhea for at least 6 months due to strenuous exercise or low weight were enrolled and were studied before and after treatment with metreleptin for 3 months. The study was carried out between 2002 and 2004; the clinical characteristics and methods have previously been described in detail (9). Subjects self-administered metreleptin (0.08 mg/kg per day) subcutaneously for 3 months, with 40% of the daily dose given at 0800 h and 60% at 2000 h to mimic the normal diurnal variation in leptin concentrations (17). They were instructed not to change their diet and exercise habits for the duration of the study. Blood samples were obtained monthly for the measurement of leptin, vaspin and visfatin concentrations.

Study 4

Chronic energy deprivation – double blinded (clinical trial registration: clinicaltrials.gov, NCT00130117).

In study 4, 18 women (age 25.0 ± 2.2 years) with hypothalamic amenorrhea for at least 6 months and relative hypoleptinemia (< 5 ng/ml) due to strenuous exercise or low weight were evaluated as part of a larger study on the effects of metreleptin on neuroendocrine function, carried out between 2008 and 2010 (20). Patient characteristics and detailed methods have been described elsewhere (20). The participants were randomized in a 1:1 ratio to receive either metreleptin or placebo. Metreleptin (0.08 mg/kg per day) or matching placebo doses were self-administered by s.c. injection daily between 1900 and 2300 h for 3 months. According to our previous studies (15–17), the dose of metreleptin was calculated based on each subject's weight to achieve physiological leptin concentrations in blood, but given as a single injection to facilitate compliance. Blood samples for the measurement of leptin, vaspin, and visfatin concentrations were obtained at baseline and after 3 months of treatment.

Hormone measurements

All samples used were stored at -80°C until assayed. Leptin concentration was measured by RIA (Millipore, Billerica, MA, USA) with a sensitivity of 0.5 ng/ml. Vaspin concentration was measured using a commercially available ELISA kit (Alpco Diagnostics, Salem, NH, USA) with a sensitivity of 12 pg/ml. Visfatin concentration was measured using a commercially available EIA kit (Alpco Diagnostics) with a sensitivity of 30 pg/ml. All samples were run in duplicate with quality controls, and inter- and intra-assay coefficients of variation for all analytes were $< 15\%$. Routine evaluation in our laboratory, as well as previously published data (21–24), suggest that long-term storage and freezing/thawing does not affect the levels of the molecules of interest.

Statistical analysis

The results are presented as mean $\pm$ S.E.M. Statistical analyses were performed by using SPSS 11.0 (SPSS, Chicago, IL, USA), and a two-tailed $P < 0.05$ was considered statistically significant. For study 1 (short-term energy deficit), paired t -tests were used to assess changes in hormone levels within each condition. To determine whether these changes varied between

conditions, we used a mixed-effects model repeated measures ANOVA. For study 2 (high-dose leptin treatment), paired t -tests were used to assess changes in hormone levels. For studies 3 and 4 (the two chronic energy-deficit studies), changes in hormone levels were analyzed by using a mixed-effects model repeated-measures ANOVA. For comparing changes between the metreleptin- and placebo-treated groups, we used repeated measures analysis as a primary analysis to evaluate changes in hormone levels by overall P values for the main effect of treatment and treatment by time interaction. One-way ANOVA followed by the protected least significant differences technique was used to compare the differences between the two treatment groups for baseline characteristics and for follow-up measurements. Statistical analyses were conducted using on-treatment data.

Results

Study 1: short-term energy deprivation

Leptin concentration was significantly reduced after 3 days of fasting (from 8.5 ± 2.3 to 1.8 ± 0.4 ng/ml, $P = 0.002$) and metreleptin administration reversed this change (both $P = 0.002$); vaspin and visfatin concentrations were not affected by short-term fasting nor leptin replacement (Table 1).

Study 2: high-dose leptin treatment

Pharmacological doses of leptin during 3 days of fasting had no significant effects on serum vaspin (0.68 ± 0.05 vs 0.68 ± 0.05 ng/ml on days 1 and 3 respectively; $P = 0.981$) and visfatin (4.36 ± 0.29 vs 4.31 ± 0.35 ng/ml on days 1 and 3 respectively; $P = 0.907$) concentrations.

Study 3: chronic energy deprivation: open label leptin treatment

Open-label metreleptin treatment for 3 months in women with hypothalamic amenorrhea significantly decreased body weight (54.7 ± 4.5 to 52.2 ± 3.5 kg; $P < 0.001$) and increased serum leptin concentrations to physiological levels during the first 2 months and to

Table 1 Effects of total fasting for 72 h with or without leptin replacement on circulating vaspin and visfatin. Values are means $\pm$ S.E.M.

	Fed		Fasting		Fasting + leptin	
	Day 1	Day 3	Day 1	Day 3	Day 1	Day 3
Leptin (ng/ml)	7.2 ± 1.6	10.1 ± 2.3	8.5 ± 2.3	$1.8 \pm 0.4^*$	7.0 ± 1.7	$15.9 \pm 3.8^*$
Vaspin (ng/ml)	0.68 ± 0.21	0.69 ± 0.18	0.61 ± 0.14	0.40 ± 0.16	0.47 ± 0.15	0.46 ± 0.15
Visfatin (ng/ml)	3.32 ± 0.82	3.66 ± 1.41	4.00 ± 1.27	2.56 ± 0.94	2.68 ± 0.67	2.78 ± 0.79

*Value is significantly different from corresponding value on day 1, $P < 0.05$.

Table 2 Effects of open-label leptin replacement for 3 months on circulating vaspin and visfatin in women with hypothalamic amenorrhea. Values are means $\pm$ S.E.M.

	Baseline	One month	Two months	Three months	P value
Leptin (ng/ml)	3.9 $\pm$ 0.8	10.0 $\pm$ 1.8*	22.1 $\pm$ 7.3*	39.1 $\pm$ 13.1*	<0.001
Vaspin (ng/ml)	1.07 $\pm$ 0.42	1.23 $\pm$ 0.39	1.34 $\pm$ 0.42	1.06 $\pm$ 0.33	0.520
Visfatin (ng/ml)	3.01 $\pm$ 1.16	2.28 $\pm$ 1.02	2.12 $\pm$ 0.60	1.54 $\pm$ 0.61	0.265

The P value (effect of time) from the repeated measures ANOVA is shown. *Value is significantly different from corresponding value at baseline, $P < 0.05$.

mildly supraphysiological levels during the third month at the higher dose (Table 2). There were no significant changes in circulating vaspin or visfatin (Table 2).

Study 4: chronic energy deprivation: double-blind, placebo-controlled, randomized leptin treatment

In the double-blind, placebo-controlled, randomized study, metreleptin treatment decreased body weight only slightly but significantly (55.6 ± 2.0 to 55.5 ± 1.7 kg; $P = 0.046$), whereas no change was observed with placebo administration (55.4 ± 2.3 to 56.9 ± 1.8 ; $P = 0.727$). Despite the significant increase in serum leptin concentration, there were no significant changes in circulating vaspin or visfatin (Table 3).

Discussion

We found that neither short-term energy deprivation (72 h of total fasting) nor chronic energy deficit (hypothalamic amenorrhea) affect serum concentrations of vaspin and visfatin. As energy deprivation states are typically associated with hypoleptinemia, we also studied whether metreleptin administration in either replacement doses to normalize circulating leptin levels or pharmacological doses that would increase serum leptin levels at or beyond the physiological range would alter the concentrations of these two adipokines. We observed that metreleptin has no significant effect on circulating vaspin and visfatin levels. We thus conclude that i) vaspin and visfatin are not likely involved in the development of metabolic and neuroendocrine changes that are typical in states of energy deficiency in humans, and ii) leptin does not modulate the levels of these two adipokines.

Leptin is a pleiotropic hormone; in addition to regulating energy homeostasis, it has important roles in neuroendocrine, reproductive and immune functions (8, 25, 26). Leptin conveys information to the central nervous system (CN) regarding energy availability in the body, a system that is particularly sensitive to energy deprivation. In both humans and animals, states of acute and chronic energy deprivation are characterized by hypoleptinemia and metabolic and neuroendocrine dysfunction, which normalizes with the administration of leptin (9, 27, 28). Leptin exerts its actions by binding to its receptors in both the CN and the periphery and activating signal transduction pathways that overlap with but are distinct from insulin signaling pathways (29). As vaspin and visfatin are novel adipokines exhibiting insulin sensitizing and/or mimetic effects (3, 4), we examined whether leptin regulates the circulating levels of these adipokines by conducting interventional studies in humans involving acute and chronic energy deficiency and leptin administration.

Vaspin is a recently discovered adipocytokine (3). The administration of recombinant vaspin in mice was shown to improve glucose tolerance and insulin sensitivity, identifying this novel adipokine as an insulin sensitizer (3). However, recent cross-sectional studies in humans do not confirm the association between circulating vaspin and glucose tolerance status or insulin sensitivity measured with the hyperinsulinemic clamp technique (30). Still, serum vaspin concentrations are elevated in obesity and type 2 diabetes, and this adipokine has thus been suggested to serve as a biomarker of obesity-related impairment in insulin sensitivity (5, 31, 32). Recent studies in humans have shown significant and direct associations between leptin and vaspin concentrations (23, 33). Also, a study in rats observed that fasting for 24 or 48 h decreases circulating vaspin and this effect is partly abolished by exogenous leptin administration (13). Contrary to these

Table 3 Effects of double-blinded placebo or leptin replacement for 3 months on circulating vaspin and visfatin in women with hypothalamic amenorrhea. Values are means $\pm$ S.E.M.

	Placebo		Leptin		P values		
	Baseline	Three months	Baseline	Three months	Treatment	Time	Interaction
Leptin (ng/ml)	2.77 $\pm$ 0.60	3.24 $\pm$ 0.30	2.67 $\pm$ 0.56	23.88 $\pm$ 5.06*	0.003	0.001	0.002
Vaspin (ng/ml)	0.94 $\pm$ 0.08	1.01 $\pm$ 0.18	0.88 $\pm$ 0.31	1.02 $\pm$ 0.28	0.952	0.278	0.715
Visfatin (ng/ml)	5.80 $\pm$ 0.92	6.35 $\pm$ 0.82	4.22 $\pm$ 0.69	6.22 $\pm$ 0.43	0.921	0.824	0.361

P values from the repeated measures ANOVA are shown. *Value is significantly different from corresponding value at baseline, $P < 0.05$.

findings, a recent observational study in a small number of human subjects reported a preprandial rise and postprandial fall in vaspin levels (34), indicating that either feeding or feeding-related factors such as insulin, which also regulates leptin levels (35) and/or leptin, may be reciprocally related to these daily changes in vaspin levels. To test and expand the hypothesis raised by the above observational study in a systematic and methodical manner, we performed interventional studies involving short- and long-term energy deficit with or without leptin replacement in physiological and pharmacological doses. We found that neither total fasting for 72 h nor leptin administration in replacement or pharmacological doses significantly affected vaspin concentrations in healthy lean and obese men and women. Similarly, we observed no effect of chronic energy deficiency *per se* or leptin replacement in a condition of chronic energy deficit (hypothalamic amenorrhea). The inconsistency of the results between animals and humans could be related to differences in the relative contribution of visceral fat (secreting vaspin) to total body fat. Alternatively, it is also possible that the cellular regulation of vaspin production and secretion in rodents is not similar to that in humans. The fact we did not detect any changes in vaspin levels over a few days of fasting and/or in response to leptin replacement may indicate that only short-term signals of food intake such as insulin may have an effect in altering vaspin levels in the short term, as previously suggested (34).

Visfatin was originally identified as an adipokine exhibiting insulin-mimetic effects via binding to and activating the insulin receptor (4). The results of observational studies on the association between circulating visfatin concentrations and obesity and insulin resistance are conflicting (4, 5, 36–40). In an observational study of obese human subjects, plasma visfatin levels increased following intestinal bypass surgery, and post-surgery visfatin levels were significantly correlated with post-surgery leptin levels and change in leptin levels following surgery (41). In addition, *in vitro* and *ex vivo* studies have shown that leptin increases visfatin production in 3T3-L1 adipocytes and adipose tissue extracts from humans and mice (14). In our short-term interventional studies, however, we found no effect of energy deficit on circulating visfatin levels with or without exogenous metreleptin in replacement or pharmacological doses. In our model of chronic energy deprivation (exercise-induced energy deficiency and hypothalamic amenorrhea), visfatin levels also remained unchanged, even with the normalization of leptin levels through administration of metreleptin for up to 3 months. These results are consistent with the absence of a significant change in circulating visfatin following a long-term, calorie-restricted diet in women (42), as well as the absence of a significant difference in serum visfatin levels

between untreated patients with anorexia nervosa and bulimia nervosa and healthy control subjects (43).

Although our studies were performed in relatively small sample sizes and thus the possibility of a type II error cannot be fully excluded, a recently published study that demonstrated effects of short-term calorie intake and/or food deprivation on vaspin levels was even smaller (34). Moreover, the various approaches taken, the convergence of similar results from several independent studies and especially the randomized, double-blinded, placebo-controlled design of both the short-term and the chronic energy deprivation studies add to the robustness of our results. In addition, we assessed circulating vaspin and visfatin in models of both acute and chronic energy deficit and hypoleptinemia, with administration of physiological and pharmacological doses of metreleptin. The range of leptin concentrations achieved and the breadth of our analysis of these hormones strengthens the validity of our results. Leptin, visfatin, and vaspin were measured using established, state-of-the art methodology with low laboratory error by technicians that were blinded to the study hypothesis. However, in this study, we only assessed a potential effect of leptin but have not studied any potential effects of insulin, as previously suggested (34). This remains to be tested in the future.

In summary, we have shown that i) short- and long-term energy deficit in humans does not affect circulating vaspin and visfatin, indicating that these adipokines are likely not involved in the metabolic and neuroendocrine abnormalities characteristic of energy deprivation states, and ii) leptin administration to alleviate hypoleptinemia does not significantly alter vaspin or visfatin levels. Thus, our results argue against the existence of a cross talk between these adipokines and leptin, which is not only of physiological significance but also may have implications on whether these hormones could be administered and/or co-administered, if and when they find their way in our therapeutic armamentarium.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by Grant Number UL1 RR025758 and M01-RR-01032- Harvard Clinical and Translational Science Center, from the National Center for Research Resources. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Research Resources or the National Institutes of Health. Funding was received from the National Institute of Diabetes and Digestive and Kidney Diseases grants DK58785, DK79929, and DK81913, AG032030; a faculty research grant from Yonsei University College of Medicine (6-2011-0000); and the National Research Foundation of Korea (Grant funded by the Korean Government MEST, Basic Research Promotion Fund, NRF-2010-013-E0008). The Mantzoros Laboratory is also supported by a discretionary grant from Beth Israel Deaconess Medical Center.

Amgen, Inc., and Amylin Pharmaceuticals, Inc., supplied metreleptin for this study and approved the design of the study but had no role in the study design; conduct of the study; collection, management, analysis, and interpretation of the data; or the preparation, review, or approval of the manuscript.

Acknowledgments

We would like to thank the nurses, technicians, and nutritionists at the Beth Israel Deaconess Medical Center General Clinical Research Center in the conduct of the study and John Chamberland for technical assistance.

References

- Athyros VG, Tziomalos K, Karagiannis A, Anagnostis P & Mikhailidis DP. Should adipokines be considered in the choice of the treatment of obesity-related health problems? *Current Drug Targets* 2010 **11** 122–135. (doi:10.2174/138945010790030992)
- Wozniak SE, Gee LL, Wachtel MS & Frezza EE. Adipose tissue: the new endocrine organ? A review article *Digestive Diseases and Sciences* 2009 **54** 1847–1856. (doi:10.1007/s10620-008-0585-3)
- Hida K, Wada J, Eguchi J, Zhang H, Baba M, Seida A, Hashimoto I, Okada T, Yasuhara A, Nakatsuka A, Shikata K, Hourai S, Futami J, Watanabe E, Matsuki Y, Hiramatsu R, Akagi S, Makino H & Kanwar YS. Visceral adipose tissue-derived serine protease inhibitor: a unique insulin-sensitizing adipocytokine in obesity. *PNAS* 2005 **102** 10610–10615. (doi:10.1073/pnas.0504703102)
- Fukuhara A, Matsuda M, Nishizawa M, Segawa K, Tanaka M, Kishimoto K, Matsuki Y, Murakami M, Ichisaka T, Murakami H, Watanabe E, Takagi T, Akiyoshi M, Ohtsubo T, Kihara S, Yamashita S, Makishima M, Funahashi T, Yamanaka S, Hiramatsu R, Matsuzawa Y & Shimomura I. Visfatin: a protein secreted by visceral fat that mimics the effects of insulin. *Science* 2005 **307** 426–430. (doi:10.1126/science.1097243)
- El-Mesallamy HO, Kassem DH, El-Demerdash E & Amin AI. Vaspin and visfatin/Nampt are interesting interrelated adipokines playing a role in the pathogenesis of type 2 diabetes mellitus. *Metabolism* 2011 **60** 63–70. (doi:10.1016/j.metabol.2010.04.008)
- Rasouli N & Kern PA. Adipocytokines and the metabolic complications of obesity. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** S64–S73. (doi:10.1210/jc.2008-1613)
- Rajala MW & Scherer PE. The adipocyte. At the crossroads of energy homeostasis, inflammation, and atherosclerosis. *Endocrinology* 2003 **144** 3765–3773. (doi:10.1210/en.2003-0580)
- Dardeno TA, Chou SH, Moon HS, Chamberland JP, Fiorenza CG & Mantzoros CS. Leptin in human physiology and therapeutics. *Frontiers in Neuroendocrinology* 2010 **31** 377–393. (doi:10.1016/j.yfrne.2010.06.002)
- Welt CK, Chan JL, Bullen J, Murphy R, Smith P, DePaoli AM, Karalis A & Mantzoros CS. Recombinant human leptin in women with hypothalamic amenorrhea. *New England Journal of Medicine* 2004 **351** 987–997. (doi:10.1056/NEJMoa040388)
- Lee JH, Chan JL, Sourlas E, Raptopoulos V & Mantzoros CS. Recombinant methionyl human leptin therapy in replacement doses improves insulin resistance and metabolic profile in patients with lipotrophy and metabolic syndrome induced by the highly active antiretroviral therapy. *Journal of Clinical Endocrinology and Metabolism* 2006 **91** 2605–2611. (doi:10.1210/jc.2005-1545)
- Oral EA, Simha V, Ruiz E, Andewelt A, Premkumar A, Snell P, Wagner AJ, DePaoli AM, Reitman ML, Taylor SI, Gordon P & Garg A. Leptin-replacement therapy for lipodystrophy. *New England Journal of Medicine* 2002 **346** 570–578. (doi:10.1056/NEJMoa012437)
- Rosenbaum M, Nicolson M, Hirsch J, Murphy E, Chu F & Leibel RL. Effects of weight change on plasma leptin concentrations and energy expenditure. *Journal of Clinical Endocrinology and Metabolism* 1997 **82** 3647–3654. (doi:10.1210/jc.82.11.3647)
- Gonzalez CR, Caminos JE, Vazquez MJ, Garces MF, Cepeda LA, Angel A, Gonzalez AC, Garcia-Rendueles ME, Sangiao-Alvarellos S, Lopez M, Bravo SB, Nogueiras R & Dieguez C. Regulation of visceral adipose tissue-derived serine protease inhibitor by nutritional status, metformin, gender and pituitary factors in rat white adipose tissue. *Journal of Physiology* 2009 **587** 3741–3750. (doi:10.1113/jphysiol.2009.172510)
- Tan BK, Chen J, Brown J, Adya R, Ramanjaneya M, Menon V, Bailey CJ, Lehnert H & Randeva HS. *In vivo* and *ex vivo* regulation of visfatin production by leptin in human and murine adipose tissue: role of mitogen-activated protein kinase and phosphatidylinositol 3-kinase signaling pathways. *Endocrinology* 2009 **150** 3530–3539. (doi:10.1210/en.2008-1655)
- Chan JL, Wong SL & Mantzoros CS. Pharmacokinetics of subcutaneous recombinant methionyl human leptin administration in healthy subjects in the fed and fasting states: regulation by gender and adiposity. *Clinical Pharmacokinetics* 2008 **47** 753–764. (doi:10.2165/00003088-200847110-00006)
- Chan JL, Wong SL, Orlova C, Raciti P & Mantzoros CS. Pharmacokinetics of recombinant methionyl human leptin after subcutaneous administration: variation of concentration-dependent parameters according to assay. *Journal of Clinical Endocrinology and Metabolism* 2007 **92** 2307–2311. (doi:10.1210/jc.2006-2864)
- Wong SL, DePaoli AM, Lee JH & Mantzoros CS. Leptin hormonal kinetics in the fed state: effects of adiposity, age, and gender on endogenous leptin production and clearance rates. *Journal of Clinical Endocrinology and Metabolism* 2004 **89** 2672–2677. (doi:10.1210/jc.2003-031931)
- Chan JL, Heist K, DePaoli AM, Veldhuis JD & Mantzoros CS. The role of falling leptin levels in the neuroendocrine and metabolic adaptation to short-term starvation in healthy men. *Journal of Clinical Investigation* 2003 **111** 1409–1421. (doi:10.1172/JCI200317490)
- Chan JL, Matarese G, Shetty GK, Raciti P, Kelesidis I, Aufiero D, De Rosa V, Perna F, Fontana S & Mantzoros CS. Differential regulation of metabolic, neuroendocrine, and immune function by leptin in humans. *PNAS* 2006 **103** 8481–8486. (doi:10.1073/pnas.0505429103)
- Chou SH, Chamberland JP, Liu X, Matarese G, Gao C, Stefanakis R, Brinkoetter MT, Gong H, Arampatzis K & Mantzoros CS. Leptin is an effective treatment for hypothalamic amenorrhea: a randomized, double-blinded, placebo-controlled trial. *PNAS* 2011 In press. (doi:10.1073/pnas.1015674108)
- Botella-Carretero JL, Luque-Ramirez M, Alvarez-Blasco F, Peromingo R, San Millan JL & Escobar-Morreale HF. The increase in serum visfatin after bariatric surgery in morbidly obese women is modulated by weight loss, waist circumference, and presence or absence of diabetes before surgery. *Obesity Surgery* 2008 **18** 1000–1006. (doi:10.1007/s11695-007-9369-7)
- Krzyzanowska K, Mittermayer F, Krugluger W, Kopp HP & Scherthaner G. Increase in visfatin after weight loss induced by gastropastic surgery. *Obesity* 2006 **14** 1886–1889. (doi:10.1038/oby.2006.219)
- Handisurya A, Riedl M, Vila G, Maier C, Clodi M, Prikoszovich T, Ludvik B, Prager G, Luger A & Kautzky-Willer A. Serum vaspin concentrations in relation to insulin sensitivity following RYGB-induced weight loss. *Obesity Surgery* 2010 **20** 198–203. (doi:10.1007/s11695-009-9882-y)
- Ingelsson E, Larson MG, Fox CS, Yin X, Wang TJ, Lipinska I, Pou KM, Hoffmann U, Benjamin EJ, Keaney JF Jr & Vasan RS. Clinical correlates of circulating visfatin levels in a community-based sample. *Diabetes Care* 2007 **30** 1278–1280. (doi:10.2337/dc06-2353)
- Mantzoros CS. Role of leptin in reproduction. *Annals of the New York Academy of Sciences* 2000 **900** 174–183. (doi:10.1111/j.1749-6632.2000.tb06228.x)

- 26 Rosenbaum M, Sy M, Pavlovich K, Leibel RL & Hirsch J. Leptin reverses weight loss-induced changes in regional neural activity responses to visual food stimuli. *Journal of Clinical Investigation* 2008 **118** 2583–2591. (doi:10.1172/JCI35055)
- 27 Ahima RS, Prabakaran D, Mantzoros C, Qu D, Lowell B, Maratos-Flier E & Flier JS. Role of leptin in the neuroendocrine response to fasting. *Nature* 1996 **382** 250–252. (doi:10.1038/382250a0)
- 28 Rosenbaum M, Goldsmith R, Bloomfield D, Magnano A, Weimer L, Heymsfield S, Gallagher D, Mayer L, Murphy E & Leibel RL. Low-dose leptin reverses skeletal muscle, autonomic, and neuroendocrine adaptations to maintenance of reduced weight. *Journal of Clinical Investigation* 2005 **115** 3579–3586. (doi:10.1172/JCI25977)
- 29 Kim YB, Uotani S, Pierroz DD, Flier JS & Kahn BB. *In vivo* administration of leptin activates signal transduction directly in insulin-sensitive tissues: overlapping but distinct pathways from insulin. *Endocrinology* 2000 **141** 2328–2339. (doi:10.1210/en.141.7.2328)
- 30 von Loeffelholz C, Mohlig M, Arafat AM, Isken F, Spranger J, Mai K, Randeve HS, Pfeiffer AF & Weickert MO. Circulating vaspin is unrelated to insulin sensitivity in a cohort of nondiabetic humans. *European Journal of Endocrinology* 2010 **162** 507–513. (doi:10.1530/EJE-09-0737)
- 31 Li Q, Chen R, Moriya J, Yamakawa J, Sumino H, Kanda T & Takahashi T. A novel adipocytokine, visceral adipose tissue-derived serine protease inhibitor (vaspin), and obesity. *Journal of International Medical Research* 2008 **36** 625–629.
- 32 Youn BS, Kloting N, Kratzsch J, Lee N, Park JW, Song ES, Ruschke K, Oberbach A, Fasshauer M, Stumvoll M & Bluher M. Serum vaspin concentrations in human obesity and type 2 diabetes. *Diabetes* 2008 **57** 372–377. (doi:10.2337/db07-1045)
- 33 Aust G, Richter O, Rohm S, Kerner C, Hauss J, Kloting N, Ruschke K, Kovacs P, Youn BS & Bluher M. Vaspin serum concentrations in patients with carotid stenosis. *Atherosclerosis* 2009 **204** 262–266. (doi:10.1016/j.atherosclerosis.2008.08.028)
- 34 Jeong E, Youn BS, Kim DW, Kim EH, Park JW, Namkoong C, Jeong JY, Yoon SY, Park JY, Lee KU & Kim MS. Circadian rhythm of serum vaspin in healthy male volunteers: relation to meals. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 1869–1875. (doi:10.1210/jc.2009-1088)
- 35 Mantzoros CS, Liolios AD, Tritos NA, Kaklamani VG, Doulgerakis DE, Griveas I, Moses AC & Flier JS. Circulating insulin concentrations, smoking, and alcohol intake are important independent predictors of leptin in young healthy men. *Obesity Research* 1998 **6** 179–186.
- 36 Berndt J, Kloting N, Kralisch S, Kovacs P, Fasshauer M, Schon MR, Stumvoll M & Bluher M. Plasma visfatin concentrations and fat depot-specific mRNA expression in humans. *Diabetes* 2005 **54** 2911–2916. (doi:10.2337/diabetes.54.10.2911)
- 37 Kloting N & Kloting I. Visfatin: gene expression in isolated adipocytes and sequence analysis in obese WOKW rats compared with lean control rats. *Biochemical and Biophysical Research Communications* 2005 **332** 1070–1072. (doi:10.1016/j.bbrc.2005.05.058)
- 38 Li L, Yang G, Li Q, Tang Y, Yang M, Yang H & Li K. Changes and relations of circulating visfatin, apelin, and resistin levels in normal, impaired glucose tolerance, and type 2 diabetic subjects. *Experimental and Clinical Endocrinology & Diabetes* 2006 **114** 544–548. (doi:10.1055/s-2006-948309)
- 39 Oki K, Yamane K, Kamei N, Nojima H & Kohno N. Circulating visfatin level is correlated with inflammation, but not with insulin resistance. *Clinical Endocrinology* 2007 **67** 796–800. (doi:10.1111/j.1365-2265.2007.02966.x)
- 40 Pagano C, Pilon C, Olivieri M, Mason P, Fabris R, Serra R, Milan G, Rossato M, Federspil G & Vettor R. Reduced plasma visfatin/pre-B cell colony-enhancing factor in obesity is not related to insulin resistance in humans. *Journal of Clinical Endocrinology and Metabolism* 2006 **91** 3165–3170. (doi:10.1210/jc.2006-0361)
- 41 Garcia-Fuentes E, Garcia-Almeida JM, Garcia-Arnes J, Garcia-Serrano S, Rivas-Marin J, Gallego-Perales JL, Rojo-Martinez G, Garrido-Sanchez L, Bermudez-Silva FJ, Rodriguez de Fonseca F & Soriguer E. Plasma visfatin concentrations in severely obese subjects are increased after intestinal bypass. *Obesity* 2007 **15** 2391–2395. (doi:10.1038/oby.2007.284)
- 42 Mitterberger MC, Mattesich M, Klaver E, Lechner S, Engelhardt T, Larcher L, Pierer G, Piza-Katzer H & Zwerschke W. Adipokine profile and insulin sensitivity in formerly obese women subjected to bariatric surgery or diet-induced long-term caloric restriction. *Journals of gerontology. Series A, Biological sciences and medical sciences* 2010 **65** 915–923. (doi:10.1093/gerona/gdq107)
- 43 Dostalova I, Sedlackova D, Papezova H, Nedvidkova J & Haluzik M. Serum visfatin levels in patients with anorexia nervosa and bulimia nervosa. *Physiological Research* 2009 **58** 903–907.

Received 8 March 2011

Accepted 21 March 2011