CLINICAL STUDY

High prevalence of *AIP* gene mutations following focused screening in young patients with sporadic pituitary macroadenomas

Maria A Tichomirowa^{1,*}, Anne Barlier^{3,*}, Adrian F Daly^{1,*}, Marie-Lise Jaffrain-Rea^{4,5}, Cristina Ronchi^{6,7}, Maria Yaneva⁸, Jonathan D Urban⁹, Patrick Petrossians¹, Atanaska Elenkova⁸, Antoine Tabarin¹⁰, Rachel Desailoud¹¹, Dominique Maiter¹², Thomas Schürmeyer¹³, Renato Cozzi¹⁴, Marily Theodoropoulou¹⁵, Caroline Sievers¹⁵, Ignacio Bernabeu¹⁶, Luciana A Naves¹⁷, Olivier Chabre¹⁸, Carmen Fajardo Montañana¹⁹, Vaclav Hana²⁰, Georges Halaby²¹, Brigitte Delemer²², José Ignacio Labarta Aizpún²³, Emmanuel Sonnet²⁴, Ángel Ferrandez Longás²³, Marie-Thérèse Hagelstein¹, Philippe Caron²⁵, Günter K Stalla¹⁵, Vincent Bours², Sabina Zacharieva⁸, Anna Spada⁷, Thierry Brue²⁶ and Albert Beckers¹

Departments of ¹Endocrinology and ²Molecular Genetics, Centre Hospitalier Universitaire de Liège, Domaine Universitaire du Sart-Tilman, University of Liège, 4000 Liège, Belgium, ³Laboratory of Biochemistry and Molecular Biology, Centre Hospitalo Universitaire Conception, Centre de Recherche en Neurobiologie Neurophysiologie de Marseille, Unité Mixte de Recherche 6231 Centre National de la Recherche Scientifique, Université de la Méditerranée, 13385 Marseille, France, ⁴Department of Experimental Medicine, University of L'Aquila, L'Aquila, Italy, ⁵Neuromed, IRCCS, 86077 Pozzilli, Italy, ⁶Endocrinology and Diabetology Unit, Department of Medicine, University Hospital Wuerzburg, Oberduerrbacher Str. 6, 97080 Wuerzburg Germany, ⁷Unit of Endocrinology, Fondazione IRCCS Ospedale Maggiore Policlinico Mangiagalli Regina Elena, Milan, Italy, ⁸Clinical Center of Endocrinology and Gerontology, 6 bd Damian Gruev, 1303 Sofia, Bulgaria, ⁹ToxStrategies, Inc., 3420 Executive Center Drive, Suite 114, Austin, Texas 78731, USA, ¹⁰Department of Endocrinology, Hôpital Haut Léveque, CHU de Bordeaux, Avenue de Magellan, 33600 Pessac, France, ¹¹Department of Endocrinology, Diabetology and Nutrition, University Medical Centre, 80054 Amiens cedex 1, France, ¹²Department of Endocrinology, Saint Luc University Hospital, Université Catholique de Louvain, B-1200 Brussels, Belgium, ¹³Department of Endocrinology, Klinikum Mutterhaus, Feldstrasse 16, 54290 Trier, Germany, ¹⁴Division of Endocrinology, Ospedale Niguarda, I-20162 Milan, Italy, ¹⁵Department of Endocrinology, Max Planck Institute of Psychiatry, 80804 Munich, Germany, ¹⁶Endocrine Division, Departamento de Medicina, Complejo Hospitalario Universitario de Santiago, Hospital Clínico Universitario de Santiago, Universidad de Santiago de Compostela, La Coruna, Spain, ¹⁷Division of Endocrinology, University of Brasilia, Brasilia, Brazil, ¹⁸Service d'Endocrinologie, Hôpital Albert Michallon, 38043 Grenoble, France, ¹⁹Department of Endocrinology, Hospital Universitario de la Ribera, 46600 Alzira, Valencia, Spain, ²⁰Third Department of Medicine, First Medical Faculty, Charles University, Prague, Czech Republic, ²¹Department of Endocrinology and Metabolism, Hôtel Dieu Hospital, Beirut, Lebanon, ²²Department of Endocrinology, University Hospital of Reims, Reims, France, ²³Department of Pediatrics, Hospital Infantil Miguel Servet, Paseo Isabel la Católica 3, 50009 Zaragoza, Spain, ²⁴Service d'Endocrinologie, CHU de Brest, Hôpital Cavale Blanche, Brest, France, ²⁵Service d'Endocrinologie, Maladies métaboliques Nutrition, CHU de Toulouse, Hôpital Larrey, Pôle Cardio-Vasculaire et Métabolique, 24 Chemin de Pouvoirville, TSA 30030, FR-31059 Toulouse cedex 9, France and ²⁶Service d'Endocrinologie, Diabète et Maladies Métaboliques, et Centre de Référence des Maladies Rares d'Origine Hypophysaires DEFHY, Hôpital de la Timone, 13385 Marseille, France

(Correspondence should be addressed to A Beckers; Email: albert.beckers@chu.ulg.ac.be)

*(M A Tichomirowa, A Barlier and A F Daly contributed equally to this work)

Abstract

Background: Aryl hydrocarbon receptor interacting protein (*AIP*) mutations (*AIPmut*) cause aggressive pituitary adenomas in young patients, usually in the setting of familial isolated pituitary adenomas. The prevalence of *AIPmut* among sporadic pituitary adenoma patients appears to be low; studies have not addressed prevalence in the most clinically relevant population. Hence, we undertook an international, multicenter, prospective genetic, and clinical analysis at 21 tertiary referral endocrine departments.

Methods: We included 163 sporadic pituitary macroadenoma patients irrespective of clinical phenotype diagnosed at <30 years of age.

Results: Overall, 19/163 (11.7%) patients had germline *AIPmut*; a further nine patients had sequence changes of uncertain significance or polymorphisms. *AIPmut* were identified in 8/39 (20.5%) pediatric patients. Ten *AIPmut* were identified in 11/83 (13.3%) sporadic somatotropinoma patients, in 7/61 (11.5%) prolactinoma patients, and in 1/16 non-functioning pituitary adenoma patients. Large genetic deletions were not seen using multiplex ligation-dependent probe amplification. Familial screening was possible in the relatives of seven patients with *AIPmut* and carriers were found in six of the seven families. In total, pituitary adenomas were diagnosed in 2/21 *AIPmut*-screened carriers; both had asymptomatic microadenomas.

Conclusion: Germline *AIPmut* occur in 11.7% of patients <30 years with sporadic pituitary macroadenomas and in 20.5% of pediatric patients. *AIPmut* mutation testing in this population should be considered in order to optimize clinical genetic investigation and management.

European Journal of Endocrinology 165 509–515

Introduction

Clinically relevant pituitary adenomas have a prevalence of one case per 1064–1239 of the population in Western Europe and an incidence of four cases per 100 000 (1–3). The majority of pituitary adenomas occur sporadically, while up to 5% of all cases occur against an inherited or familial background; more than half are due to multiple endocrine neoplasia type 1 (MEN1) (4). Familial isolated pituitary adenomas (FIPA) account for most of the remaining cases, along with rare conditions such as Carney's complex (CNC) or MEN4 (5, 6). Aryl hydrocarbon receptor interacting protein (*AIP*) mutations (*AIPmut*) occur in 15–30% of FIPA kindreds (7–9). In a recent large international study of *AIPmut*-associated pituitary adenomas, Daly *et al.* (10) observed that *AIPmut* are associated with all types of pituitary adenomas; these tumors are large, occur at a young age, and demonstrate features of aggressiveness and treatment resistance. While somatotropinomas and prolactinomas predominate, other pituitary adenoma subtypes do occur in the setting of *AIPmut*, including Cushing disease and thyrotropinoma, and the silent somatotrope subtype of non-functioning adenomas has also recently been described (10, 11).

AIPmut occur infrequently in unselected populations of patients with sporadic pituitary tumors (0–3.0%), in children with sporadic pituitary adenomas (2.7%) or Cushing disease (1.4%) or in specific sporadic tumor types (e.g. acromegaly patients: 3.2%) (12–19). This suggests that *AIPmut* screening of large, unselected sporadic pituitary adenoma populations is impractical; similarly, limited screening performed only in children could miss cases that develop in patients in their late teens and twenties (10). In the largest study of pituitary adenoma patients with *AIPmut* ($n=96$), the most clearly defined features at diagnosis were macroadenoma (93.3%) and young age at diagnosis (median: 23.0 years) (10). Combining this information, 81.2% of patients were diagnosed with a macroadenoma at <30 years of age. No study to date has used this existing knowledge on patient/tumor characteristics to determine whether focused *AIPmut* screening may be clinically informative. In order to explore the utility of focused genetic screening, we investigated the frequency of *AIPmut* among patients with sporadic pituitary macroadenomas that occurred before the age of 30 years, in whom no other known genetic cause or FIPA was present.

Methods

Patients

This genetic screening study was performed in sporadic pituitary adenoma patients with tumors ≥ 10 mm in maximal diameter on magnetic resonance imaging (MRI) that were diagnosed before the age of 30 years. Patients

had to be free of known diagnosis of genetic causes or familial pituitary adenomas (e.g. MEN1, CNC, MEN4, McCune–Albright syndrome, or FIPA). The study was conducted in centers in Belgium, Brazil, Bulgaria, Czech Republic, France, Germany, Italy, Lebanon, and Spain.

A total of 163 patients provided informed consent in their own local language to undergo *AIP* gene sequencing (72.0% of patients meeting the criteria at the study centers agreed to undergo genetic testing). In cases where an *AIPmut* was found, genetic counseling and testing were offered to family members. After counseling, consenting *AIPmut* carriers underwent MRI and hormonal testing to assess for the presence of pituitary disease. Basic demographic, clinical, and therapeutic response data were collected for patients with *AIPmut*. All subjects gave their written informed consent for themselves or for their minor children, while assent of the child was also obtained if they were between 11 and 17 years (no child was under the age of 11 years at the time of genetic testing) and the study was approved by local ethics committees and the Ethics Committee of the University of Liège.

Genetic analysis

DNA extracted from peripheral blood was analyzed for *AIPmut* by direct sequencing. In patients with normal *AIP* sequencing, multiplex ligation-dependent probe amplification (MLPA) was used to search for extensive deletions, as described previously (12). The control DNA from 200 healthy individuals included 30 healthy controls of North African origin to cover this previously unaddressed population, which was represented in our European sample.

The potential effect of each intronic or silent variation on *AIP* protein was evaluated *in silico* using a battery of different tools. For the *in silico* analyses, a 180–240 base portion of the *AIP* gene sequence surrounding each of the nucleotide variants was evaluated for potential splicing and/or functional effects using the following bioinformatic tools: Human Splicing Finder (<http://www.umd.be/HSF/HSF.html>), FANS (<http://fans.ngc.sinica.edu.tw/fans/input.do>), and FastSNP (<http://fastsnp.ibms.sinica.edu.tw>). These tools generate predictions based on algorithms that compare consensus sequences for certain motifs (e.g. sites for intronic and exonic silencers and enhancers) within sequences of interest. As individual tools provide only an estimation of potential effect, it is generally recommended that more than one tool be used to evaluate genomic variations of interest (20). PolyPhen2 (<http://genetics.bwh.harvard.edu/pph2/index.shtml>) was used to evaluate the potential effects of missense mutations on *AIP* structure. *AIP* sequence variants were compared with human single nucleotide polymorphism (SNP) databases (dbSNP, http://www.ncbi.nlm.nih.gov/SNP/snp_summary.cgi) and also against recently published results in genetically diverse populations (21).

Results

Patient characteristics

A total of 163 patients with sporadic pituitary macroadenomas diagnosed before the age of 30 years were included. Eighty-three (50.9%) patients had somatotropinomas (46 females; mean $\pm$ s.d. age at diagnosis: 23 ± 6.2 years, mean $\pm$ s.d. tumor diameter 29.2 ± 14.7 mm); 61 (37.4%) had prolactinomas (39 males; mean $\pm$ s.d. age at diagnosis: 19.6 ± 4.6 years, mean $\pm$ s.d. tumor diameter 34.5 ± 15.8 mm); 16 (9.8%) had non-functioning pituitary adenomas (NFPA; nine females; mean $\pm$ s.d. age at diagnosis: 22.3 ± 5.1 years, mean $\pm$ s.d. tumor diameter 29.5 ± 16 mm); two (1.2%) had Cushing disease (both female, ages at diagnosis: 30.0 and 12.0 years; tumor diameter 12.0 mm in both); and one patient had a thyrotropinoma (male, 15.0 years at diagnosis, tumor diameter 32 mm). Two patients had concomitant brain tumors, one somatotropinoma patient had a glioblastoma of which he later died, and a prolactinoma patient had a meningioma.

AIP sequencing

AIP sequence variations were observed in 28/163 patients (17.2%), of which 19 (11.7%) would likely result in altered AIP expression or function (Table 1). MLPA did not reveal any large deletions. The c.100–18C>T change in one patient is a previously reported rare SNP (rs117691341).

Ten AIPmut were identified in 11/83 (13.3%) sporadic somatotropinoma patients. Eight patients were males and seven presented with invasive tumors. Six of the ten AIPmut led to premature stop codons (p.Arg22X, p.Glu82fsX7, p.Gly117AlafsX39, p.Gln184X, p.Tyr261X, and p.Arg304X ($n=2$ unrelated patients)). The previously described p.Arg271Trp mutation occurs at an amino acid with an established structure–function relationship (8, 10, 22, 23). Two other missense mutations were observed *in silico* to have a highly probable effect on protein structure (p.Glu84Lys and p.Ala277Pro). *In silico* analyses also strongly supported a pathological role for the synonymous p.Glu197Glu variant, via alternative splicing at a splicing enhancer site within an important structural domain. A c.993+60G>C variant in the 3'-untranslated region (UTR) had no predicted effect on splicing, but *in silico* analyses predicted a new transcription factor-binding site, resulting in a low-to-medium likelihood of a deleterious effect. The c.468+16C>T variant predicted a new intronic branch site and also the generation of a new transcription factor-binding site *in silico*, but of a low-to-moderate likelihood of a deleterious effect. The p.Arg16His variant in one patient has previously been reported (8, 15), but recently, it was noted clearly not to segregate with pituitary adenomas

in a FIPA kindred (24). Other AIP sequence variants had no predicted effect on AIP structure/function (p.Arg128His, c.468+15G>T, and c.993+63C>T).

AIPmut were found in 7/61 (11.5%) sporadic prolactinoma patients, six of whom were males. One frameshift/truncation (p.Gly117AlafsX39) was pathogenic. Based on *in silico* analyses, four missense mutations predicted deleterious effects (high likelihood: p.Arg56Cys, p.Lys58Asn, and p.Tyr268Cys; moderate likelihood: p.Val195Ala). One synonymous change (p.Phe269Phe) was also deleterious as characterized by Igreja *et al.* (9). The p.Arg304Gln AIP variant did not strongly predict a deleterious effect on the protein *in silico*, but previous characterization *in vitro* suggests that it is indeed a mutation (9). The p.Arg16His variant was found in one patient (see above).

One AIPmut (c.88_89delGA and p.Asp30Trpfs14X) was found in 1/16 (6.25%) NFPA patients. No AIP variants were seen in the Cushing disease or thyrotropinoma patients.

Familial screening

Familial screening studies were possible in the relatives of seven patients with AIPmut. Mutation carriers were found in six of the seven families. In the family where no mutation carrier was identified, the asymptomatic mother of a somatotropinoma patient with a p.Gly82fsX7 AIPmut declined genetic analysis. Among a total of 21 AIPmut carriers, MRI revealed undiagnosed pituitary adenomas in two members of one family with a p.Arg304X mutation. In that family, three mutation carriers were identified and the patient's mother and sister had pituitary microadenomas, neither of which was associated with pituitary hormonal abnormalities.

AIPmut in pediatric patients with pituitary adenomas

Considering only pediatric patients (i.e. under the age of 18 years), AIP variations were identified in ten of 39 patients (25.6%) diagnosed before 18 years of age (five somatotropinomas and four prolactinomas). Eight variations were likely to be pathogenic in this pediatric cohort (20.5%), which was a significantly higher rate than in the patients aged ≥ 18 years (11/124; $P < 0.01$).

Therapeutic responses

In terms of responses to therapy, the results obtained are in line with those reported elsewhere in AIPmut-bearing patients with pituitary adenomas (10). In patients with somatotropinomas, 4/11 (36.4%) underwent two or more surgical interventions, while secondary (post-operative) somatostatin analog therapy achieved disease control in 1/9 (11%) treated patients. Tumor size before and after somatostatin analog therapy was

Table 1 Clinical characteristic of patients with different *AIP* gene mutations and variants. Pathological *AIP* gene mutations (*AIPmut*) are shown in bold, while variations of unlikely molecular impact and polymorphisms are in normal font.

Patient number	AIPmut/ variant	AN or ref.*	Sex	At diagnosis		Max. tum. diameter (mm)	ES (Y/N)	Invasion (Y/N)	Evidence of del. effect		Familial study
				Age (y)	Hor. levels (ng/ml)				In silico	In vitro (ref.)	
Acromegaly											
1	p.Arg22X	(12)	M	10	GH 6.9 IGF1 N/A	38	Y	Y	Y	N	One mutation carrier, no pituitary tumor
2	p.Glu82fsX7	FJ514477.1	M	15	GH 48.6 IGF1 1404	60	Y	Y	Y	N	No other mutation carrier ^a
3	p.Glu84Lys	GQ403801.1	F	20	GH 14.9 IGF1 556	11	N	N	Y	N	N
4	p.Gly117Alafs39X c.350delG	GQ847774.1	M	18	GH N/A IGF1 810	17	N	N	Y	N	N
5	p.Gln184X	GQ403802.1	M	21	GH 30.0 IGF1 1231	N/A	Y	N/A	Y	N	N
6	p.Glu197Glu	JN561683	F	23	GH 0.5 IGF1 660	26	Y	Y	Y	N	N
7	p.Tyr261X	GQ403803.1	M	28	GH 77.0 IGF1 2443	38	N	Y	Y	N	N
8	p.Arg271Trp	(8)	M	18	GH 180.0 IGF1 N/A	45	Y	Y	Y	Y (9, 17, 22)	N
9	p.Ala277Pro	(32)	M	12	GH 13.6 IGF1 530	27	Y	Y	Y	N	One mutation carrier, no pituitary tumor
10	p.Arg304X	(7)	F	17	GH 9.0 IGF1 905	N/A	N/A	N/A	Y	Y (17)	N
11	p.Arg304X	(7)	M	25	IGF1 578	40	Y	Y	Y	Y (17)	Three mutation carriers, two non-functioning pituitary lesions
12	p.Arg16His	(8)	F	29	N/A	N/A	N/A	N/A	Y	Y; minor (9)	N
13	p.Arg128His	(32)	M	27	GH 58.5 IGF1 909	20	N	N	N	N	N
14	c.468 + 15C>T	(15)	F	19	GH 54.8 IGF1 535	13	N	N	N	N	Two other carrier, no tumor
15	c.468 + 16G>T	(15)	F	15	N/A	23	Y	Y	Y	N	N
16	c.993 + 60 G>C	(15, 29)	F	34 ^b	N/A	15	Y	Y	Y	N	N
17	c.993 + 60 G>C	(15, 29)	M	22	N/A	N/A	N/A	N/A	Y	N	N
18	c.993 + 63 C>T	(15, 29)	F	27	GH 26.0 IGF 643	25	N	Y	N	N	N
Prolactinoma											
19	p.Arg56Cys	GU969040	M	26	PRL 5242	70	Y	Y	Y	N	N
20	p.Lys58Asn	GQ847773.1	M	20	PRL > 10000	72	Y	Y	Y	N	N
21	p.Gly117Alafs39X c.350delG	GQ847774.1	M	16	PRL 2478	N/A	N/A	N/A	Y	N	N
22	p.Val195Ala	(32)	M	12	PRL 10560	40	Y	Y	Y	N	Five mutation carriers, no pituitary tumor
23	p.Tyr268Cys	GQ412196.1	M	28	PRL 6340	85	Y	Y	Y	N	Two mutation carriers, no pituitary tumor
24	p.Phe269Phe	(9)	M	10	PRL 20 000	N/A	Y	Y	Y	Y (9)	No
25	p.Arg304Gln	(15)	M	15	PRL 257	27	Y	Y	Y	Y; minor (9)	Seven mutation carriers, no pituitary tumor
26	p.Arg16His	(8)	M	20	PRL 3867	45	Y	NA	Y	N	N
27	c.100 – 18C>T	rs117691341	M	15	PRL 459	28	Y	Y	Y	N	N
Non-secreting adenoma											
28	p.Asp307TrpfsX14 c.88_89delGA	GQ847775.1	M	19	Normal	39	Y	Y	N	N	N

N/A, not available; AN, Accession number; Ref., reference; Hor., hormone; tum., tumor; ES, extrasellar; del., deleterious.

*Reference of original mutation description.

^aMother of affected patient declined genetic screening.^bAge at first symptoms is 28 years.

measurable in six cases; in one case, there was 60% tumor shrinkage, and in the remaining cases, there was no size change ($n=4$) or tumor growth ($n=1$). Among patients with prolactinomas, 3/6 of cases that received cabergoline did not achieve disease control with usual clinical doses (3–4 mg/week); there were two cases of resistance to high-dose cabergoline (7 mg/week), and in one of these cases, secondary resistance occurred after initial control. Five prolactinoma patients underwent neurosurgery (three patients had two interventions each; one required four interventions) and radiotherapy was used in three patients. The 19-year-old patient with a non-secreting pituitary adenoma underwent neurosurgery twice and radiotherapy thereafter; the tumor was negative for all pituitary hormones on immunohistochemistry.

Discussion

This is the first study to specifically investigate the prevalence of AIP sequence variations among patients thought to be at the greatest risk, namely those diagnosed with pituitary macroadenomas as children or young adults (<30 years). The finding of AIPmut in 11.7% of the entire cohort and among 20.5% of patients aged <18 years at diagnosis strongly supports the contention that screening for AIPmut among young sporadic patients with macroadenomas may be clinically useful. We noted no large AIP gene deletions using MLPA, which suggests that such changes occur in a minority of cases that are negative for AIPmut on sequencing (9, 12, 25).

These results expand on those obtained in previous studies performed using different criteria. Studies in unselected sporadic pituitary adenoma populations demonstrated a very low rate of AIPmut status (12–14, 16). Other studies undertook screening that was restricted to children/adolescents (18, 26), to those with FIPA (8, 9), to sporadic somatotropinoma patients (7, 14, 15), or were limited to a subset of AIPmut (19, 27). These studies identified many novel AIPmut mutations and, along with more recent work (10), have characterized disease features and responses to therapy.

We identified AIPmut in 13.3% of sporadic somatotropinoma patients in our targeted cohort, including six distinct truncating mutations; an Italian acromegalic patient had a previously reported missense mutation (p.Arg271Trp) (8, 10, 23) that was shown to abolish the interaction of AIP with phosphodiesterase 4A5 (17, 22). Two other missense mutations (p.Glu84Lys and p.Ala277Pro) were predicted *in silico* to have probably damaging effects on AIP protein. Also, a new synonymous p.Glu197Glu mutation was strongly predicted to have a pathological effect via interrupting a splicing enhancer site. Notably, the frequency of AIPmut among prolactinoma patients (11.5%) was similar to that of somatotropinoma patients. AIPmut in prolactinoma patients were detected mostly in males and were

relatively difficult to treat with dopamine agonists, which is in line with other recent work (10). As males with prolactinomas can represent a particularly difficult group to treat (28), it is possible that AIPmut status may partially contribute to this poorer disease control, although specific studies of AIPmut status in large prolactinoma cohorts are lacking. The silent variation, p.Phe269Phe, was previously reported as pathogenic by disrupting splicing enhancer sites, causing a loss of exon 6 in a FIPA family with two somatotropinoma patients (9). One novel truncating AIPmut (p.Gly117AlafsX39) occurred in a 16-year-old patient with a prolactinoma and also in an unrelated 18-year-old unrelated acromegalic patient. The p.Lys58Asn, p.Arg56Cys, p.Tyr268Cys, and p.Val195Ala missense mutations scored as being likely to affect AIP protein on *in silico* analysis. A truncating AIPmut was found in 1/16 NFPA (6.3%) patients and is to our knowledge the first report of a pathogenic variation in sporadic NFPA. The patient was aged 19 years and presented with visual field alteration and panhypopituitarism due to mass effects. A previous study identified two AIP variations in a cohort of 55 NFPA patients, but these were found to be polymorphisms (16). The number of NFPA patients included in our study is small, due to the relative rarity of NFPA in younger individuals. No AIPmut were found in Cushing disease or thyrotropinoma patients, but this is again attributed to the small sample size. Corticotropinomas are mainly microadenomas and recently, sporadic Cushing disease in the young has been shown to be very rarely associated with AIPmut (18). Thyrotropinomas are very rare tumors in the general population, and to date, only one case associated with AIPmut has been reported (10).

Among the AIP variants, a number were non-pathogenic on *in silico* analyses (e.g. c.468+15 C>T and c.993+63 C>T). With others, the evidence is less certain or is contradictory. The p.Arg16His variant occurs in an evolutionary conserved residue and is predicted to have a deleterious effect on AIP protein on *in silico* analysis. In contrast, Igraja *et al.* (9) reported only a modest effect of this variant on AIP function in an *in vitro* assay. Also arguing strongly against it being a mutation is the recent publication from Guaraldi & Salvatori (24) showing that this change did not segregate uniformly with pituitary adenomas in a large FIPA kindred. On balance, the evidence currently suggests that p.Arg16His is most likely a rare polymorphism. A p.Arg304Gln mutation was observed in a 15-year-old boy with a prolactinoma whose family screening revealed another seven mutation carriers. No carrier had evidence of pituitary tumor, but one carrier (his mother) was diagnosed with breast cancer and the other (his maternal aunt) has a history of stomach cancer. This mutation occurs at a CpG island hotspot and was previously reported in two families with GH-secreting adenomas and in two sporadic cases of corticotropinoma and one acromegaly patient (13, 15, 17). The strong

clinical data (i.e. multiple affected subjects) and the experimental data are apparently divergent, with only modest effects being seen on *in vitro* AIP functional assays (9) and low potential for deleterious effects *in silico*. The c.993+60G>C variant in the 3'-UTR, while it predicts a low-to-moderate likelihood of a deleterious effect via a new transcription factor binding site and has been reported in colorectal cancer (29), was also present in normal control subjects and is possibly not a pathological variant. *In vitro* testing of AIP variants of uncertain nature using a more expanded range of assays of AIP function beyond phosphodiesterase 4A5-based constructs may be useful to confirm or rule out the pathogenicity of AIP variants.

As AIPmut are most commonly found either in a FIPA or familial setting (7–10, 18) and as pituitary adenomas can be diagnosed in asymptomatic AIPmut carriers via the screening process (30), we performed screening in families of AIPmut pituitary adenoma patients. The completeness of this particular data set is less than optimal for determining penetrance as family members in only seven of the 18 AIPmut patients consented to genetic screening. Mutation carriers were found in six families while in the seventh family the mother of the patient declined genetic studies. MRI revealed two microadenomas among the 21 (9.5%) carriers, which is in keeping with the relatively low disease penetrance of pituitary adenomas. While the tumors in these two AIPmut carriers are clinically silent at present, longer term follow-up is required similar to recent recommendations on incidentaloma surveillance (31). However, it is too early to determine whether these are AIPmut-related tumors or incidentalomas and a high clinical index of suspicion is warranted. Determining true disease penetrance in the setting of AIPmut requires studying a large number of kindreds to allow for detailed surveillance of a sizeable number of mutation carriers through childhood/adolescence and early adulthood.

In conclusion, screening for germline AIPmut in patients aged <30 years with pituitary macroadenomas was positive in 11.7% of cases and in 20.5% of those aged <18 years. Testing for AIPmut in children and young adults diagnosed with macroadenomas could be a useful addition to screening in FIPA kindreds considering the relatively high incidence of AIPmut in this well-defined and readily encountered sub-population (33).

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This study was supported by the Fonds d'Investissement pour la Recherche Scientifique 2008 (FIRS) du CHU de Liège, University of Liège, Belgium, and by the Oncogenetic Network of the French Ministry of Health, CNRS.

References

- 1 Daly AF, Rixhon M, Adam C, Dempegioti A, Tichomirowa MA & Beckers A. High prevalence of pituitary adenomas: a cross-sectional study in the Province of Liege, Belgium. *Journal of Clinical Endocrinology and Metabolism* 2006 **91** 4769–4775. (doi:10.1210/jc.2006-1668)
- 2 Fernandez A, Karavitaki N & Wass JA. Prevalence of pituitary adenomas: a community-based, cross-sectional study in Banbury (Oxfordshire, UK). *Clinical Endocrinology* 2010 **72** 377–382. (doi:10.1111/j.1365-2265.2009.03667.x)
- 3 Raappana A, Koivukangas J, Ebeling T & Pirlä T. Incidence of pituitary adenomas in Northern Finland in 1992–2007. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 4268–4275. (doi:10.1210/jc.2010-0537)
- 4 Scheithauer BW, Laws ER Jr, Kovacs K, Horvath E, Randall RV & Carney JA. Pituitary adenomas of the multiple endocrine neoplasia type I syndrome. *Seminars in Diagnostic Pathology* 1987 **4** 205–211.
- 5 Daly AF, Jaffrain-Rea ML, Ciccirelli A, Valdes-Socin H, Rohmer V, Tamburrano G, Borson-Chazot C, Estour B, Ciccirelli E, Brue T, Ferolla P, Emy P, Colao A, De Menis E, Lecomte P, Penforis F, Delemer B, Bertherat J, Wémeau JL, De Herder W, Archambeaud F, Stevenaert A, Calender A, Murat A, Cavagnini F & Beckers A. Clinical characterization of familial isolated pituitary adenomas. *Journal of Clinical Endocrinology and Metabolism* 2006 **91** 3316–3323. (doi:10.1210/jc.2005-2671)
- 6 Daly AF, Tichomirowa MA & Beckers A. The epidemiology and genetics of pituitary adenomas. *Best Practice & Research. Clinical Endocrinology & Metabolism* 2009 **23** 543–554. (doi:10.1016/j.beem.2009.05.008)
- 7 Vierimaa O, Georgitsi M, Lehtonen R, Vahteristo P, Kokko A, Raitila A, Tuppurainen K, Ebeling TM, Salmela PI, Paschke R, Gundogdu S, De Menis E, Makinen MJ, Launonen V, Karhu A & Aaltonen LA. Pituitary adenoma predisposition caused by germline mutations in the AIP gene. *Science* 2006 **312** 1228–1230. (doi:10.1126/science.1126100)
- 8 Daly AF, Vanbellinghen JF, Khoo SK, Jaffrain-Rea ML, Naves LA, Guitelman MA, Murat A, Emy P, Gimenez-Roqueplo AP, Tamburrano G, Raverot G, Barlier A, De Herder W, Penforis A, Ciccirelli E, Estour B, Lecomte P, Gatta B, Chabre O, Sabaté MI, Bertagna X, Garcia Basavilbaso N, Stalder G, Colao A, Ferolla P, Wémeau JL, Caron P, Sadoul JL, Oneto A, Archambeaud F, Calender A, Sinilnikova O, Montañana CF, Cavagnini F, Hana V, Solano A, Delettières D, Luccio-Camelo DC, Basso A, Rohmer V, Brue T, Bours V, Teh BT & Beckers A. Aryl hydrocarbon receptor-interacting protein gene mutations in familial isolated pituitary adenomas: analysis in 73 families. *Journal of Clinical Endocrinology and Metabolism* 2007 **92** 1891–1896. (doi:10.1210/jc.2006-2513)
- 9 Igreja S, Chahal HS, King P, Bolger GB, Srirangalingam U, Guasti L, Chapple JP, Trivellin G, Gueorguiev M, Guegan K, Stals K, Khoo B, Kumar AV, Ellard S, Grossman AB, Korbonits M & International FIPA Consortium. Characterization of aryl hydrocarbon receptor interacting protein (AIP) mutations in familial isolated pituitary adenoma families. *Human Mutation* 2010 **31** 950–960. (doi:10.1002/humu.21292)
- 10 Daly AF, Tichomirowa MA, Petrossians P, Heliövaara E, Jaffrain-Rea ML, Barlier A, Naves LA, Ebeling T, Karhu A, Raappana A, Cazabat L, De Menis E, Montañana CF, Raverot G, Weil RJ, Sane T, Maier D, Neggers S, Yaneva M, Tabarin A, Verrua E, Eloranta E, Murat A, Vierimaa O, Salmela PI, Emy P, Toledo RA, Sabaté MI, Villa C, Popelier M, Salvatori R, Jennings J, Ferrandez Longás A, Labarta Aizpún JI, Georgitsi M, Paschke R, Ronchi C, Valimaki M, Saloranta C, De Herder W, Cozzi R, Guitelman M, Magri F, Lagonigro MS, Halaby G, Corman V, Hagelstein MT, Vanbellinghen JF, Barra GB, Gimenez-Roqueplo AP, Cameron FJ, Borson-Chazot F, Holdaway I, Toledo SP, Stalla GK, Spada A, Zacharieva S, Bertherat J, Brue T, Bours V, Chanson P, Aaltonen LA & Beckers A. Clinical characteristics and therapeutic

- responses in patients with germ-line AIP mutations and pituitary adenomas: an International Collaborative Study. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** E373–E383. (doi:10.1210/jc.2009-2556)
- 11 Villa C, Lagonigro MS, Magri F, Koziak M, Jaffrain Rea ML, Brauner R, Bouligand J, Junier MP, Di Rocco F, Sainte-Rose C, Beckers A, Roux FX, Daly AF & Chiovato L. Hyperplasia-adenoma sequence in pituitary tumorigenesis related to aryl hydrocarbon receptor interacting protein (AIP) gene mutation. *Endocrine-Related Cancer* 2011 **18** 347–356. (doi:10.1530/ERC-11-0059)
 - 12 Barlier A, Vanbellinghen JF, Daly AF, Silvy M, Jaffrain-Rea ML, Trouillas J, Tamagno G, Cazabat L, Bours V, Brue T, Enjalbert A & Beckers A. Mutations in the aryl hydrocarbon receptor interacting protein gene are not highly prevalent among subjects with sporadic pituitary adenomas. *Journal of Clinical Endocrinology and Metabolism* 2007 **92** 1952–1955. (doi:10.1210/jc.2006-2702)
 - 13 Georgitsi M, Raitila A, Karhu A, Tuppurainen K, Makinen MJ, Vierimaa O, Paschke R, Saeger W, van der Luijt RB, Sane T, Robledo M, De Menis E, Weil RJ, Wasik A, Zielinski G, Lucewicz O, Lubinski J, Launonen V, Vahteristo P & Aaltonen LA. Molecular diagnosis of pituitary adenoma predisposition caused by aryl hydrocarbon receptor-interacting protein gene mutations. *PNAS* 2007 **104** 4101–4105. (doi:10.1073/pnas.0700004104)
 - 14 Iwata T, Yamada S, Mizusawa N, Golam H, Sano T & Yoshimoto K. The aryl hydrocarbon receptor-interacting protein gene is rarely mutated in sporadic GH-secreting adenomas. *Clinical Endocrinology* 2007 **66** 499–502. (doi:10.1111/j.1365-2265.2007.02758.x)
 - 15 Cazabat L, Libè R, Perlemoine K, René-Corail F, Burnichon N, Gimenez-Roqueplo AP, Dupasquier-Fediaevsky L, Bertagna X, Clauser E, Chanson P, Bertherat J & Raffin-Sanson ML. Germline inactivating mutations of the aryl hydrocarbon receptor-interacting protein gene in a large cohort of sporadic acromegaly: mutations are found in a subset of young patients with macroadenomas. *European Journal of Endocrinology* 2007 **157** 1–8. (doi:10.1530/EJE-07-0181)
 - 16 Buchbinder S, Bierhaus A, Zorn M, Nawroth PP, Humpert P & Schilling T. Aryl hydrocarbon receptor interacting protein gene (AIP) mutations are rare in patients with hormone secreting or non-secreting pituitary adenomas. *Experimental and Clinical Endocrinology & Diabetes* 2008 **116** 625–628. (doi:10.1055/s-2008-1065366)
 - 17 Leontiou CA, Gueorguiev M, van der Spuy J, Quinton R, Lolli F, Hassan S, Chahal HS, Igreja SC, Jordan S, Rowe J, Stolbrink M, Christian HC, Wray J, Bishop-Bailey D, Berney DM, Wass JA, Popovic V, Ribeiro-Oliveira A Jr, Gadelha MR, Monson JP, Akker SA, Davis JR, Clayton RN, Yoshimoto K, Iwata T, Matsuno A, Eguchi K, Musat M, Flanagan D, Peters G, Bolger GB, Chapple JP, Frohman LA, Grossman AB & Korbonits M. The role of the aryl hydrocarbon receptor-interacting protein gene in familial and sporadic pituitary adenomas. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** 2390–2401. (doi:10.1210/jc.2007-2611)
 - 18 Stratakis CA, Tichomirowa MA, Boikos S, Azevedo ME, Lodish M, Martari M, Verma S, Daly AF, Raygada M, Keil ME, Papademetriou J, Drori-Herishanu L, Horvath A, Tsang KM, Nesterova M, Franklin S, Vanbellinghen J-F, Bours V, Salvatori R & Beckers A. The role of germline AIP, MEN1, PRKAR1A, CDKN1B and CDKN2C mutations in children and adolescents with pituitary adenoma. *Clinical Genetics* 2010 **78** 457–463. (doi:10.1111/j.1399-0004.2010.01406.x)
 - 19 DiGiovanni R, Serra S, Ezzat S & Asa SL. AIP Mutations are not identified in patients with sporadic pituitary adenomas. *Endocrine Pathology* 2007 **18** 76–78. (doi:10.1007/s12022-007-0010-z)
 - 20 Coassin S, Brandstätter A & Kronenberg F. Lost in the space of bioinformatic tools: a constantly updated survival guide for genetic epidemiology. The GenEpi Toolbox. *Atherosclerosis* 2010 **209** 321–335. (doi:10.1016/j.atherosclerosis.2009.10.026)
 - 21 Rowlands JC, Urban JD, Wikoff DS & Budinsky RA. An evaluation of single nucleotide polymorphisms in the human aryl hydrocarbon receptor-interacting protein (AIP) gene. *Drug Metabolism and Pharmacokinetics*, 2011 In press. (doi:10.2133/dmpk.DMPK-11-SC-013)
 - 22 Bolger GB, Peden AH, Steele MR, MacKenzie C, McEwan DG, Wallace DA, Huston E, Baillie GS & Houslay MD. Attenuation of the activity of the cAMP-specific phosphodiesterase PDE4A5 by interaction with the immunophilin XAP2. *Journal of Biological Chemistry* 2003 **278** 33351–33363. (doi:10.1074/jbc.M303269200)
 - 23 Jennings JE, Georgitsi M, Holdaway I, Daly AF, Tichomirowa M, Beckers A, Aaltonen LA, Karhu A & Cameron FJ. Aggressive pituitary adenomas occurring in young patients in a large Polynesian kindred with a germline R271W mutation in the AIP gene. *European Journal of Endocrinology* 2009 **161** 799–804. (doi:10.1530/EJE-09-0406)
 - 24 Guaraldi F & Salvatori R. Familial isolated pituitary adenomas: from genetics to therapy. *Clinical and Translational Science* 2011 **4** 55–62. (doi:10.1111/j.1752-8062.2010.00254.x)
 - 25 Georgitsi M, Heliövaara E, Paschke R, Kumar AV, Tischkowitz M, Vierimaa O, Salmela P, Sane T, De Menis E, Cannavò S, Gündogdu S, Lucassen A, Izatt L, Aylwin S, Bano G, Hodgson S, Koch CA, Karhu A & Aaltonen LA. Large genomic deletions in AIP in pituitary adenoma predisposition. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** 4146–4151. (doi:10.1210/jc.2008-1003)
 - 26 Georgitsi M, De Menis E, Cannavò S, Mäkinen MJ, Tuppurainen K, Pauletto P, Curtò L, Weil RJ, Paschke R, Zielinski G, Wasik A, Lubinski J, Vahteristo P, Karhu A & Aaltonen LA. Aryl hydrocarbon receptor interacting protein (AIP) gene mutation analysis in children and adolescents with sporadic pituitary adenomas. *Clinical Endocrinology* 2008 **69** 621–627. (doi:10.1111/j.1365-2265.2008.03266.x)
 - 27 Yu R, Bonert S, Saporta I, Raffel LJ & Melmed S. Aryl hydrocarbon receptor interacting protein variants in sporadic pituitary adenomas. *Journal of Clinical Endocrinology and Metabolism* 2006 **91** 5126–5129. (doi:10.1210/jc.2006-1731)
 - 28 Ciccarelli A, Daly AF & Beckers A. The epidemiology of prolactinomas. *Pituitary* 2005 **8** 3–6. (doi:10.1007/s11102-005-5079-0)
 - 29 Georgitsi M, Karhu A, Winqvist R, Visakorpi T, Waltering K, Vahteristo P, Launonen V & Aaltonen LA. Mutation analysis of aryl hydrocarbon receptor interacting protein (AIP) gene in colorectal, breast, and prostate cancers. *British Journal of Cancer* 2007 **96** 352–356. (doi:10.1038/sj.bjc.6603573)
 - 30 Chahal HS, Stals K, Unterländer M, Balding DJ, Thomas MG, Kumar AV, Besser GM, Atkinson AB, Morrison PJ, Howlett TA, Levy MJ, Orme SM, Akker SA, Abel RL, Grossman AB, Burger J, Ellard S & Korbonits M. AIP mutation in pituitary adenomas in the 18th century and today. *New England Journal of Medicine* 2011 **364** 43–50. (doi:10.1056/NEJMoa1008020)
 - 31 Freda PU, Beckers AM, Katznelson L, Molitch ME, Montori VM, Post KD & Vance ML. Pituitary incidentaloma: an endocrine society clinical practice guideline. *Journal of Clinical Endocrinology and Metabolism* 2011 **96** 894–904. (doi:10.1210/jc.2010-1048)
 - 32 Jaffrain-Rea ML, Angelini M, Gargano D, Tichomirowa MA, Daly AF, Vanbellinghen JF, D'Innocenzo E, Barlier A, Giangaspero F, Esposito V, Ventura L, Arcella A, Theodoropoulou M, Naves LA, Fajardo C, Zacharieva S, Rohrer V, Brue T, Gulino A, Cantore G, Alesse E & Beckers A. Expression of aryl hydrocarbon receptor (AHR) and AHR-interacting protein in pituitary adenomas: pathological and clinical implications. *Endocrine-Related Cancer* 2009 **16** 1029–1043. (doi:10.1677/ERC-09-0094)
 - 33 Jaffrain-Rea ML, Daly AF, Angelini M, Petrossians P, Bours V & Beckers A. Genetic susceptibility in pituitary adenomas: from pathogenesis to clinical implications. *Expert Review of Endocrinology and Metabolism* 2011 **6** 195–214. (doi:10.1586/eem.10.87)

Received 6 April 2011

Revised version received 15 June 2011

Accepted 13 July 2011