ERRATUM

Impact of metformin versus the prandial insulin secretagogue, repaglinide, on fasting and postprandial glucose and lipid responses in non-obese patients with type 2 diabetes

Søren S Lund¹, Lise Tarnow¹, Merete Frandsen¹, Ulla M Smidt¹, Oluf Pedersen^{1,2}, Hans-Henrik Parving^{2,3} and Allan A Vaag^{1,4}

¹Steno Diabetes Center, Niels Steensens Vej 2, 2820 Gentofte, Denmark, ²Faculty of Health Sciences, University of Aarhus, Aarhus, Denmark, ³Department of Medical Endocrinology, Rigshospitalet, University of Copenhagen, 2100 Copenhagen, Denmark and ⁴Department of Endocrinology, University of Lund, S-20502 Malmö, Sweden

(Correspondence should be addressed to S S Lund who is now at Boehringer Ingelheim, Ingelheim, Germany; Email: sslund@dadlnet.dk)

The authors and the journal apologise for an error in this article published in the *European Journal of Endocrinology* 2008 Vol **158** pp 35–46. In the second paragraph under the section 'Statistical analysis', on page 37, the formula for calculating the AUC should be as follows:

$$1.5 \text{ h} \times (0.5 \times \text{concentration}_{t=0 \text{ h}} + \text{concentration}_{t=1.5 \text{ h}} + \text{concentration}_{t=3.0 \text{ h}} + \text{concentration}_{t=4.5 \text{ h}} + 0.5 \times \text{concentration}_{t=6.0 \text{ h}})$$

and not as published. The full sentence should therefore read:

Total area under the curve (AUC) during 0–6 h was estimated as the postprandial summary variable and calculated by the trapezoidal rule in units of concentration $\times$ hours, i.e. $1.5 \text{ h} \times (0.5 \times \text{concentration}_{t=0 \text{ h}} + \text{concentration}_{t=1.5 \text{ h}} + \text{concentration}_{t=3.0 \text{ h}} + \text{concentration}_{t=4.5 \text{ h}} + 0.5 \times \text{concentration}_{t=6.0 \text{ h}})$.

European Journal of Endocrinology **165** 831