CLINICAL STUDY

Body weight loss reverts obesity-associated hypogonadotropic hypogonadism: a systematic review and meta-analysis

Giovanni Corona^{1,2}, Giulia Rastrelli¹, Matteo Monami³, Farid Saad⁴, Michaela Luconi⁵, Marcello Lucchese⁶, Enrico Facchiano⁶, Alessandra Sforza², Gianni Forti⁵, Edoardo Mannucci⁷ and Mario Maggi¹

¹Sexual Medicine Andrology Unit, Department of Experimental, Clinical and Biomedical Sciences, University of Florence, Florence, Italy,

²Endocrinology Unit, Medical Department, Azienda Usl Bologna Maggiore-Bellaria Hospital, Bologna, Italy, ³Diabetes Section Geriatric Unit,

Department of Critical Care, University of Florence, Florence, Italy, ⁴Bayer Schering Pharma AG, Berlin, Germany, ⁵Endocrinology Unit,

Department of Clinical Physiopathology, University of Florence, Florence, Italy and ⁶Bariatric and Metabolic Surgery Unit, University of Florence, Florence, Italy and ⁷Diabetes Agency, Careggi Hospital, Florence, Italy

(Correspondence should be addressed to M Maggi; Email: m.maggi@dfc.unifi.it)

Abstract

Objective: Few randomized clinical studies have evaluated the impact of diet and physical activity on testosterone levels in obese men with conflicting results. Conversely, studies on bariatric surgery in men generally have shown an increase in testosterone levels. The aim of this study is to perform a systematic review and meta-analysis of available trials on the effect of body weight loss on sex hormones levels.

Design: Meta-analysis.

Methods: An extensive Medline search was performed including the following words: 'testosterone', 'diet', 'weight loss', 'bariatric surgery', and 'males'. The search was restricted to data from January 1, 1969 up to August 31, 2012.

Results: Out of 266 retrieved articles, 24 were included in the study. Of the latter, 22 evaluated the effect of diet or bariatric surgery, whereas two compared diet and bariatric surgery. Overall, both a low-calorie diet and bariatric surgery are associated with a significant ($P < 0.0001$) increase in plasma sex hormone-binding globulin-bound and -unbound testosterone levels (total testosterone (TT)), with bariatric surgery being more effective in comparison with the low-calorie diet (TT increase: 8.73 (6.51–10.95) vs 2.87 (1.68–4.07) for bariatric surgery and the low-calorie diet, respectively; both $P < 0.0001$ vs baseline). Androgen rise is greater in those patients who lose more weight as well as in younger, non-diabetic subjects with a greater degree of obesity. Body weight loss is also associated with a decrease in estradiol and an increase in gonadotropins levels. Multiple regression analysis shows that the degree of body weight loss is the best determinant of TT rise ($B = 2.50 \pm 0.98$, $P = 0.029$).

Conclusions: These data show that weight loss is associated with an increase in both bound and unbound testosterone levels. The normalization of sex hormones induced by body weight loss is a possible mechanism contributing to the beneficial effects of surgery in morbid obesity.

European Journal of Endocrinology 168 829–843

Introduction

Obesity is a complex condition with serious social and psychological dimensions, which affects virtually all age and socioeconomic groups and threatens to overwhelm both developed and developing countries. Recently, Finucane *et al.* (1) estimated the trends of BMI for adults aged 20 years and older in 199 countries and territories. Between 1980 and 2008, age-standardized mean BMI for men increased in every region apart from Central Africa and South Asia, with USA and UK being highest. The global change worldwide was 0.4 kg/m² per decade for men. Accordingly, the worldwide, age-standardized prevalence of obesity was 9.8% in men in 2008, which was nearly twice the 1980 prevalence of 4.8% (1).

Contrary to conventional wisdom, the obesity epidemic is not restricted to developed societies. Accordingly, a recent large population-based study involving more than 3400 subjects from eight European centers (EMAS study) showed that about half of subjects were overweight or obese with an even higher prevalence in Eastern transitional countries (2). Excess body weight is a crucial risk factor for mortality and morbidity not only for cardiovascular diseases (CVD) but also for type 2 diabetes mellitus (T2DM), cancer, and musculoskeletal disorders; these complications cause nearly 3 million deaths every year worldwide (3, 4, 5, 6, 7, 8). Despite this evidence, obesity is one of today's most blatantly visible – yet most neglected – public health problems.

Several studies have demonstrated that, in individuals at risk, intensive lifestyle interventions, which include nutritional counseling and physical activity, are able to reduce body weight and insulin resistance, preventing the progression to overt T2DM and CVD (9, 10, 11, 12). However, many observational studies have shown that most of the weight lost with diet and exercise is regained in the long term, at least in the majority of patients (13). Bariatric surgery is another possibility to rapidly reduce body weight in individuals at risk, with beneficial effects in obesity-related metabolic disturbances. A meta-analysis of observational studies of bariatric surgery involving 3188 patients with T2DM has reported that 78% had normalization of blood glucose in the absence of medications (14). However, available randomized trials are still very few and of limited size and duration (15, 16). Furthermore, some recent observational studies have suggested caution about the benefits of bariatric surgery, showing no improvement in mortality when compared with standard care (17) and a high rate of long-term surgical complications especially for laparoscopic gastric banding procedures (18).

It is known that obesity in men is associated with low testosterone and reduced sex hormone-binding globulin (SHBG) levels (19, 20, 21, 22, 23). An obesity-associated decline in SHBG might partially explain the observed fall in testosterone levels (19, 20, 21, 22, 23). However, it is important to note that an increased BMI was associated with a low measured, or calculated, free and bioavailable testosterone (24). Specific pathogenetic mechanisms involved in this phenomenon are complex and not completely understood. Evidence indicates that testosterone deficiency induces increased adiposity while increased adiposity induces hypogonadism (22, 25). Few randomized clinical studies have specifically evaluated the impact of diet and physical activity on testosterone levels in obese men. The results of these studies are essentially conflicting: some of them showed an increase of testosterone (26, 27, 28, 29, 30, 31, 32, 33, 34), others showed no change (35, 36, 37, 38, 39), and one small study has shown even a decrease in testosterone levels (40). Similarly, in the last 10 years, several trials have evaluated the impact of bariatric surgery on testosterone levels in men generally, showing an increase in testosterone levels (41, 42, 43, 44, 45, 46, 47, 48).

The aim of this study was to perform a systematic review and meta-analysis of available trials on the effect of body weight loss on sex hormones levels.

Materials and methods

A meta-analysis was performed including all studies, comparing testosterone levels before and after the diet or bariatric surgery. An extensive Medline, Embase, and Cochrane search was performed including the following

words: 'testosterone', 'diet', 'weight loss', and 'bariatric surgery'. The search, which accrued data from January 1, 1969 up to August 31, 2012, was restricted to English language articles and studies of human participants. Only studies with an observational period of at least 2 weeks were considered.

The identification of relevant abstracts, the selection of studies based on the criteria described above, and the subsequent data extraction were performed independently by two of the authors (G Corona and G Rastrelli), and conflicts resolved by the third investigator (M Monami). The quality of studies was assessed using the Cochrane criteria (49).

Completed but still unpublished trials were identified through a search of the www.clinicaltrials.gov website and the results, when not already available on the website, were obtained through a formal request to the authors.

Statistical analysis

Heterogeneity studies were assessed using the I^2 statistics for total testosterone (TT). Considering that heterogeneity could not be excluded ($I^2 = 51.236\%$), mean differences in TT, SHBG, estradiol (E_2), calculated free testosterone (cFT), LH, and FSH between subjects before and after the diet or bariatric surgery were calculated using both a random- and fixed-effect model. For a more conservative approach, results of random-effect models are presented; fixed-effect model is presented only when it differed from the random-effect model. In addition, in some trials, significance of between-group comparisons (P) was not reported; these studies could not be included in the analysis described above. As a consequence, a sensitivity analysis was performed, evaluating endpoint values of each parameter in different treatment groups, in a non-paired fashion (non-paired analysis). Given that most of the studies that did not describe P values reported a non-significant difference across groups, the mean (paired) analysis, which excludes these data, is likely to overestimate the effect of treatments. On the other hand, the non-paired analysis is a very conservative approach, which could underestimate the treatment effect. Meta-regression analysis was performed to test the effect of age, BMI, diabetes, and delta of weight before or after any body-weight-loss interventions on TT and E_2 levels. In addition, different linear age-adjusted regression analysis models, weighting each study for the number of subjects enrolled, were performed to verify the independent effect of baseline BMI and DM as well as percent reduction in BMI (Δ -BMI) and E_2 (Δ - E_2) on TT after weight loss. A further multivariate model was performed, considering Δ -TT as the dependent variable and baseline TT, baseline BMI, and Δ -BMI as covariates.

All analyses were performed using Comprehensive Meta-analysis Version 2, Biostat (Englewood, NJ, USA).

Multivariate analyses were performed on Statistical Package for the Social Sciences (SPSS) for Windows 17.1.

Results

Out of 266 retrieved articles, 24 were included in the study. Of the included studies (see the trial flow summary in Fig. 1), 22 evaluated the effect of diet or bariatric surgery, whereas two compared diet and bariatric surgery. The characteristics of the trials included in the meta-analysis are summarized in Table 1.

Effect of Diet or bariatric surgery on hormonal modifications

Studies comparing the effect of diet or bariatric surgery on hormonal parameters enrolled 479 patients with a mean follow-up of 38 weeks. In particular, 13 studies evaluated the effect of low-calorie diet, whereas 11 investigated the role of bariatric surgery (two gastroplasty, six biliopancreatic diversion, and three mixed interventions respectively). Information on TT, SHBG, and cFT, according to Vermeulen's formula, was available in 23, 15, and nine studies respectively. In addition, data on E₂, FSH, and LH were available in 14, 10, and 10 studies respectively. The mean percent body weight loss at endpoint was $9.8 \pm 4.5\%$ in the low-calorie diet and $32.0 \pm 5.4\%$ in bariatric surgery.

The Begg adjusted rank correlation test (Kendall's $\tau=0.02$, $P=0.88$), calculated on the basis of TT, suggested no major publication bias.

The results of fixed-effect models were not different from those of random-effect models; therefore, only the latter will be reported. When considering data on TT, both low-calorie diet and bariatric surgery were associated with a significant ($P<0.0001$) increase in plasma TT, with bariatric surgery being more significantly effective in comparison with the low-calorie diet

(Fig. 2A). In particular, malabsorptive surgery (including studies which performed biliopancreatic diversion, see references (35, 43, 44, 46, 50)) determined a more consistent increase in TT when compared with non-malabsorptive surgery (including studies which performed gastroplasty approach, see references (41, 42)) (11.58 (7.12 – 16.03) vs 6.08 (2.38 – 9.78) nmol/l, respectively; both $P<0.0001$). However, the latter was not confirmed after adjusting for percent weight loss of BMI (Δ -BMI).

Meta-regression analysis showed that differences in TT, after any approach for body weight loss, were significantly higher in patients with higher TT at baseline as well as in younger and more obese subjects (Fig. 3A, B and C). Furthermore, a higher Δ -BMI was associated with a higher testosterone increase (Fig. 3D). The prevalence of diabetes at baseline was, conversely, inversely related to the TT increase after body weight loss (Fig. 3E). However, meta-regression analysis did not show any association between the Δ -E₂ and TT modifications after weight loss (data not shown).

In order to evaluate the major determinants of the percent increase in TT after weight loss (Δ -TT), alternative, age-adjusted, stepwise, logistic regression analyses were explored using Δ -TT as the dependent variable and baseline TT, BMI, and DM as well as percent reduction in BMI (Δ -BMI) and E₂ (Δ -E₂) after body weight loss as putative predictors. Baseline TT, BMI, and Δ -BMI as well as the percent increase in SHBG (Δ -SHBG) were all significantly associated with Δ -TT ($B=5.16 \pm 2.13$, $P=0.022$; $B=2.51 \pm 0.84$, $P=0.007$; $B=2.76 \pm 0.71$, $P=0.001$; and $B=0.70 \pm 0.24$, $P=0.013$ respectively), whereas no association between Δ -TT and Δ -E₂ or DM at baseline was observed after adjusting for age (data not shown). When a further model was performed considering Δ -TT as the dependent variable and baseline TT, BMI, and Δ -BMI as covariates, only Δ -BMI retained a significant association with Δ -TT ($B=2.50 \pm 0.98$, $P=0.029$).

Figure 1 Trial flow diagram. Unpublished studies were identified through a search of the www.clinicaltrials.gov website.

Table 1 Moderators and outcome variables in individual studies included in the meta-analysis. All data are reported as mean $\pm$ S.D.

Study	Location	Int.	Type of study	FU (weeks)	Age (years)	BMI (kg/m ²)	DM (%)	TT (nmol/l)	LH (U/l)	FSH (U/l)	SHBG (nmol/l)	cFT (pmol/l)	E ₂ (pg/ml)
Stanik <i>et al.</i> (26)	Sepulveda, CA, USA	LCD	BA	8	—	—	4.2	13.9 $\pm$ 3.4	—	—	—	—	36.0 $\pm$ 14.7
Hoffer <i>et al.</i> (27)	Boston, MA, USA	LCD	BA	4	34.0 $\pm$ 7.0	33.1	0	13.0 $\pm$ 1.0	—	—	—	—	—
Pasquali <i>et al.</i> (28)	Bologna, Italy	LCD	BA	10	35.2 $\pm$ 5.6	43.4 $\pm$ 6.3	22.2	11.9	8.4	8.8	9.0	—	45
Strain <i>et al.</i> (29)	New York, NY, USA	LCD	BA	68	34.0 $\pm$ 11.0	—	0	8.3 $\pm$ 4.0	10.3	6.5 $\pm$ 4.7	9.2 $\pm$ 3.2	—	54.0 $\pm$ 26.0
Leenen <i>et al.</i> (36)	Utrecht, The Netherlands	LCD	BA	13	40.0 $\pm$ 6.0	30.7 $\pm$ 2.2	0	12.7 $\pm$ 3.2	—	—	17.0 $\pm$ 6.0	—	27.7 $\pm$ 7.9
Pritchard <i>et al.</i> (30)	Laval, Canada	LCD	BA	13	21.0 $\pm$ 0.8	26.2 $\pm$ 5.5	0	12.3 $\pm$ 4.1	—	—	—	—	—
Kraemer <i>et al.</i> ^a (37)	Pennsylvania, USA	LCD	RCT	12	39.7 $\pm$ 6.6	33.1	0	15.9 $\pm$ 7.7	—	—	—	—	—
Kraemer <i>et al.</i> ^b (37)	Pennsylvania, USA	LCD	RCT	12	38.0 $\pm$ 8.2	31.3	0	17.4 $\pm$ 4.2	—	—	—	—	—
Kraemer <i>et al.</i> ^c (37)	Pennsylvania, USA	LCD	RCT	12	39.6 $\pm$ 6.6	29.2	0	18.1 $\pm$ 5.2	—	—	—	—	—
Volek <i>et al.</i> (38)	Storrs, CT, USA	LCD	BA	6	36.0 $\pm$ 11.9	—	0	21.5 $\pm$ 7.6	—	—	42.8 $\pm$ 32.0	—	—
Mingrone <i>et al.</i> (70)	Rome, Italy	LCD	RCT	54	—	47.8 $\pm$ 8.8	0	—	—	—	19.7 $\pm$ 11.0	—	—
Kaukua <i>et al.</i> (31)	Helsinki, Finland	LCD	RCT	10	45.9 $\pm$ 9.0	39.3 $\pm$ 3.3	5.3	11.1 $\pm$ 3.4	3.6 $\pm$ 2.2	4.9 $\pm$ 2.8	29.0 $\pm$ 14.0	201.3 $\pm$ 58.9	—
Niskanen <i>et al.</i> (32)	Kuopio, Finland	LCD	BA	9	46.3 $\pm$ 7.5	36.1 $\pm$ 3.8	24.0	11.2 $\pm$ 3.9	—	—	27.6 $\pm$ 11.9	185.0 $\pm$ 66.0	12.1 $\pm$ 1.1
Hufelder <i>et al.</i> (33)	Berlin, Germany	LCD	RCT	52	55.9 $\pm$ 6.0	32.5 $\pm$ 2.4	100	10.4 $\pm$ 0.8	—	—	39.7 $\pm$ 8.0	—	—
Reis <i>et al.</i> (35)	São Paulo, Brazil	LCD	RCT	104	42.2 $\pm$ 10.5	54.0 $\pm$ 6.7	0	11.7 $\pm$ 5.1	4.7 $\pm$ 1.5	3.4 $\pm$ 1.6	—	—	37.2 $\pm$ 11.2
Kho0 <i>et al.</i> (34)	Adelaide, Australia	LCD	CBA	8	49.6 $\pm$ 11.0	34.4 $\pm$ 4.0	38.8	20.2 $\pm$ 6.9	—	—	22.2 $\pm$ 8.7	507.0 $\pm$ 300.6	—
Kho0 <i>et al.</i> (39)	Adelaide, SA, Australia	LCD	RCT	52	62.3 $\pm$ 5.9	35.6 $\pm$ 4.8	100	13.9 $\pm$ 3.3	—	—	30.4 $\pm$ 13.9	296.0 $\pm$ 47.0	—
Kho0 <i>et al.</i> ^d (39)	Adelaide, SA, Australia	LCD	RCT	52	58.1 $\pm$ 11.4	35.1 $\pm$ 4.3	100	11.7 $\pm$ 3.6	—	—	22.5 $\pm$ 9.3	285.0 $\pm$ 87.0	—
Bastounis <i>et al.</i> (41)	Athens, Greece	BS ^e	BA	52	34.7 $\pm$ 7.7	57.1 $\pm$ 7.4	0	12.3 $\pm$ 6.2	2.4 $\pm$ 1.6	2.9 $\pm$ 2.0	20.3 $\pm$ 16.8	—	64.4 $\pm$ 24.9
Mingrone <i>et al.</i> (70)	Rome, Italy	BS ^f	RCT	54	—	48.0 $\pm$ 5.4	0	—	—	—	13.2 $\pm$ 6.5	—	—
Globerman <i>et al.</i> (42)	Tel-Aviv, Israel	BS ^e	BA	46.4	38.2 $\pm$ 10.1	44.3 $\pm$ 6.8	35.3	13.4 $\pm$ 7.2	4.9 $\pm$ 3.2	8.0 $\pm$ 5.6	—	—	11.1 $\pm$ 3.4
Alagna <i>et al.</i> (43)	Sassari, Italy	BS ^f	BA	52	—	47.3	75	9.6 $\pm$ 3.7	2.4 $\pm$ 1.6	2.9 $\pm$ 1.9	—	—	44.0 $\pm$ 29.0
Hammoud <i>et al.</i> (44)	Salt Lake City, UT, USA	BS ^f	CBA	104	48.9 $\pm$ 9.6	46.2 $\pm$ 7.2	—	11.8 $\pm$ 5.9	—	—	244.3 $\pm$ 9.9	—	39.0 $\pm$ 26.7
Omana <i>et al.</i> (45)	St Louis, MO, USA	BS ^g	BA	52	48	48.3	43.2	10.6 $\pm$ —	5.1	13.4	35.9	199.0	34.4
Reis <i>et al.</i> (35)	São Paulo, Brazil	BS ^f	RCT	104	36.7 $\pm$ 11.5	55.7 $\pm$ 7.8	0	11.7 $\pm$ 4.5	4.8 $\pm$ 1.7	4.0 $\pm$ 3.4	—	—	47.8 $\pm$ 15.5
Botella <i>et al.</i> ^h (46)	Madrid, Spain	BS ^f	BA	24	40.0 $\pm$ 10.3	47.1 $\pm$ 6.0	35	10.1 $\pm$ 3.5	3.2 $\pm$ 2.2	4.2 $\pm$ 2.2	18.3 $\pm$ 8.3	258.0 $\pm$ 100.4	37.6 $\pm$ 13.4
Fachiano <i>et al.</i> (47)	Florence, Italy	BS ^g	BA	24	39.2	45.7 $\pm$ 6.3	25	8.6 $\pm$ 3.1	2.8 $\pm$ 1.9	3.7 $\pm$ 2.5	19.6 $\pm$ 7.6	219.0 $\pm$ 62.6	43.8 $\pm$ 13.4
Pellitiero <i>et al.</i> (48)	Barcelona, Spain	BS ^g	BA	52	40.5 $\pm$ 9.9	50.3 $\pm$ 6.1	15.2	8.6 $\pm$ 3.2	—	—	18.3 $\pm$ 11.8	230.2 $\pm$ 82.6	42.3 $\pm$ 9.9
Woodard <i>et al.</i> (50)	Stanford, CA, USA	BS ^f	BA	52	48.1 $\pm$ 10.4	48.2 $\pm$ 12.0	42.0	9.0 $\pm$ 4.0	—	—	—	—	—

Int., intervention; LCD, low-calorie diet; BS, bariatric surgery; FU, follow-up (weeks); DM, diabetes mellitus; TT, total testosterone; E₂, estradiol; SHBG, sex hormone-binding globulin; cFT, calculated free testosterone; BA, controlled cohort before-and-after comparisons in the same group of patients; RCT, randomized controlled trials; CBA, controlled before-and-after study between two or more groups of participants receiving different interventions.

^aLow-calorie diet-only group.

^bLow-calorie diet and aerobic exercise three-times-a-week group.

^cLow-calorie diet and aerobic and strength training three-times-a-week group.

^dLow-energy diet group.

^eGastroplasty approach.

^fBiliopancreatic diversion.

^gMixed interventions.

^hHigh-protein, low-fat diet group.

Figure 2 (Legend continued.)

Figure 2 (Legend continued.)

Figure 2 Weighted differences (with 95% CI) of mean total testosterone (TT), sex hormone-binding globulin (SHBG), and calculated free testosterone (cFT) before and after weight loss. Data are presented as derived from the paired *t*-test (A, B and C) and from non-paired analyses (D, E and F). ^Bariatric surgery group; ^^high-protein, low-fat diet group; *low-calorie diet and aerobic exercise three-times-a-week group; and **low-calorie diet and aerobic and strength training three-times-a-week group. LL, lower limit; UL, upper limit.

Figure 3 Influence of baseline total testosterone (TT; A), age (B), BMI (C), percent BMI loss (D), and diabetes mellitus (DM; E) on TT weighted mean differences before and after body weight loss. I, intercept; S, slope.

Similar to what was observed for TT, SHBG ($P < 0.0001$) and cFT levels significantly increased after body weight loss ($P < 0.0001$; see also Fig. 2B and C). No subgroup analyses for the body-weight-loss reduction approach were performed for SHBG and cFT, due to insufficient available data. All these data were confirmed when non-paired analyses were applied (Fig. 2D, E and F).

When considering data on E_2 , body weight loss was associated with a significant reduction in plasma E_2 (mean changes -8.87 (-15.22 ; -2.52) pmol/l, $P < 0.0001$; Fig. 4A). The data were confirmed when non-paired analysis was applied (Fig. 4B). No subgroup analyses for the body-weight-loss reduction approach and no meta-regression analyses were performed for E_2 , due to insufficient data.

Finally, body weight loss, whatever obtained, was significantly associated with a significant increase in both LH and FSH levels (1.31 (0.80 – 1.82) and 1.79 (1.28 – 2.30) U/l for LH and FSH, respectively; both $P < 0.0001$, see also Fig. 5A and B). No subgroup analyses for the body-weight-loss reduction approach were performed for LH and FSH due to insufficient available data. All these data were confirmed when non-paired analyses were applied (Fig. 5C and D).

Discussion

This meta-analysis shows that body weight loss is associated with a relevant increase in gonadotropins and in bound and unbound testosterone, with a decline in the estrogen level. Androgen rise is greater in those patients who lose more weight. The beneficial effects of body weight loss are more apparent in younger, non-diabetic subjects with a greater degree of obesity; however, this could be due to a greater body weight loss.

It is well known that in morbidly obese men, LH levels and pulse amplitude were attenuated, when compared with normal-weight controls (51, 52). Conversely, no modification of LH pulse frequency has been reported in obese men (51, 52). Based on these and other studies (53, 54), obesity-related male hypogonadism is now considered as a form of hypogonadotropic hypogonadism (see for a review, references (55, 56)). Obesity is characterized by a relative abundance of estrogens since P450 aromatase is highly expressed by fat tissue. The increased amount of estrogen levels might, in turn, play a negative feedback on both the hypothalamus and the pituitary, leading to decreased LH secretion (22, 55). Accordingly, it has been reported that a weekly low dose (2.5 mg) of the aromatase inhibitor letrozole can restore testosterone levels and increase LH levels in severely obese hypogonadal men (57). In line with this view, we now show that body weight loss, obtained either by a lifestyle or bariatric intervention, is associated with a decline in estrogen levels and a rise in gonadotropins and testosterone. However, no effect of ΔE_2 on TT after weight loss was observed in the meta-regression analysis. In addition, when ΔE_2 was introduced in a multiple regression model along with other covariates, Δ -BMI, and not ΔE_2 , resulted as the most important determinant of the testosterone increase. Hence, other fat-associated factors, besides estrogens, should mediate a weight reduction-induced improvement in testosterone levels. It should be recognized, however, that local

Figure 4 Weighted differences (with 95% CI) of mean estradiol (E₂) before and after weight loss. Data are presented as derived from the paired *t*-test (A) and from non-paired analysis (B). *Bariatric surgery group.

Figure 5 (Legend continued.)

Figure 5 Weighted differences (with 95% CI) of mean LH and FSH before and after weight loss. Data are presented as derived from the paired *t*-test (A and B) and from non-paired analysis (C and D). *Bariatric surgery group.

aromatization at the hypothalamus–pituitary unit could explain the effect of aromatase inhibitors on LH and testosterone, rather than circulating E_2 itself and that, therefore, the best predictor of local E_2 concentrations is systemic testosterone concentrations.

A series of adipokines has been proposed as a link between obesity and reproductive axis disorders and among them, the most extensively studied are leptin, ghrelin, and adiponectin (58). However, information on adipokine variation levels after a lifestyle or bariatric intervention is not available.

An alternative possibility is that weight reduction is associated with increased insulin sensitivity, with insulin having a permissive role on GnRH neuron activity (59). Accordingly, patients with T2DM and insulin resistance show a higher prevalence of hypogonadotropic hypogonadism (54, 55, 60), and in a mouse model of central depletion of the insulin receptor, the phenotype recapitulates both the metabolic syndrome (MetS) and hypogonadotropic hypogonadism (61). Interestingly, we now report that the diabetic condition has a negative effect on testosterone rise after the intervention; however, this difference does not retain statistical significance after adjusting for Δ -BMI, suggesting that it could be due to a greater body weight loss in non-diabetic patients.

In line with this data, the Endocrine Society Guidelines (62) and the Third International Consultation on Sexual Medicine (63) emphasized that lifestyle modifications should be strongly encouraged in hypogonadal subjects with obesity, T2DM, and the MetS. This meta-analysis demonstrates that, in fact, body weight loss significantly increases testosterone levels. However, the testosterone rise induced by lifestyle interventions was only modest, probably reflecting the relatively modest results of the targeted diet and physical activity on body weight loss (9.8% with diet vs 32% with surgery in the studies included in this meta-analysis).

Our knowledge of the long-term effects of bariatric surgery is based only on observational, non-randomized studies, often of poor methodological quality. These studies showed a relevant beneficial effect of surgery on body weight loss, survival, and obesity-related comorbidities (64, 65). These effects are evident only in patients with severe forms of morbid obesity (64). This meta-analysis shows that bariatric surgery is associated with a greater increase in testosterone levels than lifestyle interventions, particularly in those with a higher baseline BMI and who also show a greater body weight loss. It is possible that the consistent testosterone rise in patients with higher degrees of obesity might mediate some of the aforementioned positive effects of surgical interventions on health status such as the lower incidence of diabetes, the MetS, CVD, and mortality (66, 67, 68).

Male hypogonadism can be considered as one of the many adverse consequences of overweight and obesity.

Body weight loss and lifestyle interventions should be the first approach offered to obese hypogonadal men. However, their effect on testosterone levels is modest. An interesting alternative is the combination of lifestyle interventions and testosterone supplementation. In a small randomized controlled trial (RCT) involving 16 subjects with newly diagnosed T2DM and the MetS, Heufelder *et al.* (33) demonstrated that the combination of testosterone-replacement therapy (TRT) and lifestyle interventions leads to greater therapeutic improvements in glycemic control and reverses the MetS condition after 52 weeks of treatment in comparison with lifestyle interventions only. Similarly, an 18-week, placebo-controlled RCT in obese patients with obstructive sleep apnea demonstrated that the combination of TRT and lifestyle modifications improved insulin sensitivity, reduced liver fat, and increased muscle mass in comparison with placebo and lifestyle modifications alone (69). Hence, although the combination of TRT and the diet would be expected to further improve weight loss-associated testosterone rise, larger studies are advisable to confirm this hypothesis.

Some limitations should be recognized. First, hormonal assays and reference values were not often comparable. In addition, local aromatization at the hypothalamus–pituitary unit could explain the effect of aromatase inhibitors on LH and testosterone, rather than circulating E_2 itself and, therefore, the best predictor of local E_2 concentrations should be systemic testosterone concentrations. Accordingly, our data suggest that circulating E_2 does not represent the major determinant of testosterone increase after weight loss.

In conclusion, body weight loss induces a consistent increase in TT levels, which is greater in those who lose more body weight. Furthermore, a reduction in body weight is associated with an increase in SHBG, cFT, LH, and FSH and a reduction in E_2 . The normalization of sex hormones induced by body weight loss is a possible mechanism contributing to the beneficial effects of surgery in morbid obesity.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by the Italian Minister of University, Research and Instruction (grant nos 2009WLNXT_002 and RBFR10VJ56_002).

References

- 1 Finucane MM, Stevens GA, Cowan MJ, Danaei G, Lin JK, Paciorek CJ, Singh GM, Gutierrez HR, Lu Y, Bahalim AN *et al.* National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and

- epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* 2011 **377** 557–567. (doi:10.1016/S0140-6736(10)62037-5)
- 2 Corona G, Lee DM, Forti G, O'Connor DB, Maggi M, O'Neill TW, Pendleton N, Bartfai G, Boonen S, Casanueva FF *et al.* Age-related changes in general and sexual health in middle-aged and older men: results from the European Male Ageing Study (EMAS). *Journal of Sexual Medicine* 2010 **1** 1362–1380. (doi:10.1111/j.1743-6109.2009.01601.x)
 - 3 Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S, Murray CJ & the Comparative Risk Assessment Collaborating Group. Selected major risk factors and global and regional burden of disease. *Lancet* 2002 **360** 1347–1360. (doi:10.1016/S0140-6736(02)11403-6)
 - 4 Ni Mhurchu C, Rodgers A, Pan WH, Gu DF & Woodward M. Body mass index and cardiovascular disease in the Asia-Pacific region: an overview of 33 cohorts involving 310 000 participants. *International Journal of Epidemiology* 2004 **33** 751–758. (doi:10.1093/ije/dyh163)
 - 5 Prospective Studies Collaboration. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* 2009 **373** 1083–1096. (doi:10.1016/S0140-6736(09)60318-4)
 - 6 World Health Organization. Global health risks: mortality and burden of disease attributable to selected major risks. Geneva: World Health Organization, 2009.
 - 7 Corona G, Monami M, Boddi V, Balzi D, Melani C, Federico N, Balzi D, Sforza A, Rotella CM, Forti G *et al.* Is obesity a further cardiovascular risk factor in patients with erectile dysfunction? *Journal of Sexual Medicine* 2010 **7** 2538–2546.
 - 8 Corona G, Monami M, Boddi V, Cameron-Smith M, Lotti F, de Vita G, Melani C, Balzi D, Sforza A, Forti G *et al.* Male sexuality and cardiovascular risk. A cohort study in patients with erectile dysfunction. *Journal of Sexual Medicine* 2010 **7** 1918–1927. (doi:10.1111/j.1743-6109.2010.01744.x)
 - 9 Tuomilehto J, Lindström J, Eriksson JG, Valle TT, Hämäläinen H, Ilanne-Parikka P, Keinänen-Kiukaanniemi S, Laakso M, Louheranta A, Rastas M *et al.* Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *New England Journal of Medicine* 2001 **344** 1343–1350. (doi:10.1056/NEJM200105033441801)
 - 10 Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM & Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *New England Journal of Medicine* 2002 **346** 393–403. (doi:10.1056/NEJMoa012512)
 - 11 Yamaoka K & Tango T. Efficacy of lifestyle education to prevent type 2 diabetes: a meta-analysis of randomized controlled trials. *Diabetes Care* 2005 **28** 2780–2786. (doi:10.2337/diacare.28.11.2780)
 - 12 Gillies CL, Abrams KR, Lambert PC, Cooper NJ, Sutton AJ, Hsu RT & Khunti K. Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance: systematic review and meta-analysis. *BMJ* 2007 **334** 29912–29915. (doi:10.1136/bmj.39063.689375.55)
 - 13 Bray GA. Uses and misuses of the new pharmacotherapy of obesity. *Annals of Medicine* 1999 **31** 1–3. (doi:10.3109/07853899909019257)
 - 14 Buchwald H, Estok R, Fährbach K, Banel D, Jensen MD, Pories WJ, Bantle JP & Sledge I. Weight and type 2 diabetes after bariatric surgery: systematic review and meta-analysis. *American Journal of Medicine* 2009 **122** 248–256. (doi:10.1016/j.amjmed.2008.09.041)
 - 15 Mingrone G, Panunzi S, De Gaetano A, Guidone C, Iaconelli A, Leccesi L, Nanni G, Pomp A, Castagneto M, Ghirlanda G *et al.* Bariatric surgery versus conventional medical therapy for type 2 diabetes. *New England Journal of Medicine* 2012 **366** 1577–1585. (doi:10.1056/NEJMoa1200111)
 - 16 Dixon JB, Schachter LM, O'Brien PE, Jones K, Grima M, Lambert G, Brown W, Bailey M & Noughton MT. Surgical vs conventional therapy for weight loss treatment of obstructive sleep apnea: a randomized controlled trial. *Journal of the American Medical Association* 2012 **308** 1142–1149. (doi:10.1001/2012.jama.11580)
 - 17 Maciejewski ML, Livingston EH, Smith VA, Kavee AL, Kahwati LC, Henderson WG & Arterburn DE. Survival among high-risk patients after bariatric surgery. *Journal of the American Medical Association* 2011 **305** 2419–2426. (doi:10.1001/jama.2011.817)
 - 18 Himpens J, Cadière GB, Bazi M, Vouche M, Cadière B & Dapri G. Long-term outcomes of laparoscopic adjustable gastric banding. *Archives of Surgery* 2011 **146** 802–807. (doi:10.1001/archsurg.2011.45)
 - 19 Corona G, Rastrelli G, Forti G & Maggi M. Update in testosterone therapy for men. *Journal of Sexual Medicine* 2011 **8** 639–654. (doi:10.1111/j.1743-6109.2010.02200.x)
 - 20 Corona G, Rastrelli G, Vignozzi L & Maggi M. Emerging medication for the treatment of male hypogonadism. *Expert Opinion on Emerging Drugs* 2012 **17** 239–292. (doi:10.1517/14728214.2012.683411)
 - 21 Corona G, Rastrelli G, Vignozzi L, Mannucci E & Maggi M. How to recognize late-onset hypogonadism in men with sexual dysfunction. *Asian Journal of Andrology* 2012 **14** 251–259. (doi:10.1038/aja.2011.138)
 - 22 Corona G, Rastrelli G, Morelli A, Vignozzi L, Mannucci E & Maggi M. Hypogonadism and metabolic syndrome. *Journal of Endocrinological Investigation* 2011 **34** 557–567.
 - 23 Corona G, Mannucci E, Forti G & Maggi M. Following the common association between testosterone deficiency and diabetes mellitus, can testosterone be regarded as a new therapy for diabetes? *International Journal of Andrology* 2009 **32** 431–441. (doi:10.1111/j.1365-2605.2009.00965.x)
 - 24 Wu FC, Tajar A, Pye SR, Silman AJ, Finn JD, O'Neill TW, Bartfai G, Casanueva F, Forti G, Giwercman A *et al.* Hypothalamic–pituitary–testicular axis disruptions in older men are differentially linked to age and modifiable risk factors: the European Male Aging Study. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** 2737–2745. (doi:10.1210/jc.2007-1972)
 - 25 Saad F, Aversa A, Isidori AM & Gooren LJ. Testosterone as potential effective therapy in treatment of obesity in men with testosterone deficiency: a review. *Current Diabetes Reviews* 2012 **8** 131–143. (doi:10.2174/157339912799424573)
 - 26 Stanik S, Dornfeld LP, Maxwell MH, Viosca SP & Korenman SG. The effect of weight loss on reproductive hormones in obese men. *Journal of Clinical Endocrinology and Metabolism* 1981 **53** 828–832. (doi:10.1210/jcem-53-4-828)
 - 27 Hoffer LJ, Beitins IZ, Kyung NH & Bistrian BR. Effects of severe dietary restriction on male reproductive hormones. *Journal of Clinical Endocrinology and Metabolism* 1986 **62** 288–292. (doi:10.1210/jcem-62-2-288)
 - 28 Pasquali R, Casimirri F, Melchionda N, Fabbri R, Capelli M, Platé L, Patrono D, Balestra V & Barbara L. Weight loss and sex steroid metabolism in massively obese man. *Journal of Endocrinological Investigation* 1988 **11** 205–210.
 - 29 Strain GW, Zumoff B, Miller LK, Rosner W, Levit C, Kalin M, Hershcovf RJ & Rosenfeld RS. Effect of massive weight loss on hypothalamic–pituitary–gonadal function in obese men. *Journal of Clinical Endocrinology and Metabolism* 1988 **66** 1019–1023. (doi:10.1210/jcem-66-5-1019)
 - 30 Pritchard J, Després JP, Gagnon J, Tchernof A, Nadeau A, Tremblay A & Bouchard C. Plasma adrenal, gonadal, and conjugated steroids following long-term exercise-induced negative energy balance in identical twins. *Metabolism* 1999 **48** 1120–1127. (doi:10.1016/S0026-0495(99)90125-7)
 - 31 Kaukua J, Pekkarinen T, Sane T & Mustajoki P. Sex hormones and sexual function in obese men losing weight. *Obesity Research* 2003 **11** 689–694. (doi:10.1038/oby.2003.98)
 - 32 Niskanen L, Laaksonen DE, Punnonen K, Mustajoki P, Kaukua J & Rissanen A. Changes in sex hormone-binding globulin and testosterone during weight loss and weight maintenance in

- abdominally obese men with the metabolic syndrome. *Diabetes, Obesity & Metabolism* 2004 **6** 208–215. (doi:10.1111/j.1462-8902.2004.00335.x)
- 33 Heufelder AE, Saad F, Bunck MC & Gooren L. Fifty-two-week treatment with diet and exercise plus transdermal testosterone reverses the metabolic syndrome and improves glycemic control in men with newly diagnosed type 2 diabetes and subnormal plasma testosterone. *Journal of Andrology* 2009 **30** 726–733. (doi:10.2164/jandrol.108.007005)
 - 34 Khoo J, Piantadosi C, Worthley S & Wittert GA. Effects of a low-energy diet on sexual function and lower urinary tract symptoms in obese men. *International Journal of Obesity* 2010 **34** 1396–1403. (doi:10.1038/ijo.2010.76)
 - 35 Reis LO, Favaro WJ, Barreiro GC, de Oliveira LC, Chaim EA, Fregonesi A & Ferreira U. Erectile dysfunction and hormonal imbalance in morbidly obese male is reversed after gastric bypass surgery: a prospective randomized controlled trial. *International Journal of Andrology* 2010 **33** 736–744. (doi:10.1111/j.1365-2605.2009.01017.x)
 - 36 Leenen R, van der Kooy K, Seidell JC, Deurenberg P & Koppeschaar HP. Visceral fat accumulation in relation to sex hormones in obese men and women undergoing weight loss therapy. *Journal of Clinical Endocrinology and Metabolism* 1994 **78** 1515–1520. (doi:10.1210/jc.78.6.1515)
 - 37 Kraemer WJ, Volek JS, Clark KL, Gordon SE, Puhl SM, Koziris LP, McBride JM, Triplett-McBride NT, Putukian M, Newton RU *et al.* Influence of exercise training on physiological and performance changes with weight loss in men. *Medicine and Science in Sports and Exercise* 1999 **31** 1320–1329. (doi:10.1097/00005768-199909000-00014)
 - 38 Volek JS, Sharman MJ, Love DM, Avery NG, Gómez AL, Scheett TP & Kraemer WJ. Body composition and hormonal responses to a carbohydrate-restricted diet. *Metabolism* 2002 **51** 864–870. (doi:10.1053/meta.2002.32037)
 - 39 Khoo J, Piantadosi C, Duncan R, Worthley SG, Jenkins A, Noakes M, Worthley MI, Lange K & Wittert GA. Comparing effects of a low-energy diet and a high-protein low-fat diet on sexual and endothelial function, urinary tract symptoms, and inflammation in obese diabetic men. *Journal of Sexual Medicine* 2011 **10** 2868–2875. (doi:10.1111/j.1743-6109.2011.02417.x)
 - 40 Klibanski A, Beitins IZ, Badger T, Little R & McArthur JW. Reproductive function during fasting in men. *Journal of Clinical Endocrinology and Metabolism* 1981 **53** 258–263. (doi:10.1210/jcem-53-2-258)
 - 41 Bastounis EA, Karayiannakis AJ, Syrigos K, Zbar A, Makri GG & Alexiou D. Sex hormone changes in morbidly obese patients after vertical banded gastroplasty. *European Surgical Research* 1998 **30** 43–47. (doi:10.1159/00008556)
 - 42 Globerman H, Shen-Orr Z, Karnieli E, Aloni Y & Charuzi I. Inhibin B in men with severe obesity and after weight reduction following gastroplasty. *Endocrine Research* 2005 **31** 17–26. (doi:10.1080/07435800500228971)
 - 43 Alagna S, Cossu ML, Gallo P, Tilocca PL, Pileri P, Alagna G, Maninchedda P, Sini AL, Pilo L, Rovasio PP *et al.* Biliopancreatic diversion: long-term effects on gonadal function in severely obese men. *Surgery for Obesity and Related Diseases* 2006 **2** 82–86. (doi:10.1016/j.soard.2006.01.005)
 - 44 Hammoud A, Gibson M, Hunt SC, Adams TD, Carrell DT, Kolotkin RL & Meikle AW. Effect of Roux-en-Y gastric bypass surgery on the sex steroids and quality of life in obese men. *Journal of Clinical Endocrinology and Metabolism* 2009 **94** 1329–1332. (doi:10.1210/jc.2008-1598)
 - 45 Omana JJ, Tamler R, Strohmayer E, Herron D & Kini S. Sex hormone levels in men undergoing bariatric surgery. *Journal of the American College of Surgeons* 2009 **35** S22–S23. (doi:10.1016/j.jamcollsurg.2009.06.042)
 - 46 Botella-Carretero JJ, Balsa JA, Gómez-Martin JM, Peromingo R, Huerta L, Carrasco M, Arrieta F, Zamarron I, Martín-Hidalgo A & Vazquez C. Circulating free testosterone in obese men after bariatric surgery increases in parallel with insulin sensitivity. *Journal of Endocrinological Investigation* 2013. In press.
 - 47 Facchiano E, Scaringi S, Veltri M, Samavat J, Maggi M, Forti G, Luconi M & Lucchese M. Age as a predictive factor of testosterone improvement in male patients after bariatric surgery: preliminary results of a monocentric prospective study. *Obesity Surgery* 2012 **23** 167–172. (doi:10.1007/s11695-012-0753-6)
 - 48 Pellitero S, Olazola I, Alastrue A, Martínez E, Granada ML, Balibrea JM, Moreno P, Serra A, Navarro-Díaz M, Romero R *et al.* Hypogonadotropic hypogonadism in morbidly obese males is reversed after bariatric surgery. *Obesity Surgery* 2012 **22** 1835–1842.
 - 49 Eds Higgins JPT & Green S. In *Cochrane Handbook for Systematic Reviews of Interventions*, version 5.0.1 (updated September 2008). The Cochrane Collaboration, 2008. (Available from www.cochrane-handbook.org).
 - 50 Woodard G, Ahmed S, Podelski V, Hernandez-Boussard T, Presti J Jr & Morton JM. Effect of Roux-en-Y gastric bypass on testosterone and prostate-specific antigen. *British Journal of Surgery* 2012 **99** 693–698. (doi:10.1002/bjs.8693)
 - 51 Giagulli VA, Kaufman JM & Vermeulen A. Pathogenesis of the decreased androgen levels in obese men. *Journal of Clinical Endocrinology and Metabolism* 1994 **79** 997–1000. (doi:10.1210/jc.79.4.997)
 - 52 Vermeulen A, Kaufman JM, Deslypere JP & Thomas G. Attenuated luteinizing hormone (LH) pulse amplitude but normal LH pulse frequency, and its relation to plasma androgens in hypogonadism of obese men. *Journal of Clinical Endocrinology and Metabolism* 1993 **76** 1140–1146. (doi:10.1210/jc.76.5.1140)
 - 53 Corona G, Mannucci E, Fisher AD, Lotti F, Petrone L, Balercia G, Bandini E, Forti G & Maggi M. Low levels of androgens in men with erectile dysfunction and obesity. *Journal of Sexual Medicine* 2008 **5** 2454–2463.
 - 54 Dhindsa S, Prabhakar S, Sethi M, Bandyopadhyay A, Chaudhuri A & Dandona P. Frequent occurrence of hypogonadotropic hypogonadism in type 2 diabetes. *Journal of Clinical Endocrinology and Metabolism* 2004 **89** 5462–5468. (doi:10.1210/jc.2004-0804)
 - 55 Dandona P & Dhindsa S. Update: hypogonadotropic hypogonadism in type 2 diabetes and obesity. *Journal of Clinical Endocrinology and Metabolism* 2011 **96** 2643–2651. (doi:10.1210/jc.2010-2724)
 - 56 Saad F, Aversa A, Isidori AM, Zafalon L, Zitzmann M & Gooren L. Onset of effects of testosterone treatment and time span until maximum effects are achieved. *European Journal of Endocrinology* 2011 **165** 675–685. (doi:10.1530/EJE-11-0221)
 - 57 Loves S, Ruinemans-Koerts J & de Boer H. Letrozole once a week normalizes serum testosterone in obesity-related male hypogonadism. *European Journal of Endocrinology* 2008 **158** 741–747. (doi:10.1530/EJE-07-0663)
 - 58 Michalakis K, Mintzioti G, Kaprara A, Tarlatzis BC & Goulis DG. The complex interaction between obesity, metabolic syndrome and reproductive axis: a narrative review. *Metabolism* 2013. In press. (doi:10.1016/j.metabol.2012.08.012)
 - 59 Burcelin R, Thorens B, Glauser M, Gaillard RC & Pralong FP. Gonadotropin-releasing hormone secretion from hypothalamic neurons: stimulation by insulin and potentiation by leptin. *Endocrinology* 2003 **144** 4484–4491. (doi:10.1210/en.2003-0457)
 - 60 Corona G, Mannucci M, Petrone L, Ricca V, Balercia G, Mansani R, Chiarini V, Giommi R, Forti G & Maggi M. Association of hypogonadism and type 2 diabetes in men attending an outpatient erectile dysfunction clinic. *International Journal of Impotence Research* 2006 **18** 190–197. (doi:10.1038/sj.ijir.3901391)
 - 61 Brüning JC, Gautam D, Burks DJ, Gillette J, Schubert M, Orban PC, Klein R, Krone W, Müller-Wieland D & Kahn CR. Role of brain insulin receptor in control of body weight and reproduction. *Science* 2000 **289** 2122–2125. (doi:10.1126/science.289.5487.2122)
 - 62 Bhasin S, Cunningham GR, Hayes FJ, Matsumoto AM, Snyder PJ, Swerdloff RS, Montori VM, Task Force & Endocrine Society. Testosterone therapy in men with androgen deficiency

- syndromes: an Endocrine Society Clinical Practice Guideline. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 2536–2559. (doi:10.1210/jc.2009-2354)
- 63 Buvat J, Maggi M, Gooren L, Guay AT, Kaufman J, Morgentaler A, Schulman C, Tan HM, Torres LO, Yassin A *et al.* Endocrine aspects of male sexual dysfunctions. *Journal of Sexual Medicine* 2010 **7** (4 Pt 2) 1627–1656. (doi:10.1111/j.1743-6109.2010.01780.x)
 - 64 Sjöström L, Narbro K, Sjöström CD, Karason K, Larsson B, Wedel H, Lystig T, Sullivan M, Bouchard C, Carlsson B *et al.* Effects of bariatric surgery on mortality in Swedish obese subjects. *New England Journal of Medicine* 2007 **357** 741–752. (doi:10.1056/NEJMoa066254)
 - 65 Sjöström L, Lindroos AK, Peltonen M, Torgerson J, Bouchard C, Carlsson B, Dahlgren S, Larsson B, Narbro K, Sjöström CD *et al.* Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery. *New England Journal of Medicine* 2004 **351** 2683–2693. (doi:10.1056/NEJMoa035622)
 - 66 Corona G, Rastrelli G, Monami M, Guay A, Buvat J, Sforza A, Forti G, Mannucci E & Maggi M. Hypogonadism as a risk factor for cardiovascular mortality in men: a meta-analytic study. *European Journal of Endocrinology* 2011 **165** 687–701. (doi:10.1530/EJE-11-0447)
 - 67 Corona G, Monami M, Rastrelli G, Aversa A, Sforza A, Lenzi A, Forti G, Mannucci E & Maggi M. Type 2 diabetes mellitus and testosterone: a meta-analysis study. *International Journal of Andrology* 2011 **34** (6 Pt 1) 528–540. (doi:10.1111/j.1365-2605.2010.01117.x)
 - 68 Corona G, Monami M, Rastrelli G, Aversa A, Tishova Y, Saad F, Lenzi A, Forti G, Mannucci E & Maggi M. Testosterone and metabolic syndrome: a meta-analysis study. *Journal of Sexual Medicine* 2011 **8** 272–283. (doi:10.1111/j.1743-6109.2010.01991.x)
 - 69 Hoyos CM, Yee BJ, Phillips CL, Machan EA, Grunstein RR & Liu PY. Body compositional and cardiometabolic effects of testosterone therapy in obese men with severe obstructive sleep apnea: a randomised placebo-controlled trial. *European Journal of Endocrinology* 2012 **167** 531–541. (doi:10.1530/EJE-12-0525)
 - 70 Mingrone G, Greco AV, Giancaterini A, Scarfone A, Castagneto M & Pugeat M. Sex hormone-binding globulin levels and cardiovascular risk factors in morbidly obese subjects before and after weight reduction induced by diet or malabsorptive surgery. *Atherosclerosis* 2002 **161** 455–462. (doi:10.1016/S0021-9150(01)00667-0)

Received 30 October 2012

Revised version received 2 March 2013

Accepted 12 March 2013