IGF1 and IGFBP3 in acute respiratory distress syndrome

Amy M Ahasic^{1,2}, Rihong Zhai², Li Su², Yang Zhao², Konstantinos N Aronis³,
B Taylor Thompson⁴, Christos S Mantzoros^{2,3} and David C Christiani^{2,4}

¹Section of Pulmonary and Critical Care Medicine, Department of Medicine, Yale University School of Medicine, 300 Cedar Street, PO Box 208057, New Haven, Connecticut 06520-8057, USA, ²Department of Environmental Health, Harvard School of Public Health, 665 Huntington Avenue, Building 1-1407, Boston, Massachusetts 02115, USA, ³Division of Endocrinology, Diabetes, and Metabolism, Department of Medicine, Harvard Medical School, Beth Israel Deaconess Medical Center, Feldberg 875, 330 Brookline Avenue, Boston, Massachusetts 02215, USA and ⁴Pulmonary and Critical Care Unit, Department of Medicine, Harvard Medical School, Massachusetts General Hospital, Bulfinch 1-148, 20 Fruit Street, Boston, Massachusetts 02114, USA

Correspondence
should be addressed
to A M Ahasic
Email
amy.ahasic@yale.edu

The authors apologise for error in their article titled 'IGF1 and IGFBP3 in acute respiratory distress syndrome' published in the *European Journal of Endocrinology* 2012, Vol 166 Iss 1, pp 121–129. The error pertains to the units used for reporting IGF1 and IGFBP3 levels. This error, in units only, has no impact on the results or the conclusions of the study.

On page 124 of the manuscript, the median IGF1 concentration in the study population should have been reported as 67.1 ng/ml (IQR, 59.6), and median IGFBP3 concentration as 1989.7 ng/ml (IQR, 1616.0).

In Table 2, the IGF1 and IGFBP3 values for controls and cases respectively should have been presented as follows:

Table 2 Baseline characteristics of cohort with plasma available by ARDS status.

Characteristic	Controls (n = 356)	ARDS cases (n = 175)	P value*
IGF1 (ng/ml), median (IQR)	72.6 (65.9)	53.1 (52.2)	<0.0001
IGFBP3 (mg/ml), median (IQR)	2083 (1596)	1707 (1495)	0.0009

IQR, interquartile range. *P value reflects comparison between controls and cases.

In Table 4, the IGF1 and IGFBP3 values for survivors and non-survivors respectively should have read as follows:

Table 4 Baseline cohort characteristics of ARDS cases by 60-day all-cause mortality.

Characteristic	Survivors (n = 97)	Non-survivors (n = 78)	P value*
IGF1 (ng/ml), median (IQR)	60.1 (54.4)	44.3 (39.2)	0.0005
IGFBP3 (mg/ml), median (IQR)	2129 (1465)	1328 (1423)	0.0016

IQR, interquartile range. *P value reflects comparison between survivors and non-survivors.

The authors emphasise that as this is a factor of 10, a constant, it does not affect the results reported in any way.