From first symptoms to final diagnosis of Cushing's disease: experiences of 176 patients

Ilonka Kreitschmann-Andermahr¹, Tsambika Psaras², Maria Tsiogka³, Daniel Starz⁴,
Bernadette Kleist¹, Sonja Siegel^{1,4}, Monika Milian⁵, Johannes Kohlmann⁴,
Christa Menzel⁴, Dagmar Führer-Sakel³, Jürgen Honegger⁵, Ulrich Sure¹,
Oliver Müller¹ and Michael Buchfelder⁴

¹Department of Neurosurgery, University Hospital Essen, University of Essen-Duisburg, Hufelandstrasse 55, 45147 Essen, Germany, ²Department of Neurosurgery, Hospital of Cologne-Merheim, Cologne, Germany, ³Department of Endocrinology, University of Essen-Duisburg, Essen, Germany, ⁴Department of Neurosurgery, University of Erlangen-Nuremberg, Erlangen, Germany and ⁵Department of Neurosurgery, University of Tuebingen, Tuebingen, Germany

Correspondence
should be addressed to
I Kreitschmann-Andermahr
Email
ilonka.kreitschmann@
uk-essen.de

The authors and the journal apologise for an error in Figure 1 in the above article published in the *European Journal of Endocrinology* vol 172, issue 3, pp 285–289. In the figure, the reasons for medical consultation of an internist before the diagnosis of Cushing's disease have been erroneously presented to be the same as to those for consultation of a gynaecologist. The figure legend had also been inadvertently cropped. The correct figure with the legend is published in full below.

Figure 1

Most frequent symptom categories and the medical specialists to whom they were presented. *CD, Cushing's disease. **This

cluster contains symptoms such as buffalo-hump, moon-face, plethora, petechiae, hirsutism, striae and muscle atrophy.