EJE PRIZE 2015

How does insulin resistance arise, and how does it cause disease? Human genetic lessons

R K Semple

University of Cambridge Metabolic Research Laboratories, Wellcome Trust-MRC Institute of Metabolic Science, Level 4, Box 289, Addenbrooke's Treatment Centre, Cambridge CB2 0QQ, UK

Correspondence should be addressed to R K Semple
Email
rks16@cam.ac.uk

Abstract

Insulin orchestrates physiological responses to ingested nutrients; however, although it elicits widely ramifying metabolic and trophic responses from diverse tissues, 'insulin resistance (IR)', a pandemic metabolic derangement commonly associated with obesity, is usually defined solely by blunting of insulin's hypoglycaemic effect. Recent study of monogenic forms of IR has established that biochemical subphenotypes of IR exist, clustering into those caused by primary disorders of adipose tissue and those caused by primary defects in proximal insulin signalling. IR is often first recognised by virtue of its associated disorders including type 2 diabetes, dyslipidaemia (DL), fatty liver and polycystic ovary syndrome (PCOS). Although these clinically observed associations are confirmed by cross-sectional and longitudinal population-based studies, causal relationships among these phenomena have been more difficult to establish. Single gene IR is important to recognise in order to optimise clinical management and also permits testing of causal relationships among components of the IR syndrome using the principle of Mendelian randomisation. Thus, where a precisely defined genetic defect is identified that directly produces one component of the syndrome, then phenomena that are causally linked to that component should be seen. Where this is not the case, then a simple causal link is refuted. This article summarises known forms of monogenic severe IR and considers the lessons to be learned about the pathogenic mechanisms both upstream from common IR and those downstream linking it to disorders such as DL, fatty liver, PCOS and cancer.

European Journal of Endocrinology
(2016) **174**, R209–R223

Origins of the concept of insulin resistance

The discovery and clinical introduction of purified insulin in the early 1920s transformed diabetes mellitus from a rapidly fatal childhood malady into a manageable chronic disease. However, as the effort to translate therapeutic proof of principle into cheap, widely available insulin

treatment gathered momentum in ensuing years, it became apparent that a subgroup of people with diabetes did not exhibit the dramatic response of the emaciated, ketotic patients first treated. Most prominent in studying and articulating this was Harold Himsworth who, in the late 1930s, discriminated between 'sensitive diabetics' and

Invited Author's profile

Robert Semple is a Reader in Endocrinology and Metabolism and Honorary Consultant Endocrinologist at the University of Cambridge, UK. He is also a Fellow of Clare College, Cambridge, and Director of the Cambridge MB PhD programme. For the past 12 years, his research has focussed on rare human disorders of insulin action and growth. His research aims to identify novel, acquired or genetic defects underlying IR and related conditions, both to accelerate diagnosis and to enhance treatment of affected patients and draw inferences, through physiological study of affected patients, about the pathobiology of common forms of metabolic disease. This work has played a key part in the establishment of a National Severe Insulin Resistance Service in Cambridge.

Review	R K Semple	How many types of insulin resistance?	174:5	R210
--------	------------	---------------------------------------	-------	------

‘insensitive diabetics’ (generally older, obese and hypertensive, commonly with arteriosclerosis and exhibiting insidious disease onset with rare ketosis and insulin-induced hypoglycaemia) (1). Physiological studies showed that ‘insulin insensitive’ patients were slow to show a hypoglycaemic response to exogenous insulin (1). Upon development of radioimmunoassays in 1960, formal proof was obtained that insulin-resistant individuals do indeed have high or normal insulin levels in the face of hyperglycaemia (2, 3).

Now, nearly 100 years after the first use of insulin in diabetes, although the scourge of childhood death due to insulin deficiency is nearly vanquished, IR and the attendant hyperinsulinaemia sit at the centre of a web of burgeoning pandemic disease. They are associated strongly with obesity (4), type 2 diabetes (5), fatty liver disease (6), ‘metabolic’ DL (4), polycystic ovary syndrome (PCOS) (7) and some cancers (8). However, despite the clear documentation of these associations, understanding of the pathogenesis of IR, and whether and how it causes all the clinical conditions with which it is associated, remains incomplete.

Mechanisms of insulin action

Teasing out how insulin exerts metabolic and growth stimulating effects on target tissues has been the subject of painstaking biochemical study for many decades. Around 40 years ago, it was shown that insulin could bind competitively to hepatocyte membranes and thereby alter cellular metabolism (9), and shortly afterwards, the insulin receptor was identified biochemically (10). Sequencing of the gene for the human insulin receptor in 1985 demonstrated it to be a transmembrane dimeric tyrosine kinase (11), a member of what is now known as a large family of receptor tyrosine kinases of immense medical significance in endocrinology, cancer and beyond.

Cellular and animal studies have subsequently delineated many of the signalling events that pass the insulin signal from its receptor on the cell surface in order to change cellular metabolism and growth. A detailed account of these, which have comprehensively been reviewed elsewhere (12, 13), is beyond the scope of this article; however, a simplified schematic is shown in Fig. 1. In brief, insulin binding to the receptor induces phosphorylation of its intracellular domains, leading to recruitment of several insulin receptor substrates (IRS), the most important of which are IRS 1 and 2. These large phosphoproteins serve as a ‘platform’ that initiates downstream signalling pathways, among which the

phosphatidylinositol-3-kinase (PI3K)/AKT pathway and the MEK/ERK (formerly MAP kinase) pathway have received most attention. The PI3K pathway has traditionally been viewed as the key metabolic effector arm of the insulin signalling response, exerting critical metabolic actions in particular through AKT2, one of three isoforms of the serine–threonine kinase AKT that is enriched in insulin-responsive tissues (14).

Several general features of the insulin signalling ‘pathway’ bear particular note. Firstly, although signalling downstream from the receptor is often conceptualised as a series of branching pathways emanating from the receptor or IRS proteins, there is both extensive crosstalk among branches of the signalling pathway and also negative feedback within them so that they function as a rather more complex network than often implied by simple schematics. Moreover, most of the signalling pathways downstream from the insulin receptor are shared by other receptor tyrosine kinases, and in particular, the insulin-like growth factor 1 (IGF1) receptor, whose signalling network is nearly indistinguishable from that of insulin. How distinct biological responses are exerted by different growth factors and RTK ligands is not fully understood, though this is likely to depend in part on different patterns of ligand and receptor expression, and probably on different subcellular location of signalling molecules. Secondly, it has previously been pointed out that at several key points of insulin signal transduction, the signal may be transmitted by several different homologous proteins, some of which function as dimeric products of more than one gene (12). This introduces the potential for huge combinatorial signalling complexity. Finally, cellular studies are beginning to unpick the importance of different temporal sequences of signalling events after receptor stimulation, which, allied to different thresholds for activation of metabolic endpoints, introduces yet more tiers of signalling complexity (15).

All these considerations mean that, even at the refined level of cellular signalling, it is difficult to formulate one comprehensive, biologically meaningful definition of ‘insulin resistance (IR)’, and doing so with reference to only one or even a handful of downstream readouts may mask a more complex and patchy perturbation of cellular signalling. As will be discussed, this may well be important for common diseases related to IR.

Quantitative definitions of IR

The catastrophic short-term consequences of severe hypoglycaemia, and the damaging longer-term consequences of hyperglycaemia, have meant that it is the

Figure 1

Simplified schema of insulin signalling pathway with known human monogenic disorders. Genes that have been implicated in Mendelian disorders of insulin signalling are shown in red with the corresponding disease and year of discovery in linked boxes. Cellular processes stimulated by insulin are shown in green, while those inhibited by insulin are shown in red italics. INS, insulin; INSR, the insulin receptor; IRS1/2, insulin receptor substrate 1/2; PI3K is illustrated as a heterodimer of one of two catalytic subunits (p110 α or p110 β) and one of four regulatory subunits, three of which (p85 α , p55 α and p50 α) are encoded by the *PIK3R1* gene mutated in SHORT syndrome. PIP₂, phosphatidylinositol-(4,5)-bisphosphate;

PIP₃, phosphatidylinositol-(3,4,5)-trisphosphate; PDK1, 3-phosphoinositide-dependent protein kinase 1; AKT, protein kinase B; GSK3, glycogen synthase kinase 3; FOXO1, forkhead box protein O1; mTORC1/2, mammalian target of rapamycin complex 1/2; TBC1D4, gene encoding AKT substrate of 160kDa; BAD, BCL2-associated agonist of cell death; PDE3B, phosphodiesterase 3B; SREBP-1c, sterol regulatory element binding transcription factor 1; SHC, Src homology 2-containing protein; GRB2, growth factor receptor-bound protein 2; SOS, SOS Ras/Rho guanine nucleotide exchange (Son of Sevenless); MEK, MAPK/ERK kinase; ERK.

action of insulin on blood glucose, rather than on other aspects of intermediary metabolism or growth, that have been by far the major focus of attention over the past century (16). Thus, it is by reference to the ability of insulin to lower blood glucose that insulin sensitivity, and

thus resistance, is generally defined (17). The simplest definitions rely on measurement of fasting plasma insulin together with blood glucose. When blood glucose is in the normal range, IR may simply be defined using an arbitrary threshold established with reference to fasting insulin in

a reference population. Where beta cell decompensation and hence hyperglycaemia have arisen, use of an empirical index such as the homeostatic model assessment–IR index is of use. More complicated dynamic testing may also be used including oral or intravenous glucose tolerance testing with derivation of one of a variety of indices, while determination of the amount of insulin required in the face of a fixed infusion of glucose to maintain normal blood glucose (so-called hyperinsulinaemic euglycaemic clamping) is generally held to be the gold standard. However, although each of these approaches has utility, and while several permit assessment of different aspects of insulin's glucose lowering action, each neglects the effect of insulin on other processes such as amino acid and lipid metabolism. Although efforts have been made to develop surrogate indices of IR based on other plasma analytes (18, 19), none of these has yet been widely clinically adopted.

As insulin sensitivity is a continuous trait, thresholds for diagnosis of 'severe' IR are arbitrary and, ideally, should be defined with reference to appropriate controls, taking into account influences on insulin sensitivity such as age, sex, pubertal stage, obesity and concomitant medical illness. Operational examples of thresholds for defining severe IR are a requirement for exogenous insulin of greater than 3 units/kg per day to maintain normoglycaemia in those with absolute insulin deficiency or a fasting insulin greater than 150 pmol/l in lean adults without diabetes. However, given the complexity of setting valid numerical thresholds, especially once diabetes has developed, clinical evidence of severe IR from history and examination are of tremendous importance in timely diagnosis.

Mendelian randomisation to test causality in human disease

A key challenge in studies of IR-associated disease in humans arises from the difficulty in discerning cause and effect, if any, in observed associations. Causality may be implied though not proven from the sequence of appearance of clinical phenomena; however, the most direct test of whether a disease is caused by an associated disorder in humans is to directly introduce that perturbation and to observe whether the disease ensues. In most cases, however, even where this might technically be possible, the postulated deleterious nature of the first perturbation and/or the time course over which the disease of interest develops render this impractical or unethical. Natural genetic variation may sometimes be used opportunistically to undertake equivalent studies, where a primary genetic cause of a candidate mediator of

disease can be identified. An often cited paradigm for this so-called 'Mendelian randomisation' approach lies in Brown's and Goldstein's use of patients with LDL receptor mutations and thus primary elevation of LDL cholesterol to provide strong evidence for the causal link between LDL cholesterol and atherosclerosis (20). The principle of Mendelian randomisation is illustrated in Fig. 2.

Mendelian randomisation would also be of value in principle in testing which of the IR-associated diseases are indeed caused by IR, as long as primary genetic causes of IR, with no independent link to the diseases being studied, could be identified. Such groups of patients with primary

Figure 2

Principle of Mendelian randomisation. Algorithm for the use of Mendelian disorders to test possible causality underlying biological association, illustrated by the example of LDL cholesterol and atherosclerosis.

Review	R K Semple	How many types of insulin resistance?	174:5	R213
--------	------------	---------------------------------------	-------	------

monogenic IR are indeed now known and continue to be discovered, and their value in informing studies of common IR-related disease will be the main focus of the rest of this article.

An analytic 'Toolkit': subtypes of human monogenic IR

It is 27 years since the first single gene form of severe IR, caused by mutations in the gene encoding the insulin receptor, was discovered (21, 22), and in the past 17 years, further disorders have been added to this, latterly with increasing rapidity as modern genetic technologies have been brought to bear (23). As the number of known monogenic disorders has increased, it has become apparent that IR is not monomorphic and, indeed, that most single gene causes of severe IR do not directly affect insulin cellular signalling pathways. Instead, many defects do directly affect the development or function of adipose tissue, with secondary effects on insulin action *in vivo*. Single gene causes of severe IR may thus be divided into groups loosely defined by the tissue or cellular process affected. It is these groupings which are lending increasing power to Mendelian randomisation.

Primary insulin signalling defects

The first human loss-of-function mutations were reported in the insulin receptor in 1988, both in Donohue syndrome, an extreme infantile form of failure to thrive with IR (22), and with the less severe 'type A' form of IR (21), which usually becomes manifest peripubertally. Since this time, more than 150 different mutations within the insulin receptor have been discovered, with relatively few recurring mutations, and the phenotypic spectrum has been well described elsewhere (23, 24, 25). A single family with four members affected by severe IR due to a loss-of-function mutation in AKT2 was also described in 2004 and was the second example of a highly penetrant Mendelian form of severe IR due to mutation of a canonical insulin signalling gene (26). Mutations in TBC1D4, involved in regulating exocytosis of glucose-transporting SLC2A4 (GLUT4) vesicles in response to insulin, were also described in association with an apparently muscle-selective form of severe IR (27), and this was later confirmed in a Greenland population, where a similar genetic defect accounts for a significant part of the variance of insulin sensitivity (28).

The latest form of severe IR established to be caused by mutation of an insulin signalling gene is seen in the

context of SHORT syndrome, denoted by short stature, joint hyperextensibility and hernias, ocular depression, Rieger anomaly and teething delay. Although not part of the acronym, severe IR has been shown to be common in this disorder, and in 2012, the underlying cause was established to be mutations in the *PIK3R1* gene (29, 30, 31). *PIK3R1* encodes three of the regulatory subunits of type 1A PI3K, which are critical for the coupling of insulin binding to its receptor to activation of AKT2 and other downstream signalling molecules (12). Primary insulin signalling disorders are schematised in Fig. 1.

Primary lipodystrophies

Inherited disorders of adipose tissue development, or lipodystrophies, were described clinically from at least the 1920s (32); however, it has only been in this century that they have yielded to genetic study, with many different causal genetic defects now identified. Inherited lipodystrophies are most commonly divided into those that feature complete, or near complete absence of all adipose tissue, so-called congenital generalised lipodystrophy (CGL) and those in which only some adipose tissue depots are affected, so-called familial partial lipodystrophy.

CGL, often known as Berardinelli-Seip congenital lipodystrophy, is caused in around 95% of cases by biallelic mutations in one of two genes, namely, *AGPAT2*, encoding 1-acylglycerol-3-phosphate O-acyltransferase 2 (*AGPAT2*) (33, 34), and *BSCL2*, which encodes a transmembrane endoplasmic reticulum protein named seipin (35). Seipin plays a role in preadipocyte differentiation (36) and in the regulation of lipid droplet synthesis and function (37, 38). Smaller numbers of patients with CGL have also been reported with loss-of-function mutations in *CAV1* (39) or *PTRF* (40), the latter associated also with myopathy. Both these genes encode key structural components of small plasma membrane invaginations, so-called *caveolae*, which account for up to 40% of the adipocyte surface area, and play an important role in organising cell signalling and lipid trafficking. Most recently, genetic disruption of *PCYT1A*, encoding phosphate cytidylyltransferase 1, which plays a key role in *de novo* phosphatidylcholine biosynthesis, has also been demonstrated in patients with CGL (41).

Familial partial lipodystrophies (FPLDs) are collectively far more common than CGL. They are usually not clinically manifest until puberty when perturbation of pubertal adipose accretion unmasks the underlying abnormality. Women are more severely affected than men, likely due in large part to the naturally greater adiposity of women than men. At least seven genes with

Review	R K Semple	How many types of insulin resistance?	174:5	R214
--------	------------	---------------------------------------	-------	------

diverse roles in adipocyte biology have now been linked to partial lipodystrophy (Table 1). The first of these, *LMNA*, encodes a nearly ubiquitously expressed intermediate filament protein, Lamin A/C, which forms part of a structural network of proteins supporting the nuclear membrane (42). Different *LMNA* mutations have also been linked to other disorders including muscular dystrophy, dilated cardiomyopathy, Charcot–Marie–Tooth neuropathy, premature aging syndromes, restrictive dermopathy and various overlap syndromes (43). The mechanism(s) linking *LMNA* mutations to lipodystrophy are not fully understood, but roles have been proposed for structural defects in the nuclear envelope and/or abnormal binding of the nuclear lamina to chromatin and transcription factors, thus altering gene expression.

The second gene to be affected in familial partial lipodystrophy is *PPARG* (44), encoding PPAR γ , a nuclear hormone receptor most highly expressed in adipose tissue where it is essential for adipocyte differentiation and which is the target of the thiazolidinedione class of antidiabetic agents. All known pathogenic mutations are heterozygous, and most have been shown to inhibit the function of co-expressed WT protein.

Although the remaining genes now known to cause partial lipodystrophy collectively account for only a tiny proportion of cases, their discovery has highlighted the critical importance of lipid droplet dynamics in human metabolic homeostasis, with defects in either *CIDEA* (45) or perilipin 1 (perilipin) (46), each of which resides on the surface of the lipid droplet and plays a regulatory role in triglyceride mobilisation, now implicated in human partial lipodystrophy.

Of note, several of the primary insulin signalling disorders also feature some degree of lipodystrophy, in keeping with an important role for insulin and IGF1 signalling in adipocyte differentiation. This is particularly

pronounced in the case of *AKT2* loss of function (26) and in SHORT syndrome (29, 31), but a generalised paucity of adipose tissue is also commonly described in the more severe insulin receptoropathies. The potential importance of this for the IR subphenotype that is clinically expressed is discussed below.

Complex monogenic disorders featuring severe IR

A clinically heterogeneous group of more complex monogenic conditions also exists that features severe IR disproportionate to whole body adiposity. These include the premature ageing syndromes Werner syndrome (47, 48) and Bloom syndrome (49), both caused by defects in DNA helicases, and a very recently described syndrome of primordial dwarfism caused by mutations in the *NSMCE2* gene, which encodes a component of the structural maintenance of chromosomes 5/6 complex (50). Each of these genes plays some role in permitting dividing cells to tolerate DNA damage and in the maintenance of telomeres. Another recently described syndrome featuring loss of s.c. fat as well as severe IR is caused by heterozygous loss-of-function mutations in *POLD1*, encoding DNA polymerase delta, the dominant lagging strand DNA synthase in human cells (51). This adds to the impression that some aspects of DNA damage repair or replication are critical to metabolic homeostasis. This notion gives further credence by the observation that childhood irradiation as part of cancer therapy predisposes to later DL and IR, which may sometimes be severe (52). Modelling of this phenomenon in leptin deficient, severely obese mice suggests that irradiation leads to adipocyte and preadipocyte death, effectively reducing adipose tissue expandability, leading to a worsened metabolic state if, rather than despite, a reduction in adipose gain after irradiation (53).

Table 1 Known genetic forms of partial lipodystrophy.

Clinical disorder	Gene	Inheritance	Distribution of adipose loss	Comment
FPLD2	<i>LMNA</i>	AD	S/C sparing head, neck and labia majora	Commonest cause of FPLD. May be mistaken for Cushing's syndrome due to preserved head and neck fat LD may be subtle, limited to limbs and associated with centripetal obesity
FPLD3	<i>PPARG</i>	AD	Limbs	
FPLD4	<i>PLIN1</i>	AD	Limbs	Single family reported to date Single patient reported Not always seen. Not associated with dyslipidaemia Also loss of limb muscle, scleroderma-like features, deafness and male hypogonadism
FPLD6	<i>AKT2</i>	AD	Limbs	
FPLD5	<i>CIDEA</i>	AR	Limbs	
SHORT syndrome	<i>PIK3R1</i>	AD or sporadic	S/C, variable	
MDP syndrome	<i>POLD1</i>	AD or sporadic	S/C	

MDP, mandibular hypoplasia, deafness, progeroid features; AD, autosomal dominant; AR, autosomal recessive.

Review	R K Semple	How many types of insulin resistance?	174:5	R215
--------	------------	---------------------------------------	-------	------

A further tantalising hint of a specific relationship between a particular cellular function and whole body metabolic IR comes from the observation of severe IR in patients with genetic defects in different centrosomal components including *ALMS1* (54), *PCNT* (55, 56) and *POC1A* (57). Teasing out the pathogenic link between the defined molecular defect and severe systemic IR in these disorders promises to yield novel insights in future into the mechanistic basis of IR. Given the complex and pleiotropic nature of the genetic defects, however, these conditions are not suitable as a basis for the Mendelian randomisation approach.

Application of Mendelian randomisation to obesity-associated metabolic disease

Many of the disorders described above, where the cellular function and immediate physiological consequences of the genetic defect can be defined with confidence, are appropriate instruments to use in a Mendelian randomisation-based attempt to address important questions about the causes and consequences of human IR. Several of these questions will be considered in turn, and the lessons to be drawn from monogenic disease will be assessed.

How does obesity give rise to IR?

Cross-sectional and longitudinal studies strongly suggest that obesity is the proximate cause of most pandemic IR; however, the question then arises of the mechanistic link between these phenomena. Study of humans with single gene defects producing severe, early onset, hyperphagic obesity, for example, due to genetic deficiency of either leptin or its receptor, confirm that IR is common in this group; however, it is often not severe, and its severity is often somewhat less than that of the concomitant obesity. This implies that there are factors other than severity of obesity which influence the degree of obesity-related IR. Importantly, at the opposite end of the adipose spectrum, patients with CGL, who have negligible adipose tissue, develop extremely severe IR, DL and fatty liver; thus, monogenic lipodystrophy has been argued to be a potentially aetiologically informative model of the common metabolic syndrome (58).

These human observations give strong support to the hypothesis that it is relative adipose failure that is critical to the pathogenesis of the obesity-related metabolic syndrome rather than the degree of obesity. According to this idea, the capacity of adipose tissue to continue expanding, while safely storing excess calories in the form

of triglyceride, is finite, and when this capacity, which shows considerable inter-individual variation, is exceeded, then 'adipose failure', with knock-on effects on other, insulin-sensitive tissues, ensues (59, 60, 61). CGL represents an extreme example of this, where the adipose energy buffering capacity is negligible. In this setting, patients remain lean; however, loss of the capacity of adipose tissue to buffer excess caloric intake by serving as a sump for free fatty acids and, to a much lesser extent, as a site of disposal of glucose is likely to result in remodelling of systemic free fatty acid flux, with increased uptake in other metabolically critically tissues such as liver, skeletal muscle and pancreatic beta cells, where harmful effects including loss of insulin sensitivity, secretory dysfunction and apoptosis may result (62). This series of maladaptive changes are collectively unified by the term lipotoxicity.

In many monogenic partial lipodystrophies, significant adipose depots remain, and indeed, it is not uncommon for whole body adiposity to be elevated, as assessed by direct measurement and by surrogate indices such as plasma leptin levels. In these conditions, it is adipose topography which appears to be critical to metabolic homeostasis. In all known genetic forms of partial lipodystrophy which feature severe IR, femorogluteal depots are affected, while central and omental fats are often preserved or increased, meaning that Cushing's syndrome is commonly invoked as a differential diagnosis at first assessment. The impression that lower body s.c. fat is critical for metabolic health is bolstered by the observation that in acquired partial lipodystrophy, where s.c. fat is usually lost in a craniocaudal direction to the level of the umbilicus, there is only a low rate of systemic metabolic derangement (63). Distributions of fat loss in partial lipodystrophies are illustrated in Fig. 3.

Importantly, although some of the genes involved in lipodystrophy are expressed in cells other than adipocytes, raising concerns that they may exert phenotypic effects unrelated to lipodystrophy, thus invalidating them as instruments for Mendelian randomisation-based analysis, the closely similar metabolic phenotype seen in patients with acquired, autoimmune, lipodystrophy and the fact that at least one of the genes implicated in genetic lipodystrophy, *PLIN*, is adipocyte-specific (46), argues against this.

Additional mechanisms beyond loss of energy buffering have been invoked to account for the link between increased adipose tissue and IR. One of these is perturbation of circulating levels of adipose-derived hormones, so-called adipokines. Evidence for a large number of these has now been put forward, but in most cases, evidence for

Figure 3

Distribution of adipose loss in lipodystrophies. Areas of deficient adipose tissue are illustrated in red, with preserved

adipose tissue represented in green. WAT, white adipose tissue; BM, bone marrow; NAFLD, non-alcoholic fatty liver disease.

a critical role in setting systemic insulin sensitivity is mixed, and most are best viewed as playing an accessory role only. Apart from leptin, best established of the *bona fide* adipokines, adiponectin, a large protein with homology to complement factor C1q, has attracted the most sustained and intense attention. It circulates at high levels in plasma as a complex mixture of oligomeric or multimeric forms. Unusually, increasing levels of adiposity and IR are associated with lower plasma adiponectin concentration, with a concomitant shift from higher molecular weight species to lower molecular weight forms. Much of the evidence that adiponectin is an insulin sensitiser come from murine studies. Genetic ablation of the adiponectin-encoding gene in mice produced IR in some, though not all reports, while administration of various different oligomeric adiponectin preparations has been shown in mice to improve insulin sensitivity (64, 65).

The metabolic role of adiponectin in humans is less clear. Until recently evidence for a causal role of low adiponectin in IR was largely correlative, based on studies of association of plasma adiponectin with metabolic disorders (64). However, several genetic studies, involving Mendelian randomisation, have also now been brought to bear on this issue. Plasma adiponectin is usually either preserved or frankly elevated, sometimes dramatically so, in the face of either genetic or acquired defects in insulin receptor function (66, 67, 68), and more limited evidence suggests that this is also true in SHORT syndrome, arguing that proximal defects in insulin signalling raise plasma

adiponectin. This observation is not explained, but demonstrates that suppressed plasma adiponectin is not a consequences of reduced insulin signalling. Indeed, it may be viewed as an attempt to ameliorate insulin sensitivity in the face of a fixed signalling defect. A more direct test of the potential of adiponectin to influence insulin sensitivity comes from true Mendelian randomisation, whereby association with IR of genetic variants at the locus of the adiponectin gene, *ADIPOQ*, that induces primary changes in plasma adiponectin, are assessed. This approach has been adopted by several groups with mixed results. Two studies found that genetically determined lowering of plasma adiponectin did indeed associate with reduced insulin sensitivity (69, 70), while the largest study, with the greatest statistical power, found no such association (71). One family has been reported in which a rare missense *ADIPOQ* variant suppressing plasma adiponectin was found to associate with IR (72). More systematic metabolic study of more families with rare *ADIPOQ* alleles identified in large-scale exome sequencing studies will be of value to identify any that suppress plasma adiponectin and to clarify the association further.

Does IR cause fatty liver and DL?

Understanding the relationship between plasma adiponectin levels and IR is important for the understanding of the pathogenesis of IR, and this is given extra weight by the potential exploitation of adiponectin for therapeutic

Review	R K Semple	How many types of insulin resistance?	174:5	R217
--------	------------	---------------------------------------	-------	------

benefit. Nevertheless low plasma adiponectin is a biochemical surrogate rather than a clinical disease. In contrast, the fatty liver disease spectrum, which is intimately linked to a pattern of DL characterised by low plasma HDL cholesterol and high plasma triglyceride, exacts an immense toll of human suffering. To this, too, Mendelian randomisation using rare disorders of insulin action may be applied. Before diabetes supervenes, primary IR due to genetic defects in the insulin receptor is a model of global, compensated IR with extreme hyperinsulinaemia. Strikingly, however, patients with genetic or acquired INSR defects appear to be entirely protected from 'metabolic' DL (73). Moreover, they do not exhibit increased liver fat, and rates of hepatic *de novo* lipogenesis, which has been proposed to be an important contributor to fatty liver and DL in human IR (74, 75), are similar to controls (73). This argues that fatty liver disease and DL can be accounted for neither by generalised IR nor by severe hyperinsulinaemia acting through non-insulin receptor-dependent pathways.

Analysis of IR due to downstream signalling defects produces contrasting results: preliminary data suggests that in SHORT syndrome due to mutations in PIK3R1, lipid profiles are normal despite severe IR (31), while in contrast, two patients studied with severe IR due to defects in AKT2, three steps down the classical insulin signalling pathway from the insulin receptor and one from PI3K, did show exaggerated metabolic DL with severe fatty liver (73).

These human observations suggest that 'partial' IR – that is, IR affecting only some parts of the insulin signalling network – may play an important role in the pathogenesis of major IR-related pathologies. This hypothesis was first widely promulgated in the 1980s (4) and has been supported previously by observations made in genetically modified mice (76). In the current context, 'partial' IR may be regarded as a cell autonomous phenomenon, whereby the pathway required for hyperinsulinaemia to drive liver fat accumulation and atherogenic patterns of VLDL secretion from liver cells depends on the insulin receptor and PIK3R1 but not AKT2, implying a signalling pathway proximal to AKT2 that drives hepatic *de novo* lipogenesis. However, this is at odds with most of the considerable body of murine data, most of which suggest that AKT2 is critical in mediating the effect of insulin to drive liver fat accumulation (77, 78). Other possibilities are that the critical elements of 'partial' IR are differential effects of the same fixed defect in insulin signalling on different downstream pathways due to differing thresholds for activation (79) and/or differential effects of the same fixed defect on insulin sensitivity of

different tissues (80). Thus, it may be that the dominant effect in humans of the AKT2 mutation reported is in adipose tissue, where AKT2 is involved in suppressing adipocyte differentiation and the suppression of lipolysis, and that the resulting 'adipose failure' swamps reduced ability of high insulin to drive *de novo* lipogenesis in the liver. Possible models of partial IR are schematised in Fig. 4.

Does IR cause PCOS?

Like type 2 diabetes, PCOS has a high prevalence, estimated to be of the order of 10% in Western populations and confers a high burden of morbidity, due to, in particular, subfertility and psychological distress related to the cosmetic effects of clinical hyperandrogenism. However, like the label 'type 2 diabetes', the label 'PCOS' does not imply aetiopathogenesis. Instead it simply describes the syndrome and is the unifying designation for the common clinical expression of a range of primary underlying defects rather than being a single entity. Indeed, although PCOS has been the focus of an extensive literature encompassing clinical, model organism and cellular studies, debate continues about key aspects of its pathogenesis.

IR and the metabolic syndrome are very well described associations of PCOS, with a prevalence in the region of 50–70%, depending on definitions used and population studied (81, 82). However, there has been uncertainty as to whether IR in PCOS has unique characteristics. Moreover primary androgen excess of a variety of causes may reduce insulin sensitivity in adipose and other tissues; so, an argument has sometimes been made that the IR in PCOS is caused by ovarian dysfunction rather than *vice versa*. Where IR is accepted as a cause of some forms of PCOS, partial IR, as in the liver, has been invoked to explain PCOS, with the ovary argued to be an non-insulin-resistant 'bystander' tissue, responding to high levels of circulating insulin set by selective defects in the metabolic actions of insulin in other tissues (83).

In the face of complex and sometimes competing hypotheses, observations in humans with single gene severe IR are simple: ovulatory dysfunction and hyperandrogenism, which may be very severe, are seen in nearly all known monogenic forms of severe IR. They are seen in exaggerated form in the context of genetic or acquired insulin receptoropathy, of downstream insulin signalling defects in AKT2 or PIK3R1, of partial or generalised lipodystrophy and pleiotropic syndromes, all of which commonly exhibit severe clinical and biochemical hyperandrogenism, oligo or amenorrhoea and classical

Figure 4

Models for role of partial IR in disease pathogenesis. Hypoglycaemic and lipogenic actions of insulin are used to illustrate each model. (A) Cell autonomous selective signalling defect model, with disproportionate attenuation of insulin action to lower glucose, leading to compensatory hyperinsulinaemia, and exposure of intact insulin signalling pathway, to higher levels of insulin action. (B) Differential dose-response model. This is a variation of the above theme whereby a fixed reduction in insulin action across all arms of the signalling pathway has a

greater effect on arms of the pathway showing a shallow dose-response curve to insulin than those showing a steep dose-response curve. (C) Differential tissue resistance to insulin model. Where an organ with the dominant role in insulin-mediated glucose disposal is more IR than other tissues, it effectively sets the level of systemic insulin, potentially exposing other more responsive tissues to abnormally high insulin action. These models are non-mutually exclusive and may all play some role *in vivo*.

Review	R K Semple	How many types of insulin resistance?	174:5	R219
--------	------------	---------------------------------------	-------	------

polycystic ovaries. In other words, while DL and fatty liver are seen only in some types of IR, the ovarian phenotype is nearly universal. There is no evidence that it differs between global IR at the level of the receptor and selective IR affecting PI3K/AKT signalling, generally regarded as the more metabolic arm of the insulin signalling pathway, in keeping with prior evidence that PI3K does not mediate the effects of insulin on human granulosa cells in culture (84). Furthermore, although perturbed androgen metabolism has been suggested to play an important role in PCOS in some forms of IR (85), the severe PCOS seen in generalised lipodystrophy argues that adipose androgen is not an obligate link between severe IR and PCOS.

In Donohue syndrome, cystic ovarian enlargement may be seen in infancy, and this can be massive even when no functional insulin receptors are expressed (86, 87). While this ovarian pathology is not identical to that of PCOS, it is plausible to suggest that it shares the same underlying mechanism, namely, synergic stimulation of follicles by extremely elevated insulin and the gonadotrophins which are elevated in the early months of life before secondary suppression. If this parallel holds true, then this suggests that the adverse effects of insulin on the ovary need not be exerted through the insulin receptor itself. These observations suggest that the ovarian component of the IR syndrome, quite unlike fatty liver and DL, do not require intact INSR function, although they do require severe hyperinsulinaemia. These findings may be reconciled by hypothesising that very high levels of insulin enhance signalling through the IGF receptor, in keeping with a large body of evidence for an important role for IGFs in follicular maturation (88, 89). PCOS is reported to be common in premenopausal women with acromegaly (90); however, it is not usually as severe as that seen in the context of severe IR. This discrepancy may be accounted for by endocrine or paracrine enhancement of IGF1 bioavailability or receptor expression in IR in addition to an IGF-like action of very high levels of circulating insulin.

Observations in defined monogenic forms of severe IR may not necessarily be extrapolatable to 'common' PCOS. On the other hand, the extremely high penetrance of a PCOS phenotype in the face of a diverse array of genetic forms of severe IR and the reversible PCOS seen in acquired states of insulin receptor dysfunction (91), which are not mimicked by insulin deficient states, show that IR with compensatory hyperinsulinaemia is sufficient to cause a severe PCOS-like state without needing to invoke additional mechanisms or the importance of a specific subtype of IR. The ability of primary hyperandrogenaemia

due to genetic abnormalities in steroid metabolism also to phenocopy prevalent PCOS (92) reaffirm, however, that there is more than one pathway to the PCOS phenotype.

Does IR cause cancer?

Increasing attention has been paid in recent years to the association in population-based studies between IR and some cancers including those of the breast, colon, prostate and endometrium (93). For some cancers, there are plausible endocrine or metabolic explanations for a causal link between IR and the tumour that do not involve direct effects of insulin on the tumour itself. For example, hyperinsulinaemia induces the oestrogen replete but oligomenorrhoeic state commonly seen in PCOS, indirectly increasing risk of endometrial cancer (94). Furthermore, it is well established that the fatty liver associated with lipodystrophy is not benign, but rather features a high risk of cirrhosis and ultimately hepatocellular cancer in lipodystrophies (95). However, it has also been argued, with some experimental evidence, that insulin may directly drive tumour growth (93, 96).

Cancer pathogenesis is complex and multifactorial, and hyperinsulinaemia is likely to be only one facilitatory element in a complex sequence of events. Nevertheless it is to be predicted that in severe monogenic forms of IR, the cancer risk may be further increased from the risk attributed to IR in the general population. Indeed, several non-malignant elements of the severe IR syndrome, namely, the dermal and epidermal hyperplasia of acanthosis nigricans, the organomegaly of infantile receptoropathy and the pseudo-acromegaloid soft tissue overgrowth common to many forms of IR, attest to the tissue growth-promoting effects of hyperinsulinaemia. In the most severe hyperinsulinaemia seen in Donohue syndrome, ovarian tumours may arise in infancy (86, 87), while colonic polyposis is also sometimes seen in either recessive or dominant insulin receptoropathies. Thus, while syndromes of severe IR, except those featuring an underlying defect in DNA damage repair such as Werner and Bloom syndromes, are not penetrant cancer predisposition syndromes, the mitogenic consequences of sustained severe hyperinsulinaemia are apparent, and this adds weight to the notion of a role for hyperinsulinaemia *per se* in increasing prevalent cancer risk.

Conclusions and future directions

IR is not a disease in itself, but rather an endocrine derangement associated with several pandemic diseases

Review	R K Semple	How many types of insulin resistance?	174:5	R220
--------	------------	---------------------------------------	-------	------

and tissue pathologies. Study of rare single gene forms of IR has yielded valuable insights into several aspects of the pathogenesis of these diseases. Current findings are consistent with the ‘adipose failure’ model linking obesity to prevalent metabolic disease and suggest that compensatory hyperinsulinaemia may play a critical role in the pathogenesis of PCOS as well as fatty liver and metabolic DL. Specifically, differences among different subtypes of insulin signalling defect demonstrate that IR with respect to glucose lowering may be uncoupled from other components of the prevalent metabolic syndrome including fatty liver and DL, depending on the precise nature of the underlying signalling defect. In contrast, the ‘trophic’ features of IR including acanthosis nigricans, soft tissue overgrowth and PCOS are seen in essentially all insulin resistant hyperinsulinaemic states. Teasing out the mechanistic basis of these findings in humans offers the possibility to identify novel strategies to isolate and treat several different components of the complex of obesity and IR-related diseases.

Declaration of interest

The author declares that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Funding

Dr Semple has received speaker fees from Novo Nordisk and Sandoz. Dr Semple is supported by a Senior Research Fellowship from the Wellcome Trust (grant number WT098498).

References

- Himsworth HP. Insulin deficiency and insulin inefficiency. *BMJ* 1940 **1** 719–722. (doi:10.1136/bmj.1.4139.719)
- Yalow RS & Berson SA. Plasma insulin concentrations in nondiabetic and early diabetic subjects. Determinations by a new sensitive immunoassay technic. *Diabetes* 1960 **9** 254–260. (doi:10.2337/diab.9.4.254)
- Karam JH, Grodsky GM & Forsham PH. Excessive insulin response to glucose in obese subjects as measured by immunochemical assay. *Diabetes* 1963 **12** 197–204. (doi:10.2337/diab.12.3.197)
- Reaven GM. Pathophysiology of insulin resistance in human disease. *Physiological Reviews* 1995 **75** 473–486. (doi:10.1002/dmr.5610090503)
- Hanson RL, Imperatore G, Bennett PH & Knowler WC. Components of the “metabolic syndrome” and incidence of type 2 diabetes. *Diabetes* 2002 **51** 3120–3127. (doi:10.2337/diabetes.51.10.3120)
- Utzschneider KM & Kahn SE. Review: the role of insulin resistance in nonalcoholic fatty liver disease. *Journal of Clinical Endocrinology and Metabolism* 2006 **91** 4753–4761. (doi:10.1210/jc.2006-0587)
- Dunaif A, Segal KR, Futterweit W & Dobrjansky A. Profound peripheral insulin resistance, independent of obesity, in polycystic ovary syndrome. *Diabetes* 1989 **38** 1165–1174. (doi:10.2337/diab.38.9.1165)
- Pollak M. Insulin and insulin-like growth factor signalling in neoplasia. *Nature Reviews. Cancer* 2008 **8** 915–928. (doi:10.1038/nrc2536)
- Freychet P, Roth J & Neville DM Jr. Insulin receptors in the liver: specific binding of (¹²⁵I)insulin to the plasma membrane and its relation to insulin bioactivity. *PNAS* 1971 **68** 1833–1837. (doi:10.1073/pnas.68.8.1833)
- Yip CC, Yeung CW & Moule ML. Photoaffinity labeling of insulin receptor of rat adipocyte plasma membrane. *Journal of Biological Chemistry* 1978 **253** 1743–1745.
- Ullrich A, Bell JR, Chen EY, Herrera R, Petruzzelli LM, Dull TJ, Gray A, Coussens L, Liao YC, Tsubokawa M *et al.* Human insulin receptor and its relationship to the tyrosine kinase family of oncogenes. *Nature* 1985 **313** 756–761. (doi:10.1038/313756a0)
- Taniguchi CM, Emanuelli B & Kahn CR. Critical nodes in signalling pathways: insights into insulin action. *Nature Reviews. Molecular Cell Biology* 2006 **7** 85–96. (doi:10.1038/nrm1837)
- Cohen P. The twentieth century struggle to decipher insulin signalling. *Nature Reviews. Molecular Cell Biology* 2006 **7** 867–873. (doi:10.1038/nrm2043)
- Whiteman EL, Cho H & Birnbaum MJ. Role of Akt/protein kinase B in metabolism. *Trends in Endocrinology and Metabolism* 2002 **13** 444–451. (doi:10.1016/S1043-2760(02)00662-8)
- Humphrey SJ, James DE & Mann M. Protein phosphorylation: a major switch mechanism for metabolic regulation. *Trends in Endocrinology and Metabolism* 2015 **26** 676–687. (doi:10.1016/j.tem.2015.09.013)
- McGarry JD. What if Minkowski had been ageusic? An alternative angle on diabetes *Science* 1992 **258** 766–770. (doi:10.1126/science.1439783)
- Muniyappa R, Lee S, Chen H & Quon MJ. Current approaches for assessing insulin sensitivity and resistance in vivo: advantages, limitations, and appropriate usage. *American Journal of Physiology - Endocrinology and Metabolism* 2008 **294** E15–E26. (doi:10.1152/ajpendo.00645.2007)
- Guerrero-Romero F, Simental-Mendia LE, Gonzalez-Ortiz M, Martinez-Abundis E, Ramos-Zavala MG, Hernandez-Gonzalez SO, Jacques-Camarena O & Rodriguez-Moran M. The product of triglycerides and glucose, a simple measure of insulin sensitivity. Comparison with the euglycemic-hyperinsulinemic clamp. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 3347–3351. (doi:10.1210/jc.2010-0288)
- Finucane FM, Luan J, Wareham NJ, Sharp SJ, O’Rahilly S, Balkau B, Flyvbjerg A, Walker M, Hojlund K, Nolan JJ *et al.* Correlation of the leptin:adiponectin ratio with measures of insulin resistance in non-diabetic individuals. *Diabetologia* 2009 **52** 2345–2349. (doi:10.1007/s00125-009-1508-3)
- Goldstein JL & Brown MS. The low-density lipoprotein pathway and its relation to atherosclerosis. *Annual Review of Biochemistry* 1977 **46** 897–930. (doi:10.1146/annurev.bi.46.070177.004341)
- Yoshimasa Y, Seino S, Whittaker J, Kakehi T, Kosaki A, Kuzuya H, Imura H, Bell GI & Steiner DF. Insulin-resistant diabetes due to a point mutation that prevents insulin proreceptor processing. *Science* 1988 **240** 784–787. (doi:10.1126/science.3283938)
- Kadowaki T, Bevins CL, Cama A, Ojamaa K, Marcus-Samuels B, Kadowaki H, Beitz L, McKeon C & Taylor SI. Two mutant alleles of the insulin receptor gene in a patient with extreme insulin resistance. *Science* 1988 **240** 787–790. (doi:10.1126/science.2834824)
- Semple RK, Savage DB, Cochran EK, Gorden P & O’Rahilly S. Genetic syndromes of severe insulin resistance. *Endocrine Reviews* 2011 **32** 498–514. (doi:10.1210/er.2010-0020)
- Semple RK, Savage DB, Brierley GV & O’Rahilly S. Syndromes of Severe Insulin Resistance and/or Lipodystrophy. In *Genetic Diagnosis of Endocrine Disorders* (Second edition), pp 307–324. Elsevier 2016. (doi:10.1016/B978-0-12-800892-8.00022-1)
- Taylor SI, Cama A, Accili D, Barbetti F, Quon MJ, de la Luz Sierra M, Suzuki Y, Koller E, Levy-Toledano R, Wertheimer E *et al.* Mutations in the insulin receptor gene. *Endocrine Reviews* 1992 **13** 566–595. (doi:10.1210/edrv-13-3-566)
- George S, Rochford JJ, Wolfrum C, Gray SL, Schinner S, Wilson JC, Soos MA, Murgatroyd PR, Williams RM, Acerini CL *et al.* A family with

Review	R K Semple	How many types of insulin resistance?	174:5	R221
<p>severe insulin resistance and diabetes due to a mutation in AKT2. <i>Science</i> 2004 304 1325–1328. (doi:10.1126/science.1096706)</p> <p>27 Dash S, Sano H, Rochford JJ, Semple RK, Yeo G, Hyden CS, Soos MA, Clark J, Rodin A, Langenberg C <i>et al.</i> A truncation mutation in TBC1D4 in a family with acanthosis nigricans and postprandial hyperinsulinemia. <i>PNAS</i> 2009 106 9350–9355. (doi:10.1073/pnas.0900909106)</p> <p>28 Moltke I, Grarup N, Jorgensen ME, Bjerregaard P, Treebak JT, Fumagalli M, Korneliusen TS, Andersen MA, Nielsen TS, Krarup NT <i>et al.</i> A common Greenlandic TBC1D4 variant confers muscle insulin resistance and type 2 diabetes. <i>Nature</i> 2014 512 190–193. (doi:10.1038/nature13425)</p> <p>29 Thauvin-Robinet C, Auclair M, Duplomb L, Caron-Debarle M, Avila M, St-Onge J, Le Merrer M, Le Luyer B, Heron D, Mathieu-Dramard M <i>et al.</i> PIK3R1 mutations cause syndromic insulin resistance with lipoatrophy. <i>American Journal of Human Genetics</i> 2013 93 141–149. (doi:10.1016/j.ajhg.2013.05.019)</p> <p>30 Dymont DA, Smith AC, Alcantara D, Schwartzentruber JA, Basel-Vanagaite L, Curry CJ, Temple IK, Reardon W, Mansour S, Haq MR <i>et al.</i> Mutations in PIK3R1 cause SHORT syndrome. <i>American Journal of Human Genetics</i> 2013 93 158–166. (doi:10.1016/j.ajhg.2013.06.005)</p> <p>31 Chudasama KK, Winnay J, Johansson S, Claudii T, Konig R, Haldorsen I, Johansson B, Woo JR, Aarskog D, Sagen JV <i>et al.</i> SHORT syndrome with partial lipodystrophy due to impaired phosphatidylinositol 3 kinase signaling. <i>American Journal of Human Genetics</i> 2013 93 150–157. (doi:10.1016/j.ajhg.2013.05.023)</p> <p>32 Seip M & Trygstad O. Generalized lipodystrophy. <i>Archives of Disease in Childhood</i> 1963 38 447–453. (doi:10.1136/adc.38.201.447)</p> <p>33 Garg A, Wilson R, Barnes R, Arioglu E, Zaidi Z, Gurakan F, Kocak N, O'Rahilly S, Taylor SI, Patel SB <i>et al.</i> A gene for congenital generalized lipodystrophy maps to human chromosome 9q34. <i>Journal of Clinical Endocrinology and Metabolism</i> 1999 84 3390–3394. (doi:10.1210/jcem.84.9.6103)</p> <p>34 Agarwal AK, Arioglu E, De Almeida S, Akkoc N, Taylor SI, Bowcock AM, Barnes RI & Garg A. AGPAT2 is mutated in congenital generalized lipodystrophy linked to chromosome 9q34. <i>Nature Genetics</i> 2002 31 21–23. (doi:10.1038/ng880)</p> <p>35 Magre J, Delepine M, Khalouf E, Gedde-Dahl T Jr, Van Maldergem L, Sobel E, Papp J, Meier M, Megarbane A, Bachy A <i>et al.</i> Identification of the gene altered in Berardinelli-Seip congenital lipodystrophy on chromosome 11q13. <i>Nature Genetics</i> 2001 28 365–370. (doi:10.1038/ng585)</p> <p>36 Payne VA, Grimsey N, Tuthill A, Virtue S, Gray SL, Dalla Nora E, Semple RK, O'Rahilly S & Rochford JJ. The human lipodystrophy gene BSCL2/seipin may be essential for normal adipocyte differentiation. <i>Diabetes</i> 2008 57 2055–2060. (doi:10.2337/db08-0184)</p> <p>37 Szymanski KM, Binns D, Bartz R, Grishin NV, Li WP, Agarwal AK, Garg A, Anderson RG & Goodman JM. The lipodystrophy protein seipin is found at endoplasmic reticulum lipid droplet junctions and is important for droplet morphology. <i>PNAS</i> 2007 104 20890–20895. (doi:10.1073/pnas.0704154104)</p> <p>38 Boutet E, El Mourabit H, Prot M, Nemani M, Khalouf E, Colard O, Maurice M, Durand-Schneider AM, Chretien Y, Gres S <i>et al.</i> Seipin deficiency alters fatty acid Delta9 desaturation and lipid droplet formation in Berardinelli-Seip congenital lipodystrophy. <i>Biochimie</i> 2009 91 796–803. (doi:10.1016/j.biochi.2009.01.011)</p> <p>39 Kim CA, Delepine M, Boutet E, El Mourabit H, Le Lay S, Meier M, Nemani M, Bridel E, Leite CC, Bertola DR <i>et al.</i> Association of a homozygous nonsense caveolin-1 mutation with Berardinelli-Seip congenital lipodystrophy. <i>Journal of Clinical Endocrinology and Metabolism</i> 2008 93 1129–1134. (doi:10.1210/jc.2007-1328)</p> <p>40 Hayashi YK, Matsuda C, Ogawa M, Goto K, Tominaga K, Mitsuhashi S, Park YE, Nonaka I, Hino-Fukuyo N, Haginoya K <i>et al.</i> Human PTRF mutations cause secondary deficiency of caveolins resulting in muscular dystrophy with generalized lipodystrophy. <i>Journal of Clinical Investigation</i> 2009 119 2623–2633. (doi:10.1172/JCI38660)</p> <p>41 Payne F, Lim K, Girousse A, Brown RJ, Kory N, Robbins A, Xue Y, Sleight A, Cochran E, Adams C <i>et al.</i> Mutations disrupting the Kennedy phosphatidylcholine pathway in humans with congenital lipodystrophy and fatty liver disease. <i>PNAS</i> 2014 111 8901–8906. (doi:10.1073/pnas.1408523111)</p> <p>42 Shackleton S, Lloyd DJ, Jackson SN, Evans R, Niermeijer MF, Singh BM, Schmidt H, Brabant G, Kumar S, Durrington PN <i>et al.</i> LMNA, encoding lamin A/C, is mutated in partial lipodystrophy. <i>Nature Genetics</i> 2000 24 153–156. (doi:10.1038/72807)</p> <p>43 Rankin J & Ellard S. The laminopathies: a clinical review. <i>Clinical Genetics</i> 2006 70 261–274. (doi:10.1111/j.1399-0004.2006.00677.x)</p> <p>44 Semple RK, Chatterjee VK & O'Rahilly S. PPAR γ and human metabolic disease. <i>Journal of Clinical Investigation</i> 2006 116 581–589. (doi:10.1172/JCI28003)</p> <p>45 Rubio-Cabezas O, Puri V, Murano I, Saudek V, Semple RK, Dash S, Hyden CS, Bottomley W, Vigouroux C, Magre J <i>et al.</i> Partial lipodystrophy and insulin resistant diabetes in a patient with a homozygous nonsense mutation in CIDEC. <i>EMBO Molecular Medicine</i> 2009 1 280–287. (doi:10.1002/emmm.200900037)</p> <p>46 Gandotra S, Le Dour C, Bottomley W, Cervera P, Giral P, Reznik Y, Charpentier G, Auclair M, Delepine M, Barroso I <i>et al.</i> Perilipin deficiency and autosomal dominant partial lipodystrophy. <i>New England Journal of Medicine</i> 2011 364 740–748. (doi:10.1056/NEJMoa1007487)</p> <p>47 Yamada K, Ikegami H, Yoneda H, Miki T & Ogihara T. All patients with Werner's syndrome are insulin resistant, but only those who also have impaired insulin secretion develop overt diabetes. <i>Diabetes Care</i> 1999 22 2094–2095. (doi:10.2337/diacare.22.12.2094)</p> <p>48 Imura H, Nakao Y, Kuzuya H, Okamoto M, Okamoto M & Yamada K. Clinical, endocrine and metabolic aspects of the Werner syndrome compared with those of normal aging. <i>Advances in Experimental Medicine and Biology</i> 1985 190 171–185. (doi:10.1007/978-1-4684-7853-2_6)</p> <p>49 Diaz A, Vogiatzi MG, Sanz MM & German J. Evaluation of short stature, carbohydrate metabolism and other endocrinopathies in Bloom's syndrome. <i>Hormone Research</i> 2006 66 111–117. (doi:10.1159/000093826)</p> <p>50 Payne F, Colnaghi R, Rocha N, Seth A, Harris J, Carpenter G, Bottomley WE, Wheeler E, Wong S, Saudek V <i>et al.</i> Hypomorphism in human NSMCE2 linked to primordial dwarfism and insulin resistance. <i>Journal of Clinical Investigation</i> 2014 124 4028–4038. (doi:10.1172/JCI73264)</p> <p>51 Weedon MN, Ellard S, Prindle MJ, Caswell R, Lango Allen H, Oram R, Godbole K, Yajnik CS, Sbraccia P, Novelli G <i>et al.</i> An in-frame deletion at the polymerase active site of POLD1 causes a multisystem disorder with lipodystrophy. <i>Nature Genetics</i> 2013 45 947–950. (doi:10.1038/ng.2670)</p> <p>52 Mayson SE, Parker VE, Schutta MH, Semple RK & Rickels MR. Severe insulin resistance and hypertriglyceridemia after childhood total body irradiation. <i>Endocrine Practice</i> 2013 19 51–58. (doi:10.4158/EP12115.OR)</p> <p>53 Ablamunits V, Weisberg SP, Lemieux JE, Combs TP & Klebanov S. Reduced adiposity in ob/ob mice following total body irradiation and bone marrow transplantation. <i>Obesity</i> 2007 15 1419–1429. (doi:10.1038/oby.2007.170)</p> <p>54 Marshall JD, Maffei P, Collin GB & Naggert JK. Alstrom syndrome: genetics and clinical overview. <i>Current Genomics</i> 2011 12 225–235. (doi:10.2174/138920211795677912)</p> <p>55 Huang-Doran I, Bicknell LS, Finucane FM, Rocha N, Porter KM, Tung YC, Szekeres F, Krook A, Nolan JJ, O'Driscoll M <i>et al.</i> Genetic defects in human pericentrin are associated with severe insulin resistance and diabetes. <i>Diabetes</i> 2011 60 925–935. (doi:10.2337/db10-1334)</p> <p>56 Rauch A, Thiel CT, Schindler D, Wick U, Crow YJ, Ekici AB, van Essen AJ, Goecke TO, Al-Gazali L, Chrzanowska KH <i>et al.</i> Mutations in the pericentrin (PCNT) gene cause primordial dwarfism. <i>Science</i> 2008 319 816–819. (doi:10.1126/science.1151174)</p>				

Review	R K Semple	How many types of insulin resistance?	174:5	R222
--------	------------	---------------------------------------	-------	------

- 57 Chen JH, Segni M, Payne F, Huang-Doran I, Sleigh A, Adams C, UK10K Consortium UK, Savage DB, O'Rahilly S, Semple RK *et al*. Truncation of POC1A associated with short stature and extreme insulin resistance. *Journal of Molecular Endocrinology* 2015 **55** 147–158. (doi:10.1530/JME-15-0090)
- 58 Gorden P, Lupsa BC, Chong AY & Lungu AO. Is there a human model for the 'metabolic syndrome' with a defined aetiology? *Diabetologia* 2010 **53** 1534–1536. (doi:10.1007/s00125-010-1719-7)
- 59 Virtue S & Vidal-Puig A. It's not how fat you are, it's what you do with it that counts. *PLoS Biology* 2008 **6** e237. (doi:10.1371/journal.pbio.0060237)
- 60 Shulman GI. Cellular mechanisms of insulin resistance. *Journal of Clinical Investigation* 2000 **106** 171–176. (doi:10.1172/JCI10583)
- 61 Danforth E Jr. Failure of adipocyte differentiation causes type II diabetes mellitus? *Nature Genetics* 2000 **26** 13. (doi:10.1038/79111)
- 62 Robbins AL & Savage DB. The genetics of lipid storage and human lipodystrophies. *Trends in Molecular Medicine* 2015 **21** 433–438. (doi:10.1016/j.molmed.2015.04.004)
- 63 Misra A, Peethambaram A & Garg A. Clinical features and metabolic and autoimmune derangements in acquired partial lipodystrophy: report of 35 cases and review of the literature. *Medicine* 2004 **83** 18–34. (doi:10.1097/01.md.0000111061.69212.59)
- 64 Cook JR & Semple RK. Hypoadiponectinemia – cause or consequence of human “insulin resistance”? *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 1544–1554. (doi:10.1210/jc.2009-2286)
- 65 Ye R & Scherer PE. Adiponectin, driver or passenger on the road to insulin sensitivity? *Molecular Metabolism* 2013 **2** 133–141. (doi:10.1016/j.molmet.2013.04.001)
- 66 Semple RK, Soos MA, Luan J, Mitchell CS, Wilson JC, Gurnell M, Cochran EK, Gorden P, Chatterjee VK, Wareham NJ *et al*. Elevated plasma adiponectin in humans with genetically defective insulin receptors. *Journal of Clinical Endocrinology and Metabolism* 2006 **91** 3219–3223. (doi:10.1210/jc.2006-0166)
- 67 Semple RK, Halberg NH, Burling K, Soos MA, Schraw T, Luan J, Cochran EK, Dunger DB, Wareham NJ, Scherer PE *et al*. Paradoxical elevation of high-molecular weight adiponectin in acquired extreme insulin resistance due to insulin receptor antibodies. *Diabetes* 2007 **56** 1712–1717. (doi:10.2337/db06-1665)
- 68 Semple RK, Cochran EK, Soos MA, Burling KA, Savage DB, Gorden P & O'Rahilly S. Plasma adiponectin as a marker of insulin receptor dysfunction: clinical utility in severe insulin resistance. *Diabetes Care* 2008 **31** 977–979. (doi:10.2337/dc07-2194)
- 69 Mente A, Meyre D, Lanktree MB, Heydarpour M, Davis AD, Miller R, Gerstein H, Hegele RA, Yusuf S, Anand SS *et al*. Causal relationship between adiponectin and metabolic traits: a Mendelian randomization study in a multiethnic population. *PLoS ONE* 2013 **8** e66808. (doi:10.1371/journal.pone.0066808)
- 70 Gao H, Fall T, van Dam RM, Flyvbjerg A, Zethelius B, Ingelsson E & Hagg S. Evidence of a causal relationship between adiponectin levels and insulin sensitivity: a Mendelian randomization study. *Diabetes* 2013 **62** 1338–1344. (doi:10.2337/db12-0935)
- 71 Yaghootkar H, Lamina C, Scott RA, Dastani Z, Hivert MF, Warren LL, Stancakova A, Buxbaum SG, Lyytikainen LP, Henneman P *et al*. Mendelian randomization studies do not support a causal role for reduced circulating adiponectin levels in insulin resistance and type 2 diabetes. *Diabetes* 2013 **62** 3589–3598. (doi:10.2337/db13-0128)
- 72 Bueno AC, Sun K, Martins CS, Elias Junior J, Miranda W, Tao C, Foss-Freitas MC, Barbieri MA, Bettiol H, de Castro M *et al*. A novel ADIPOQ mutation (p.M40K) impairs assembly of high-molecular-weight adiponectin and is associated with early-onset obesity and metabolic syndrome. *Journal of Clinical Endocrinology and Metabolism* 2014 **99** E683–E693. (doi:10.1210/jc.2013-3009)
- 73 Semple RK, Sleigh A, Murgatroyd PR, Adams CA, Bluck L, Jackson S, Vottero A, Kanabar D, Charlton-Menys V, Durrington P *et al*. Postreceptor insulin resistance contributes to human dyslipidemia and hepatic steatosis. *Journal of Clinical Investigation* 2009 **119** 315–322.
- 74 Lambert JE, Ramos-Roman MA, Browning JD & Parks EJ. Increased de novo lipogenesis is a distinct characteristic of individuals with nonalcoholic fatty liver disease. *Gastroenterology* 2014 **146** 726–735. (doi:10.1053/j.gastro.2013.11.049)
- 75 Donnelly KL, Smith CI, Schwarzenberg SJ, Jessurun J, Boldt MD & Parks EJ. Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. *Journal of Clinical Investigation* 2005 **115** 1343–1351. (doi:10.1172/JCI23621)
- 76 Brown MS & Goldstein JL. Selective versus total insulin resistance: a pathogenic paradox. *Cell Metabolism* 2008 **7** 95–96. (doi:10.1016/j.cmet.2007.12.009)
- 77 Wan M, Leavens KF, Saleh D, Easton RM, Guertin DA, Peterson TR, Kaestner KH, Sabatini DM & Birnbaum MJ. Postprandial hepatic lipid metabolism requires signaling through Akt2 independent of the transcription factors FoxA2, FoxO1, and SREBP1c. *Cell Metabolism* 2011 **14** 516–527. (doi:10.1016/j.cmet.2011.09.001)
- 78 Leavens KF, Easton RM, Shulman GI, Previs SF & Birnbaum MJ. Akt2 is required for hepatic lipid accumulation in models of insulin resistance. *Cell Metabolism* 2009 **10** 405–418. (doi:10.1016/j.cmet.2009.10.004)
- 79 Cook JR, Langlet F, Kido Y & Accili D. Pathogenesis of selective insulin resistance in isolated hepatocytes. *Journal of Biological Chemistry* 2015 **290** 13972–13980. (doi:10.1074/jbc.M115.638197)
- 80 Parker VE, Savage DB, O'Rahilly S & Semple RK. Mechanistic insights into insulin resistance in the genetic era. *Diabetic Medicine* 2011 **28** 1476–1486. (doi:10.1111/j.1464-5491.2011.03463.x)
- 81 Nandi A, Chen Z, Patel R & Poretsky L. Polycystic ovary syndrome. *Endocrinology and Metabolism Clinics of North America* 2014 **43** 123–147. (doi:10.1016/j.ecl.2013.10.003)
- 82 Goodarzi MO, Dumesic DA, Chazenbalk G & Azziz R. Polycystic ovary syndrome: etiology, pathogenesis and diagnosis. *Nature Reviews. Endocrinology* 2011 **7** 219–231. (doi:10.1038/nrendo.2010.217)
- 83 Dunaif A. Insulin resistance and the polycystic ovary syndrome: mechanism and implications for pathogenesis. *Endocrine Reviews* 1997 **18** 774–800. (doi:10.1210/er.18.6.774)
- 84 Poretsky L, Seto-Young D, Shrestha A, Dhillon S, Mirjany M, Liu HC, Yih MC & Rosenwaks Z. Phosphatidylinositol-3 kinase-independent insulin action pathway(s) in the human ovary. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 3115–3119. (doi:10.1210/jcem.86.7.7617)
- 85 O'Reilly M, Gathercole L, Capper F, Arlt W & Tomlinson J. Effect of insulin on AKR1C3 expression in female adipose tissue: *in-vivo* and *in-vitro* study of adipose androgen generation in polycystic ovary syndrome. *Lancet* 2015 **385** S16–950. (doi:10.1016/S0140-6736(15)60331-2)
- 86 Weber DR, Stancescu DE, Semple R, Holland C & Magge SN. Continuous subcutaneous IGF-1 therapy via insulin pump in a patient with Donohue syndrome. *Journal of Pediatric Endocrinology & Metabolism* 2014 **27** 1237–1241. (doi:10.1515/jpem-2013-0402)
- 87 Brisigotti M, Fabbretti G, Pesce F, Gatti R, Cohen A, Parenti G & Callea F. Congenital bilateral juvenile granulosa cell tumor of the ovary in leprechaunism: a case report. *Pediatric Pathology* 1993 **13** 549–558. (doi:10.3109/15513819309048242)
- 88 Stubbs SA, Webber LJ, Stark J, Rice S, Margara R, Lavery S, Trew GH, Hardy K & Franks S. Role of insulin-like growth factors in initiation of follicle growth in normal and polycystic human ovaries. *Journal of Clinical Endocrinology and Metabolism* 2013 **98** 3298–3305. (doi:10.1210/jc.2013-1378)
- 89 Poretsky L, Cataldo NA, Rosenwaks Z & Giudice LC. The insulin-related ovarian regulatory system in health and disease. *Endocrine Reviews* 1999 **20** 535–582. (doi:10.1210/edrv.20.4.0374)
- 90 Kaltsas GA, Androulakis II, Tziveriotis K, Papadogias D, Tsikini A, Makras P, Dimitriou K, Stathopoulou A & Piaditis G. Polycystic ovaries and the polycystic ovary syndrome phenotype in women with active acromegaly. *Clinical Endocrinology* 2007 **67** 917–922. (doi:10.1111/j.1365-2265.2007.02987.x)

Review	R K Semple	How many types of insulin resistance?	174:5	R223
--------	------------	---------------------------------------	-------	------

- 91 Arioglu E, Andewelt A, Diabo C, Bell M, Taylor SI & Gorden P. Clinical course of the syndrome of autoantibodies to the insulin receptor (type B insulin resistance): a 28-year perspective. *Medicine* 2002 **81** 87–100. (doi:10.1097/00005792-200203000-00001)
- 92 Idkowiak J, Lavery GG, Dhir V, Barrett TG, Stewart PM, Krone N & Arlt W. Premature adrenarche: novel lessons from early onset androgen excess. *European Journal of Endocrinology* 2011 **165** 189–207. (doi:10.1530/EJE-11-0223)
- 93 Giovannucci E, Harlan DM, Archer MC, Bergenstal RM, Gapstur SM, Habel LA, Pollak M, Regensteiner JG & Yee D. Diabetes and cancer: a consensus report. *Diabetes Care* 2010 **33** 1674–1685. (doi:10.2337/dc10-0666)
- 94 Nead KT, Sharp SJ, Thompson DJ, Painter JN, Savage DB, Semple RK, Barker A, Australian National Endometrial Cancer Study Group, Perry JR, Attia J *et al.* Evidence of a causal association between insulinemia and endometrial cancer: a mendelian randomization analysis. *Journal of the National Cancer Institute* 2015 **107** djv178. (doi:10.1093/jnci/djv178)
- 95 Safar Zadeh E, Lungu AO, Cochran EK, Brown RJ, Ghany MG, Heller T, Kleiner DE & Gorden P. The liver diseases of lipodystrophy: the long-term effect of leptin treatment. *Journal of Hepatology* 2013 **59** 131–137. (doi:10.1016/j.jhep.2013.02.007)
- 96 Belfiore A & Malaguarnera R. Insulin receptor and cancer. *Endocrine-Related Cancer* 2011 **18** R125–R147. (doi:10.1530/ERC-11-0074)

Received 16 November 2015

Revised version received 1 February 2016

Accepted 9 February 2016