From pseudohypoparathyroidism to inactivating PTH/PTHrP signalling disorder (iPPSD), a novel classification proposed by the EuroPHP network

Susanne Thiele¹, Giovanna Mantovani², Anne Barlier³, Valentina Boldrin², Paolo Bordogna², Luisa De Sanctis⁴, Francesca M Elli², Kathleen Freson⁵, Intza Garin⁶, Virginie Grybek^{7,8}, Patrick Hanna^{7,8}, Benedetta Izzi⁵, Olaf Hiort¹, Beatriz Lecumberri⁹, Arrate Pereda^{6,10}, Vrinda Saraff¹¹, Caroline Silve^{7,8,12}, Serap Turan¹³, Alessia Usardi^{7,14}, Ralf Werner¹, Guiomar Perez de Nanclares⁶ and Agnès Linglart^{7,8,14}

¹Division of Experimental Pediatric Endocrinology and Diabetes, Department of Pediatrics, University of Lübeck, Lübeck, Germany, ²Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Endocrinology and Diabetology Unit, Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy, ³APHM, Hôpital la Conception, Laboratory of Molecular Biology, Marseille, France, ⁴Department of Public Health and Pediatric Sciences, University of Torino, Torino, Italy, ⁵Department of Cardiovascular Sciences, Center for Molecular and Vascular Biology, University of Leuven, Leuven, Belgium, ⁶Molecular (Epi)Genetics Laboratory, BioAraba National Health Institute, OSI Araba University Hospital, Vitoria-Gasteiz, Spain, ⁷APHP, Reference Center for rare disorders of the Calcium and Phosphate Metabolism, filière OSCAR and Plateforme d'Expertise Maladies Rares Paris-Sud, Hôpital Bicêtre Paris Sud, Le Kremlin Bicêtre, France, ⁸INSERM U1169, Hôpital Bicêtre, Le Kremlin Bicêtre, et Université Paris-Saclay, Le Kremlin Bicêtre, France, ⁹Department of Endocrinology and Nutrition, La Paz University Hospital, Madrid, Spain, ¹⁰Department of Biochemistry and Molecular Biology, University of Basque Country, Leioa, Spain, ¹¹Department of Endocrinology and Diabetes, Birmingham Children's Hospital, Birmingham, UK, ¹²APHP, Service de Biochimie et Génétique Moléculaires, Hôpital Cochin, Paris, France, ¹³Department of Pediatrics, Division of Endocrinology and Diabetes, Marmara University, Istanbul, Turkey, ¹⁴APHP, Department of Paediatric Endocrinology and Diabetology, Bicêtre Paris Sud hospital, Le Kremlin Bicêtre, France

Correspondence
should be addressed
to A Linglart
Email
agnes.linglart@aphp.fr

Abstract

Objective: Disorders caused by impairments in the parathyroid hormone (PTH) signalling pathway are historically classified under the term pseudohypoparathyroidism (PHP), which encompasses rare, related and highly heterogeneous diseases with demonstrated (epi)genetic causes. The actual classification is based on the presence or absence of specific clinical and biochemical signs together with an *in vivo* response to exogenous PTH and the results of an *in vitro* assay to measure Gsa protein activity. However, this classification disregards other related diseases such as acrodysostosis (ACRDYS) or progressive osseous heteroplasia (POH), as well as recent findings of clinical and genetic/epigenetic background of the different subtypes. Therefore, the EuroPHP network decided to develop a new classification that encompasses all disorders with impairments in PTH and/or PTHrP cAMP-mediated pathway.

Design and methods: Extensive review of the literature was performed. Several meetings were organised to discuss about a new, more effective and accurate way to describe disorders caused by abnormalities of the PTH/PTHrP signalling pathway.

Results and conclusions: After determining the major and minor criteria to be considered for the diagnosis of these disorders, we proposed to group them under the term 'inactivating PTH/PTHrP signalling disorder' (iPPSD). This terminology: (i) defines the common mechanism responsible for all diseases; (ii) does not require a confirmed genetic defect; (iii) avoids ambiguous terms like 'pseudo' and (iv) eliminates the clinical or molecular overlap

between diseases. We believe that the use of this nomenclature and classification will facilitate the development of rationale and comprehensive international guidelines for the diagnosis and treatment of iPPSDs.

*European Journal of
Endocrinology*
(2016) **175**, P1–P17

Introduction

Pseudohypoparathyroidism (PHP) encompasses a group of rare, related, highly heterogeneous and deeply impairing disorders characterised by end-organ resistance to the action of parathyroid hormone (PTH) and in most instances associated with a demonstrated (epi)genetic component (1, 2, 3). PHP is historically the first hormone-resistance syndrome described by Albright *et al.* (4).

A better understanding of the PHP pathophysiology followed the identification of the PTH receptor (PTH1R)

and its signal transduction pathway (Fig. 1) (5, 6). PTH1R, through its activation by two ligands, the PTH and the PTH-related peptide (PTHrP), regulates skeletal development, bone turnover and mineral ion homeostasis. In the kidney, binding of PTH to PTH1R stimulates the production of 1,25-dihydroxy vitamin D₃, and inhibits phosphate reabsorption in the proximal tubule, while it increases calcium reabsorption in the distal nephron. In the growth plate,

Figure 1

Schematic transduction of PTH1R/Gsa/cAMP/PKA pathway. Upon ligand binding (PTH or PTHrP is mentioned in the figure), the receptor (PTH1R) activates the G protein. Then, the Gsa subunit triggers the activation of the adenylate cyclase leading to cAMP synthesis. cAMP binds to the regulatory 1A subunits (R1A) of the PKA, the most common effector of cAMP. Upon cAMP (grey diamonds) binding, the catalytic subunits (Cat) dissociate from the R1A subunits, and phosphorylate numerous target proteins including CREB (cAMP-responsive binding elements) and the phosphodiesterases (PDEs). CREB activates the transcription of cAMP-responsive genes. Intracellular cAMP is then deactivated by PDEs, among which are PDE4D and PDE3A. PTH1R: transmembrane convoluted black line; G protein: trimer α , β , γ ; cAMP: grey diamond; PKA: tetramer R1A (regulatory subunit 1A) and Cat (catalytic subunit); phosphodiesterases: ovals PDE4D or PDE3A; DNA: scale bar.

Position Statement	S Thiele and others	iPPSD, a novel classification for PHP	175:6	P3
--------------------	---------------------	---------------------------------------	-------	----

PTHrP promotes endochondral ossification, by binding to PTH1R (7).

The Blomstrand chondrodysplasia (OMIM #215045), a lethal form of dwarfism (8), was the first disorder associated with biallelic loss-of-function mutations of the *PTH1R* gene (9). Subsequently, one report has described a milder phenotype in living children affected with Eiken disease (OMIM #600002), short stature, elevated PTH and mutations of *PTH1R* (10, 11).

A defect in the response of the proximal renal tubule to PTH is the hallmark of all forms of PHP. It manifests as hypocalcaemia, hyperphosphataemia and elevated circulating levels of PTH in the absence of vitamin D deficiency (5, 7, 12).

PTH receptor couples with the stimulatory G protein (Gsa), leading to cAMP formation. Renal tubular response to exogenously administered PTH through measurement of serum and urinary cAMP levels permits the differentiation of PHP type 1 (PHP1), in which a blunted cAMP response is observed, from PHP type 2 (PHP2), where cAMP increase is conserved but the phosphaturic response is deficient (13). To date, only a handful of PHP2 cases have been reported, and the molecular defect responsible for this variant is still unknown. It has also been hypothesised that PHP2 could either be an acquired defect secondary to vitamin D deficiency (14), as calcium and vitamin D supplementation resulted in normalisation of the phosphaturic response to PTH in some patients (14, 15), or due to defects downstream the Gsa protein, as seen in patients with acrodysostosis type 1 (ACRDYS1) (16).

In 1980, deficiency in the Gsa protein activity in erythrocytes extracted from patients affected with PHP1 was demonstrated *in vitro* (17, 18). For years, this bioassay allowed the diagnosis of PHP, and contributed to PHP subclassification (see below).

PHP type 1 (PHP1) is further subdivided based on the presence (PHP1A and PHP1C; OMIM #103580 and #612462 respectively) (6, 17, 18, 19) or absence (PHP1B; OMIM #603233) (6, 20) of Albright hereditary osteodystrophy (AHO) (Table 1). AHO is a clinical entity initially described together with PHP in 1942, which encompasses heterogeneous clinical findings such as brachydactyly, rounded face, short stature, stocky build and subcutaneous ossifications (4, 21, 22). Additional features that may not directly relate to AHO, yet extensively associated with PHP1A individuals, include obesity, varying degrees of intellectual disability and resistance to several hormones, including TSH, GHRH and calcitonin (23, 24, 25, 26, 27, 28). The subcategory of PHP1C has all the characteristics of PHP1A, except that

Gsa activity in erythrocytes was found comparable to controls (29, 30).

Interestingly, patients showing the physical features of AHO without any evidence of PTH resistance were also described by Albright *et al.* (21) 10 years after their first report of PHP. This new syndrome, named pseudopseudohypoparathyroidism (PPHP; OMIM #612463) may be present either in kindreds with PHP or as an isolated defect. It is possible that the 'bone phenotype' observed in AHO is largely mediated by the resistance to PTHrP at the growth plate during foetal and postnatal growth (31).

In 1990, the first heterozygous inactivating mutation in the gene coding for Gsa (*GNAS*), responsible for PHP1A, was described (32). Since then, several Gsa-coding mutations have been identified in all of its 13 exons with different frequency, with a detection rate of about 70% (33, 34, 35, 36, 37, 38, 39). Cases of deletions of 20q, including part or the whole *GNAS* gene, and an inversion at *GNAS* have been recently reported (40, 41, 42, 43, 44). Remarkably, similar mutations when paternally inherited, or occurring *de novo* on the paternal allele of *GNAS* may lead to PPHP or to progressive osseous heteroplasia (POH, OMIM #166350), a disorder characterised by heterotopic ossifications expanding into deep muscles and connective tissues (45, 46).

GNAS is a locus encoding several transcripts through alternative splicing. In most tissues, except for Gsa, the *GNAS* transcripts are of monoallelic origin due to the control of their expression by parent-specific differentially methylated regions (DMRs) (Fig. 2) (47). In thyroid, pituitary gland and most likely in the proximal tubule (36), Gsa is predominantly expressed from the maternal allele through a yet unexplained mechanism (48, 49). In the early 2000, the molecular defect of PHP1B was characterised. The most consistent defect common to all PHP1B patients is a paternal-specific pattern of cytosine methylation within the maternal *GNAS* A/B: transcriptional start site (TSS)-DMR (*GNAS* A/B:TSS-DMR; previously known as exon A/B or 1A), which could lead to a decreased expression of Gsa in the renal proximal tubules, hence PTH resistance (50). Fifteen to twenty percent of the PHP1B cases present familial history with an autosomal dominant mode of inheritance (AD-PHP1B) through the maternal lineage. Most AD-PHP1B show loss of imprinting (LOI) limited to the *GNAS* A/B:TSS-DMR (more precisely a loss of methylation (LOM)) associated with deletions on the maternal allele of *cis*-acting control elements within *STX16* or *NESP55* (51, 52, 53, 54, 55), although other

Table 1 Former classification of PHP along with the other disorders affecting the PTH/PTHrP signalling pathway; note the overlap of phenotypes and molecular defects of the patients. Diseases included in the former classification are PHP1A, PHP1B, PHP1C and PPHP.

	PHP1A	PHP1C	PHP1B					
Clinical presentation	AHO	AHO	No AHO	AHO in some patients (brachydactyly, subcutaneous ossification) and/or obesity	AHO in very few patients	Mental retardation reported in 2 patients, lambdoid synostosis, early-onset obesity, macrocephaly	BWS	No AHO
	Obesity	Obesity	No obesity		Obesity may be present		Obesity	No obesity
	Cognitive impairment Subcutaneous ossifications	Cognitive impairment Subcutaneous ossifications	No cognitive impairment					
Hormone resistance	Resistance to PTH, TSH, GHRH, calcitonin, epinephrine, glucagon and gonadotropins	Resistance to PTH, TSH, epinephrine and gonadotropins	PTH resistance	PTH resistance, ± TSH resistance	PTH resistance, ± TSH resistance	PTH resistance, ± TSH resistance	PTH resistance	PTH resistance, ± TSH resistance
<i>In vitro</i> activity of Gsa	Significantly below controls	Similar to controls	Similar to controls	Mildly decreased when compared with controls	Mildly decreased when compared with controls			Similar to controls
LOI at the <i>GNAS</i> DMRs				LOM at the <i>GNAS</i> A/B:TSS-DMR	Broad LOI	Broad LOI	Broad LOI	Broad LOI
Genetic lesion	Heterozygous mutation in the coding sequence of <i>GNAS</i> (maternal allele)	Heterozygous mutation in the coding sequence of <i>GNAS</i> (p.E392K, p.E392X, p.L388R and p.Y391X, all in exon 13) (maternal allele)	Heterozygous mutation in the coding sequence of <i>GNAS</i> (p.Ile382del) (maternal allele)	Recurrent 3-kb <i>STX16</i> deletion or 4.2-kb deletion of <i>STX16</i>	Unknown	UPD(20)pat including <i>GNAS</i>	MLID	Maternal deletion of <i>NESP</i> and/or <i>AS</i> or duplication of <i>GNAS</i>
References	(32, 33, 34, 35, 36, 37, 38, 39)	(29, 30, 78)	(112)	(50, 51, 52, 113)	(50, 51, 63, 74)	(59, 60, 61, 62, 63)	(114)	(53, 54, 58, 115)

Position Statement	S Thiele and others	iPPSD, a novel classification for PHP	175:6	P5
---------------------------	---------------------	---------------------------------------	-------	-----------

PPHP		POH	2q37.3 Deletion Syndrome	PHP2	Acrodysostosis			Blomstrand dysplasia	Eiken disease
AHO	AHO	Subcutaneous ossifications	AHO	No AHO	Severe AHO	AHO	Severe AHO	Lethal dwarfism	Epiphyseal dysplasia
Subcutaneous ossifications	Subcutaneous ossifications		Cognitive impairment	Hypocalcaemia, osteomalacia	Cognitive impairment in some patients		Hypertension		Short stature
No	Mild	No	No	PTH resistance	PTH resistance, and TSH in some patients	PTH resistance, and TSH in some patients	No		Elevated PTH in one patient
Significantly below controls	Significantly below controls								
Heterozygous mutation in the coding sequence of <i>GNAS</i> (paternal allele)	Heterozygous mutation in the coding sequence of <i>GNAS</i> (paternal allele)	Heterozygous mutation in the coding sequence of <i>GNAS</i> (paternal allele) or no mutation identified	Deletion of the 2q37.3 chromosomal region including <i>HDAC4</i>	None	Heterozygous mutation in the coding sequence of <i>PRKAR1A</i> or <i>PDE4D</i>	Heterozygous mutation in the coding sequence of <i>PRKAR1A</i>	Heterozygous mutation in the coding sequence of <i>PDE3A</i>	Biallelic inactivating mutation in the coding sequence of <i>PTH1R</i>	Biallelic inactivating mutation in the coding sequence of <i>PTH1R</i>
(34, 36, 44, 100)	(77)	(39, 45, 46, 79, 80)	(116)	(14)	(16, 82, 90, 91, 117, 118, 119)	(90, 91)	(92)	(9, 11)	(10)

AHO, Albright's hereditary osteodystrophy; BWS, Beckwith–Wiedemann syndrome; MLID, multilocus imprinting defect; NA, not available; PHP, pseudohypoparathyroidism; PPHP, pseudopseudohypoparathyroidism.

Figure 2

The imprinted human *GNAS* locus (Hg19-chr20:57,414,795-57,486,250), on chromosome 20, close to the *STX16* gene (Hg19-chr20:57,226,309-57,254,5812) (source UCSC, Hg19). The centromeric/telomeric orientation of the chromosome is indicated. The maternal (NESP), paternal (AB, AS and XL) and biallelic (Gsa) transcripts are depicted as arrows. Maternal- and paternal-expressed transcripts are drawn above and below the horizontal line respectively. Black boxes: coding exons; grey boxes: noncoding exons; arrows: transcription (direction and parental origin). The brackets delimit the imprinting control element deletions, which have been reported. *STX16* gene: full brackets: the recurrent *STX16* deletion of 3.3 kb (38); large dotted brackets: the *STX16* deletion of 4.4 kb (39); small dotted brackets: the *STX16* deletion of 29.5 kb (42). *GNAS* locus: full brackets: the 4.7 and 4 kb deletions removing the *NESP* exon and exons 3 and 4 of *GNAS-AS1* (40); large dotted brackets: the 4.2 kb deletion removing exons 3 and 4 of *GNAS-AS1* (43); deletions of 40 pb (*) and 33 pb (#) in introns of *NESP* and *GNAS-AS1* (44); small dotted brackets: the *NESP* and *GNAS-AS1* deletion (41). cen, centromeric; Mat, maternal; Pat, paternal; tel, telomeric.

maternally inherited deletions have been identified affecting all four DMRs (*GNAS-NESP*:TSS-DMR, *GNAS-AS1*:TSS-DMR, *GNAS-XL*:Ex1-DMR and *GNAS A/B*:TSS-DMR) (56, 57, 58).

The remaining cases of PHP1B are sporadic. They present with broad LOI at *GNAS*, including the *GNAS A/B*:TSS-DMR. The molecular basis of this broad LOI is yet to be identified, with an exception of less than 10% of the patients who are affected by paternal complete or segmental uniparental disomy (UPD) of the chromosome 20, comprising the *GNAS* locus (59, 60, 61, 62, 63).

To summarise, the existing classification of PHP (Table 1) is based on the following criteria: (i) presence or absence of AHO differentiates PHP1A/PHP1C from PHP1B; (ii) presence or absence of hormonal resistance differentiates PHP1 from PPHP; (iii) *in vivo* response to exogenous PTH as for nephrogenic cAMP synthesis and phosphaturia separates PHP1 from PHP2 and (iv) *in vitro* assay measuring the Gsa protein activity from erythrocyte membranes differentiates between PHP1A and PHP1C.

As described above, the existing PHP classification does not include molecular defect as a criterion and fails to stratify PHP and AHO as well as include conditions such as acrodysostosis, POH and PTH1R-related chondrodysplasia. In this manuscript, we therefore propose to review the rationale of this nomenclature and recommend a novel classification for disorders impairing the PTH/PTHrP signalling pathway.

Methodology

The EuroPHP network met on three different occasions (October 2014, May 2015, November 2015) to discuss and agree on a novel classification. The aims of these meetings were (i) to identify the limitations in the current PHP classification; (ii) to formulate mandatory criteria for the new classification; (iii) to propose a comprehensive definition gathering all the disorders; (iv) to analyse the classifications used in other genetic/epigenetic conditions and (v) to generate a novel classification. The methodology comprised a thorough review of the current literature to facilitate comparison and form basis for the proposal of a new classification.

We have carefully considered a series of classifications proposed for various rare genetic/epigenetic disorders, including the reporting manuscripts that were taken into consideration for the design of a novel classification (summarised in Table 2). In brief, methodologies were similar. A group of experts in the field identified the deficiencies in the existing classification/terminology and the need for an update. Subsequently, agreement on a novel terminology and classification was reached and reported (64, 65, 66, 67, 68, 69, 70).

Challenges and limitations of the current classification

Recent clinical and molecular data gathered for these complex disorders have questioned the distinction

Position Statement	S Thiele and others	iPPSD, a novel classification for PHP	175:6	P7
---------------------------	---------------------	---------------------------------------	-------	-----------

Table 2 Nonexhaustive review of classifications used in other conditions.

	Methodology used to build the classification	Mode of classification	Advantages	Limitations
Primary immunodeficiency diseases (69)	2-days meeting	Groups of diseases according to the most fundamental defect presented as a table format	Allows a practical clinical framework for PID diagnosis	The complexities of these conditions cannot easily be captured in the limited table format
Skeletal dysplasia (66)	Meeting, extensive review of the literature, and circulation of drafts of the manuscript	Groups of diseases defined by molecular, biochemical and/or radiographic criteria	Disorders are caused by disturbances in related metabolic pathways or gene networks, Sheer number of conditions included	The 'hybrid' nature of the classification, not clinical, not molecular
Autosomal dominant tubule interstitial kidney disease (67)	Meeting, agreement on the manuscript	Agreement on a novel name: ADTKD Classification based on the underlying genetic defect: ADTKD-gene	Provide information on the disease	Use in communication with patients may not be easy
Endocrine diseases (68)	Literature review	Groups of diseases by organ		
Diabetes mellitus (MODY) (70)	Meetings, agreement on the manuscript	Groups of diabetes by diseases' mechanism	Provide information on the disease mechanism Allow numbering of new diabetes after identification of new genes for MODY (MODY1, MODY2, MODY3, MODY4...)	Very large groups of disease (type 2 diabetes for example)
Osteogenesis imperfecta (64)	Literature review	Phenotypes on evolution, radiology, clinics and genetics	Provide information on the disease mechanism and genetics	Confusing as one causing gene may be in different categories

of the different PHP and AHO subtypes in the existing classification (Table 1). We have selected the following limits of the current classification:

1. In a subset of patients with PHP1A and varying degree of AHO, LOI of *GNAS* identical to that of PHP1B has been reported, suggesting a molecular and clinical overlap between the two subtypes (71), further confirmed (72, 73, 74, 75).
2. PHP1B patients present with a moderate reduction in Gsa activity in erythrocyte membranes, reminiscent – yet less severe – to that observed in patients with PHP1A and PPHP (76).
3. Recently, mild resistance to PTH was described in patients affected with PPHP, carrying a paternal *GNAS* mutation (77), showing that the hormonal resistance is not only associated with maternally inherited *GNAS* mutations.
4. Different molecular defects have been identified in patients with PHP1C, i.e. LOI at *GNAS* and four loss-of-function mutations in the *GNAS* carboxyl-terminus leading to a conserved adenylyl cyclase receptor-independent activation but disrupted receptor-mediated activation (29, 30, 78).
5. Paternal *GNAS* mutations associated with progressive osseous heteroplasia are usually truncating mutations (79), yet they are identical to those found in families with PHP1A and/or PPHP (45). Also noteworthy is that a fraction of POH patients exhibits some of the typical AHO features and, conversely, some PHP1A patients carrying mutations on the maternal allele present with progressive deepening heterotopic ossifications. The hypothesis that POH should be considered as a form of PPHP is, therefore, debated (80, 81).
6. Heterozygous mutations in *PRKARIA* – coding for the regulatory subunit of the protein kinase A (PKA) – and *PDE4D* – coding for phosphodiesterase type 4 – have been found in patients with acrodysostosis (16, 82, 83). Acrodysostosis refers to a heterogeneous group of rare diseases characterised by skeletal dysplasia and characteristic features, including brachydactyly, facial dysmorphism and, in some cases, mental retardation (84, 85, 86, 87, 88). Acrodysostosis differs from PHP by more generalised osseous abnormalities (87, 89). Resistance to PTH and/or TSH is present in about 60–70% of acrodysostosis patients with a *PRKARIA* mutation, while, in case of a *PDE4D* mutation, such hormone resistances are found only in a smaller

Position Statement	S Thiele and others	iPPSD, a novel classification for PHP	175:6	P8
--------------------	---------------------	---------------------------------------	-------	----

subset of 10–20%. Interestingly, few patients bearing a *PRKARIA* mutation have been described in patients with a phenotype indistinguishable from PHP1A (90, 91).

- Heterozygous mutations in *PDE3A* have been identified in patients affected with hypertension and brachydactyly type E (hypertension and brachydactyly syndrome (HTNB): OMIM #112410) (92).
- Disorders associated with an impaired function of *PTH1R*, i.e. the Blomstrand and Eiken skeletal dysplasia, are currently not included in the classification.

Over the past two decades, it became obvious that clinical features such as AHO or *in vitro* assays such as Gsa bioactivity fail to differentiate between PHP subtypes. In addition, mutations of genes different from *GNAS* have been shown to lead to PTH and/or PTHrP resistance and *GNAS* mutations might trigger diseases different from PHP/PPHP (i.e. POH). These disorders are not encompassed by the current classification system.

For all these reasons, different independent studies from the authors of the present paper, as well as the 'EuroPHP network' concluded and agreed that a uniform terminology is required to create a functional working classification that reflects the current knowledge of the diseases (29, 93, 94).

Terminology

We propose the term of 'inactivating PTH/PTHrP signalling disorder', abbreviated as iPPSD, which encompass all disorders related to this pathway. We also propose that numbering will allow for both clinical features and molecular and genetic findings to be included. The advantages of this terminology are as follows: (i) it describes the common mechanism responsible for the diseases; (ii) it does not require a confirmed genetic defect; (iii) it avoids the ambiguous term like 'pseudo'; (iv) it eliminates the clinical or molecular overlap between diseases and (v) it is flexible to incorporate new evolving information.

We recognise that the nomenclature 'inactivating PTH/PTHrP signalling disorder' might be initially difficult for patients and caregivers to remember. It would, therefore, be helpful to rely on the abbreviation iPPSD. Equally, the former terms 'pseudohypoparathyroidism' and 'pseudopseudohypoparathyroidism' were also long and challenging to use for communication. PTH/PTHrP-specific pathway was deliberately included in the name of the classification to avoid the misperception with disorders resulting from the inactivation of G protein-coupled receptors, i.e. inactivating mutations in the TSH

receptor or in the FSH receptor. All nomenclature based on the cAMP signalling were carefully considered and rejected due to their generic nature.

Identification of mandatory criteria for the new classification

Basis for the newly proposed classification of iPPSD are:

- to provide patients with an unambiguous diagnosis;
- to base nomenclature on pathophysiology, i.e. the PTH1R/Gsa/cAMP/PKA pathway, and a standardised diagnostic pathway;
- to formulate basis to develop new therapeutic approaches;
- to be sufficiently flexible and adaptable to include emerging clinical and molecular information;
- to be simple and usable for the caregivers.

It is, therefore, of significant importance to define the category of iPPSD a patient belongs to, based on the characterisation of clinical/biochemical criteria, to facilitate a definitive diagnosis and, if possible, through molecular analysis, a more specific denomination within the classification.

We suggest three key clinical features as major criteria for the diagnosis of iPPSD. The proposed major criteria have minimum or no overlap with other conditions due to different mechanisms (Table 3, especially for the differential diagnoses).

We also propose a list of minor criteria that are associated with iPPSD. These are less specific to iPPSD compared with major criteria and can occur in other clinical conditions. Therefore, minor criteria need to be combined with one or more major criteria to establish the diagnosis of iPPSD.

Major criteria

PTH resistance

The hallmark of inactivating PTH/PTHrP signalling disorders is the resistance of the renal proximal tubule to the action of PTH. All genetic defects leading to a deficient PTH1R signalling in the kidney will, therefore, be named iPPSD.

Ectopic ossifications

Ectopic ossifications are superficial, subcutaneous nodules, defined as ectopic bone formation in the adipose or dermal tissue. Progressive osseous heterotopic calcifications often begin in the dermal and subcutaneous tissues and later

Position Statement	S Thiele and others	iPPSD, a novel classification for PHP	175:6	P9
---------------------------	----------------------------	--	--------------	-----------

Table 3 Definition of major and minor criteria for iPPSD and differential diagnoses.

		Assessment	Differential diagnosis	References
I. Major criteria	1. PTH resistance	Ionized calcium, total calcium Phosphate Magnesium PTH Vitamin D (25OHD) Creatinine Urinary calcium Urinary phosphate PTH infusion test in challenging cases	Normocalcaemic hyperparathyroidism Renal failure Vitamin D deficiency or any kind of secondary hyperparathyroidism	(16)
	2. Ectopic ossification	Detailed physical exam X-rays	Fibrodysplasia ossificans progressiva (FOP, OMIM #135100), post-traumatic osteoma cutis	
	3. Brachydactyly type E (comprises the IV)	Clinical inspection (fist), hand and feet X-rays	Turner syndrome, tricho-rhino-phalangeal syndrome (TRPS), TRPS I, (OMIM #190350), TRPS-II (OMIM #150230) and TRPS-III, (OMIM #190351)	
II. Minor criteria	1. TSH resistance	TSH, T4I, antibodies, imaging [†]	Mutations in the TSH receptor	(26, 27)
	2. Other hormonal resistances	IGF-1 (GH stimulation test if necessary), calcitonin, LH, FSH, GnRH test		(2, 27, 78, 98, 99, 100, 101)
	3. Motor and cognitive retardation or impairment	Computed tomography scan and/or MRI of the brain, psychopathological rating scales adjusted for age		(24, 25, 34, 85, 86, 102, 116)
	4. Intrauterine and postnatal growth retardation	IUGR: gestational age, birth weight, birth length, head circumference, comparison to reference charts; post-natal growth: growth charts, X-ray of the left hand for determination of the bone age		(16, 40, 92, 103, 104, 120)
	5. Obesity/overweight	Weight SDS, BMI percentile, BMI z-score		(23, 105, 106)
	6. Flat nasal bridge and/or maxillary hypoplasia and/or round face	Clinical inspection		(4, 84, 86, 90)
iPPSD clinical diagnosis	(a) Presence of one major criteria, either number 1 or 2; (b) Presence of major criteria number 3 and at least 2 minor criteria [‡]			

[†]US in adults with hypothyroidism and no evidence for autoimmunity; thyroid imaging through thyroid scintigraphy and US in neonates diagnosed through screening for congenital hypothyroidism; [‡]Minor criteria are nonspecific (obesity/cognitive impairment); for instance, the association of BDE+obesity or BDE+cognitive impairment would not be relevant for our classification. By raising the number of minor criteria from 1 to 2, we will reduce the risk of overdiagnosing patients with iPPSD.

progress to the deeper tissues, such as muscles and tendons. In children, ectopic ossifications are highly suggestive of an inactivating *GNAS* mutation, i.e. iPPSD.

Brachydactyly

Brachydactyly refers to shortening of fingers, toes or both. Brachydactyly type E (BDE, OMIM #113300) encompasses

variable shortening of the metacarpals/metatarsals, often with the involvement of phalanges (Fig. 3). It can either present in isolation or as part of a genetic disorder, most of which are included among iPPSD (95).

Brachydactyly can be challenging to identify in early childhood, and tends to become more evident during early puberty. Brachydactyly can be overlooked when all bones are short as in acrodisostosis since early childhood.

Position Statement	S Thiele and others	iPPSD, a novel classification for PHP	175:6	P10
---------------------------	---------------------	---------------------------------------	-------	------------

Figure 3

Patterns of brachydactyly type E associated with iPPSD. A, B, C, D and E, brachydactylies associated with coding mutations in the Gsa subunit of the G protein (iPPSD2). F and G, bone phenotype associated with the loss of imprinting at the *GNAS* locus (iPPSD3). H, I and J, brachydactylies associated with the molecular defect in *PRKAR1A* (iPPSD4) and *PDE4D* (iPPSD5). Note the phenotypic overlap between A, H, J and B, C, G, I respectively.

While PTH resistance and ectopic ossifications are considered major criteria for iPPSD, brachydactyly is less specific and should, therefore, be combined with at least one major or two minor criteria to trigger the diagnosis of iPPSD.

Minor criteria

Thyroid-stimulating hormone (TSH) resistance

In iPPSD, TSH resistance is often mild and characterised by elevated TSH levels associated with free thyroxine (T_4) levels in a normal or low-normal reference range. This occurs in the absence of goitre and markers of autoimmune disease (26, 27). TSH resistance can sometimes be the first detected sign of iPPSD,

especially in countries where screening for congenital hypothyroidism is routinely performed (96).

Other hormone resistances

Very few other hormone resistances have been demonstrated so far. Resistance to growth hormone-releasing hormone (GHRH), leading to growth hormone deficiency, is the most frequent additional resistance found in PHP1A, affecting as many as 60% of patients (97, 98, 99). Calcitonin resistance has been described without clinical features in patients affected with PHP1A (27). Elevated follicular-stimulating hormone (FSH) and luteinizing hormone (LH) levels were reported both by us and Namnoum *et al.* (78, 100). Glucagon and adrenaline resistances were demonstrated in patients

with features of PHP and low Gsa bioactivity through *in vivo* testing (6, 101).

Motor and cognitive retardation or impairment

Psychomotor and cognitive alterations have been described in about 40 to 70% of the patients with a maternal coding mutation of *GNAS* (25, 34), as well as in some patients affected with acrodysostosis (83, 85, 86). Psychiatric manifestations have also been reported in these patients (102). Patients with paternal mutations of *GNAS* or epigenetic modifications of the *GNAS* DMRs seem unaffected (25, 63).

Intrauterine and postnatal growth retardation

Intrauterine growth retardation (IUGR) has been frequently observed in both maternal and paternal inherited inactivating *GNAS* coding mutations. However, IUGR is more pronounced in patients harbouring mutations on the paternal *GNAS* allele, mainly when affecting *GNAS* exon 2–13 mutations, compared with patients with *GNAS* exon 1/intron 1 mutations (103). IUGR has also been described in acrodysostosis with mutations in *PRKARIA* or *PDE4D*, and in patients with mutations in *PDE3A* (16, 82, 90, 92). A LOI at the maternal *GNAS* A/B: TSS-DMR has been associated with increased intrauterine growth (104).

Postnatal growth retardation is a frequent sign in PHP1A and acrodysostosis. Growth hormone deficiency and premature closure of the epiphysis result in short stature (16, 82, 97, 105). Growth retardation has also been observed in PHP1B, although only in exceptional cases (71, 74), and in patients with Eiken dysplasia (10).

Obesity/overweight

Obesity or overweight may be the most nonspecific minor sign; however, it occurs very frequently in disorders with an impaired PTH/PTHrP signalling pathway and may help to differentiate between the different subtypes. Growth hormone deficiency, impaired lipolytic response of adrenaline (101) or decreased resting energy expenditure (106) contribute to the development of obesity in patients with mutations on the maternal allele of *GNAS* (23, 107). Obesity is also a frequent feature in patients affected with acrodysostosis (16, 90, 108).

Flat nasal bridge and/or maxillar hypoplasia and/or round face

Elements of facial dysmorphism have been associated with acrodysostosis (flat nasal bridge and/or maxillar hypoplasia) or with PHP1A (round face) (4, 86).

Diagnosis of iPPSD

We propose that a minimum of one of the major criteria is mandatory for the clinical diagnosis of iPPSD. PTH resistance or ectopic ossifications may lead to the diagnosis of iPPSD with or without the presence of minor criteria. However, brachydactyly type E (BDE) should be associated with at least one major or two minor criteria to suggest iPPSD, as it is a common feature of several other diseases and syndromes (Table 3).

The known molecular causes of PTH/PTHrP signalling disorders are:

- inactivating mutations of *PTH1R*;
- inactivating heterozygous mutations in the coding sequence of *GNAS*-Gsa;
- methylation changes of the DMRs of *GNAS* caused by
 - deletions or duplications at ICRs (*STX16*; *NESP*; *GNAS*-AS1);
 - paternal UPD of chromosome 20q;
 - unknown mechanism(s);
- heterozygous mutations of *PRKARIA*;
- heterozygous mutations of *PDE4D*;
- heterozygous mutations of *PDE3A*.

In contrast to the former diagnostic classification based solely on the phenotype, once iPPSD has been identified (using criteria described Table 3), we propose to further subtype iPPSD based on the underlying molecular (epi) genetic defect. Therefore, the term iPPSD will refer to the pathophysiology of the PTH/PTHrP signalling abnormalities, while the number will refer to the underlying molecular mechanism (responsible for the pathology). We have numbered iPPSD subtypes starting with *PTH1R* mutations.

The novel classification of iPPSD

The European PHP network proposes the following classification (Fig. 4):

- iPPSD: clinical/biochemical diagnosis based on the major/minor criteria as defined above, in the absence of genetic investigation;
- iPPSD1: loss-of-function mutation in *PTH1R*;
- iPPSD2: loss-of-function mutation in Gsa;

Figure 4

Schematic representation of the new classification proposed by the European PHP network. According to the suggested new classification Blomstrand and Eiken chondrodysplasia, PHP type 1 and 2, PPHP, AHO, POH and acrodysostosis clinically/biochemically diagnosed without genetic investigation are named iPPSD; Blomstrand and Eiken chondrodysplasia due to PTHR1-inactivating mutations are named iPPSD1; PHP1A, PHP1C, PPHP and POH clinically diagnosed and characterised by Gsa-inactivating mutations are termed iPPSD2; PHP1B clinically diagnosed and due to methylation changes at the *GNAS* DMRs is classified as iPPSD3; in the presence of acrodysostosis type 1 or *PRKAR1A* mutations, the disease is classified as iPPSD4. Acrodysostosis type 2 or *PDE4D* mutations are termed iPPSD5; *PDE3A* mutations are categorised as iPPSD6; patients lacking genetic or epigenetic defects at the known genes fall under the category of iPPSDx; any newly discovered genetic/molecular defects will be labelled as iPPSDn+1.

- iPPSD3: methylation change(s) at one or more *GNAS* DMRs, associated with or without a genetic (deletion) or cytogenetic (UPD) defect;
- iPPSD4: *PRKAR1A* mutation;
- iPPSD5: *PDE4D* mutation;
- iPPSD6: *PDE3A* mutation;
- iPPSDx: lack of genetic/epigenetic defect identified following molecular investigation of known genes described above;
- iPPSDn+1: the identification of a novel gene/molecular defect will lead to a disease named iPPSD7, then 8 and so on.

iPPSD3 encompasses all disorders associated with changes in the methylation patterns of the DMRs of *GNAS*, including UPD(20)pat and deletion within *STX16*, *NESP* etc. Of most significance is the common mechanism shared by these patients, i.e. the LOM at the *GNAS* A/B:TSS-DMR. Grouping them under iPPSD3 highlights this common mechanism. Secondly, we anticipated the difficulties in integrating the multiplicity of the epigenetic mechanisms within the classification system as this adds no further diagnostic value. However, the further specification of the epigenetic defect can remain part of a private exchange between the molecular laboratory, the patient and his/her physician.

We recommend the use of Arabic numerals to avoid the confusion with letters (II with the number 11 for example).

The advantages of this new nomenclature are: (i) it stratifies the disorders into clusters caused by the same mechanism; (ii) it is flexible and open to accommodate new defects to be discovered in the future and (iii) it simplifies the concept of the overlapping disorders under a single umbrella.

This classification, however, bears some limitations. We deliberately did not include the parental origin of the genetic/epigenetic defect, although some iPPSD are imprinting disorders – namely iPPSD2 and iPPSD3 – and their phenotypic expression depends on their parental inheritance. The main reason behind this is the association of PTH resistance and POH with both maternal and paternal inactivating *GNAS* mutations. Therefore, the mechanism of the two allelic *GNAS* mutations can be considered alike. However, in daily practice, the parental origin of the *GNAS* defect should be considered, particularly for genetic counselling. In fact, AHO and multiple hormone resistance including PTH resistance are largely associated with maternal *GNAS* coding defects, whereas isolated AHO and/or POH are more often associated with paternal *GNAS* coding defects.

Another limitation of this classification is the inability to subclassify individuals with a pure clinical suspicion of iPPSD and lack of complete (epi)genetic testing. While such patients cannot be classified as iPPSDx or with a specific number, we recommend that they are classified as iPPSD.

The inclusion of the disorders involving the two main ligands of the PTH1R, i.e. hypoparathyroidism (109) and brachydactyly type E with short stature (mutations in *PTH1R* the gene encoding PTHrP (110, 111)) to the classification may be argued. However, we decided to exclude them due to several other issues such as (i) their different biochemical pattern including low levels of PTH responsible for hypoparathyroidism; (ii) the dramatic difference in the therapy of hypoparathyroidism and defects in PTH1R signalling respectively and (iii) the difference in research goals in the two disease groups.

Perspectives

We believe that the use of the new nomenclature will facilitate a more straightforward approach to the diagnosis of iPPSD, increase awareness of the red-flag signs of PTH resistance, ectopic ossifications and brachydactyly type E. It would allow for the classification of patients into local catalogues used by the different healthcare organisations in a more homogenous way, and enable future observational and research studies in the field.

We strongly believe that too many denominations for similar diseases and patients with phenocopies (PHP, PPHP, POH, ACRDYS, TRPS, BDE, AHO) have diluted and dispersed research advance, adding undue complexity to the causative mechanism and proved challenging for the experts in building a global research network in the field.

Regular use of the classification in daily practice or for scientific purposes will allow appropriate amendments in the best interest of the patients.

While producing this novel nomenclature and classification, we have identified the need to (i) disseminate this alternative classification to be positively enriched by the clinical and scientific community; (ii) validate the major/minor criteria in a series of patients affected by different iPPSDs and (iii) develop international guidelines for the diagnosis and treatment of the iPPSDs in the near future.

Declaration of interest

All the authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work and/or members of the network Euro-Pseudohypoparathyroidism network (EuroPHP) was supported by grants obtained from the European Society for Paediatric Endocrinology (ESPE) Research Unit, the University of Luebeck (H02-2011), the Italian Ministry of Health (GR-2009-1608394), Ricerca Corrente Funds (to Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico), the Instituto de Salud Carlos III and European fund for Regional Development, FEDER, (PI13/00467 to GPdN), the Basque Department of Health (GV2014111017 to GPdN), the University of Basque Country (Ref: 48198) (to A P), the I3SNS Programme of the Spanish Ministry of Health (CP03/0064; SIVI 1395/09 to GPdN), the APHP Bicêtre Paris-Sud Reference Center for Rare Disorders of the Calcium and Phosphate Metabolism (to A L, C S, V G, P H and A U) as well as the Plateforme d'Expertise Maladies Rares Paris-Sud. All members of the EuroPHP network are members of the EUCID.net (COST action BM1208 on imprinting disorders; www.imprinting-disorders.eu), which was supported by SANDOZ FRANCE for training actions.

Author contribution statement

S Thiele, G Mantovani, G Perez de Nanclares and A Linglart contributed equally to the manuscript; all the other members of the EuroPHP are listed in alphabetical order.

References

- Mantovani G. Clinical review: pseudohypoparathyroidism: diagnosis and treatment. *Journal of Clinical Endocrinology and Metabolism* 2011 **96** 3020–3030. (doi:10.1210/jc.2011-1048)
- Weinstein LS, Yu S, Warner DR & Liu J. Endocrine manifestations of stimulatory G protein alpha-subunit mutations and the role of genomic imprinting. *Endocrine Reviews* 2001 **22** 675–705. (doi:10.1210/edrv.22.5.0439)
- Turan S & Bastepe M. The GNAS complex locus and human diseases associated with loss-of-function mutations or epimutations within this imprinted gene. *Hormone Research in Paediatrics* 2013 **80** 229–241. (doi:10.1159/000355384)
- Albright F, Burnett CH, Smith PH & Parson W. Pseudohypoparathyroidism – an example of “Seabright-Bantam syndrome”. *Endocrinology* 1942 **30** 922–932.
- Chase LR, Melson GL & Aurbach GD. Pseudohypoparathyroidism: defective excretion of 3',5'-AMP in response to parathyroid hormone. *Journal of Clinical Investigation* 1969 **48** 1832–1844. (doi:10.1172/JCI106149)
- Levine MA, Downs RW Jr, Moses AM, Breslau NA, Marx SJ, Lasker RD, Rizzoli RE, Aurbach GD & Spiegel AM. Resistance to multiple hormones in patients with pseudohypoparathyroidism. Association with deficient activity of guanine nucleotide regulatory protein. *American Journal of Medicine* 1983 **74** 545–556. (doi:10.1016/0002-9343(83)91008-2)
- Cheloha RW, Gellman SH, Vilardaga J-P & Gardella TJ. PTH receptor-1 signalling-mechanistic insights and therapeutic prospects. *Nature Reviews Endocrinology* 2015 **11** 712–724. (doi:10.1038/nrendo.2015.139)
- Blomstrand S, Claesson I & Säve-Söderbergh J. A case of lethal congenital dwarfism with accelerated skeletal maturation. *Pediatric Radiology* 1985 **15** 141–143. (doi:10.1007/BF02388725)
- Zhang P, Jobert AS, Couvineau A & Silve C. A homozygous inactivating mutation in the parathyroid hormone/parathyroid hormone-related peptide receptor causing Blomstrand chondrodysplasia. *Journal of Clinical Endocrinology and Metabolism* 1998 **83** 3365–3368. (doi:10.1210/jcem.83.9.5243)
- Duchatelet S, Ostergaard E, Cortes D, Lemainque A & Julier C. Recessive mutations in PTH1R cause contrasting skeletal dysplasias in Eiken and Blomstrand syndromes. *Human Molecular Genetics* 2005 **14** 1–5. (doi:10.1093/hmg/ddi118)

Position Statement	S Thiele and others	iPPSD, a novel classification for PHP	175:6	P14
--------------------	---------------------	---------------------------------------	-------	-----

- 11 Hoogendam J, Farih-Sips H, Wynaendts LC, Löwik CWGM, Wit JM & Karperien M. Novel mutations in the parathyroid hormone (PTH)/PTH-related peptide receptor type 1 causing Blomstrand osteochondrodysplasia types I and II. *Journal of Clinical Endocrinology and Metabolism* 2007 **92** 1088–1095. (doi:10.1210/jc.2006-0300)
- 12 Stone M, Hosking D, Garcia-Himmelstine C, White D, Rosenblum D & Worth H. The renal response to exogenous parathyroid hormone in treated pseudohypoparathyroidism. *Bone* 1993 **14** 727–735. (doi:10.1016/8756-3282(93)90204-N)
- 13 Drezner M, Neelon FA & Lebovitz HE. Pseudohypoparathyroidism type II: a possible defect in the reception of the cyclic AMP signal. *New England Journal of Medicine* 1973 **289** 1056–1060. (doi:10.1056/NEJM197311152892003)
- 14 Rao DS, Parfitt AM, Kleerekoper M, Pumo BS & Frame B. Dissociation between the effects of endogenous parathyroid hormone on adenosine 3',5'-monophosphate generation and phosphate reabsorption in hypocalcemia due to vitamin D depletion: an acquired disorder resembling pseudohypoparathyroidism type II. *Journal of Clinical Endocrinology and Metabolism* 1985 **61** 285–290. (doi:10.1210/jcem-61-2-285)
- 15 Rodriguez HJ, Villarreal H Jr, Klahr S & Slatopolsky E. Pseudohypoparathyroidism type II: restoration of normal renal responsiveness to parathyroid hormone by calcium administration. *Journal of Clinical Endocrinology and Metabolism* 1974 **39** 693–701. (doi:10.1210/jcem-39-4-693)
- 16 Linglart A, Menguy C, Couvineau A, Auzan C, Gunes Y, Cancel M, Motte E, Pinto G, Chanson P, Bougnères P *et al.* Recurrent PRKAR1A mutation in acrodysostosis with hormone resistance. *New England Journal of Medicine* 2011 **364** 2218–2226. (doi:10.1056/NEJMoa1012717)
- 17 Levine MA, Downs RW Jr, Singer M, Marx SJ, Aurbach GD & Spiegel AM. Deficient activity of guanine nucleotide regulatory protein in erythrocytes from patients with pseudohypoparathyroidism. *Biochemical and Biophysical Research Communications* 1980 **94** 1319–1324. (doi:10.1016/0006-291X(80)90563-X)
- 18 Farfel Z, Brickman AS, Kaslow HR, Brothers VM & Bourne HR. Defect of receptor-cyclase coupling protein in pseudohypoparathyroidism. *New England Journal of Medicine* 1980 **303** 237–242. (doi:10.1056/NEJM198007313030501)
- 19 Radeke HH, Auf'mkolk B, Jüppner H, Krohn HP, Keck E & Hesch RD. Multiple pre- and postreceptor defects in pseudohypoparathyroidism (a multicenter study with twenty four patients). *Journal of Clinical Endocrinology and Metabolism* 1986 **62** 393–402. (doi:10.1210/jcem-62-2-393)
- 20 Silve C, Santora A, Breslau N, Moses A & Spiegel A. Selective resistance to parathyroid hormone in cultured skin fibroblasts from patients with pseudohypoparathyroidism type Ib. *Journal of Clinical Endocrinology and Metabolism* 1986 **62** 640–644. (doi:10.1210/jcem-62-4-640)
- 21 Albright F, Forbes AP & Henneman PH. Pseudo-pseudohypoparathyroidism. *Transactions of the Association of American Physicians* 1952 **65** 337–350.
- 22 Eyre WG & Reed WB. Albright's hereditary osteodystrophy with cutaneous bone formation. *Archives of Dermatology* 1971 **104** 634–642. (doi:10.1001/archderm.1971.04000240058008)
- 23 Long DN, McGuire S, Levine MA, Weinstein LS & Germain-Lee EL. Body mass index differences in pseudohypoparathyroidism type Ia versus pseudopseudohypoparathyroidism may implicate paternal imprinting of Galpha(s) in the development of human obesity. *Journal of Clinical Endocrinology and Metabolism* 2007 **92** 1073–1079. (doi:10.1210/jc.2006-1497)
- 24 Farfel Z & Friedman E. Mental deficiency in pseudohypoparathyroidism type I is associated with Ns-protein deficiency. *Annals of Internal Medicine* 1986 **105** 197–199. (doi:10.7326/0003-4819-105-2-197)
- 25 Mouallem M, Shaharabany M, Weintrob N, Shalitin S, Nagelberg N, Shapira H, Zadik Z & Farfel Z. Cognitive impairment is prevalent in pseudohypoparathyroidism type Ia, but not in pseudopseudohypoparathyroidism: possible cerebral imprinting of Gsalpha. *Clinical Endocrinology* 2008 **68** 233–239. (doi:10.1111/j.1365-2265.2007.03025.x)
- 26 Balavoine AS, Ladsous M, Velayoudom FL, Vlaeminck V, Cardot-Bauters C, d'Herbomez M & Wemeau JL. Hypothyroidism in patients with pseudohypoparathyroidism type Ia: clinical evidence of resistance to TSH and TRH. *European Journal of Endocrinology* 2008 **159** 431–437. (doi:10.1530/EJE-08-0111)
- 27 Vlaeminck-Guillem V, D'Herbomez M, Pigny P, Fayard A, Bauters C, Decoulx M & Wemeau JL. Pseudohypoparathyroidism Ia and hypercalcaemia. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 3091–3096. (doi:10.1210/jcem.86.7.7690)
- 28 Linglart A, Maupetit-Méhouas S & Silve C. GNAS-related loss-of-function disorders and the role of imprinting. *Hormone Research in Paediatrics* 2013 **79** 119–129. (doi:10.1159/000348516)
- 29 Brix B, Werner R, Staedt P, Struve D, Hiort O & Thiele S. Different pattern of epigenetic changes of the GNAS gene locus in patients with pseudohypoparathyroidism type Ic confirm the heterogeneity of underlying pathomechanisms in this subgroup of pseudohypoparathyroidism and the demand for a new classification of GNAS-related disorders. *Journal of Clinical Endocrinology and Metabolism* 2014 **99** E1564–E1570. (doi:10.1210/jc.2013-4477)
- 30 Thiele S, de Sanctis L, Werner R, Grotzinger J, Aydin C, Juppner H, Bastepe M & Hiort O. Functional characterization of GNAS mutations found in patients with pseudohypoparathyroidism type Ic defines a new subgroup of pseudohypoparathyroidism affecting selectively Gsalpha-receptor interaction. *Human Mutation* 2011 **32** 653–660. (doi:10.1002/humu.21489)
- 31 Hirai T, Chagin AS, Kobayashi T, Mackem S & Kronenberg HM. Parathyroid hormone/parathyroid hormone-related protein receptor signaling is required for maintenance of the growth plate in postnatal life. *PNAS* 2011 **108** 191–196. (doi:10.1073/pnas.1005011108)
- 32 Patten JL, Johns DR, Valle D, Eil C, Gruppiso PA, Steele G, Smallwood PM & Levine MA. Mutation in the gene encoding the stimulatory G protein of adenylate cyclase in Albright's hereditary osteodystrophy. *New England Journal of Medicine* 1990 **322** 1412–1419. (doi:10.1056/NEJM199005173222002)
- 33 Elli FM, deSanctis L, Ceoloni B, Barbieri AM, Bordogna P, Beck-Peccoz P, Spada A & Mantovani G. Pseudohypoparathyroidism type Ia and pseudo-pseudohypoparathyroidism: the growing spectrum of GNAS inactivating mutations. *Human Mutation* 2013 **34** 411–416. (doi:10.1002/humu.22265)
- 34 Thiele S, Werner R, Grötzinger J, Brix B, Staedt P, Struve D, Reiz B, Farida J & Hiort O. A positive genotype-phenotype correlation in a large cohort of patients with pseudohypoparathyroidism type Ia and pseudo-pseudohypoparathyroidism and 33 newly identified mutations in the GNAS gene. *Molecular Genetics & Genomic Medicine* 2015 **3** 111–120. (doi:10.1002/mgg3.2015.3.issue-2)
- 35 Lemos MC & Thakker RV. GNAS mutations in pseudohypoparathyroidism type Ia and related disorders. *Human Mutation* 2015 **36** 11–19. (doi:10.1002/humu.2015.36.issue-1)
- 36 Weinstein LS, Gejman PV, Friedman E, Kadowaki T, Collins RM, Gershon ES & Spiegel AM. Mutations of the Gs alpha-subunit gene in Albright hereditary osteodystrophy detected by denaturing gradient gel electrophoresis. *PNAS* 1990 **87** 8287–8290.
- 37 Mantovani G, Bondioni S, Linglart A, Maghnie M, Cisternino M, Corbetta S, Lania AG, Beck-Peccoz P & Spada A. Genetic analysis and evaluation of resistance to thyrotropin and growth hormone-releasing hormone in pseudohypoparathyroidism type Ib. *Journal of Clinical Endocrinology and Metabolism* 2007 **92** 3738–3742. (doi:10.1210/jc.2007-0869)
- 38 Ham H-J, Baek K-H, Lee J-Y, Kim SY, Mo EY, Kim ES, Han JH & Moon SD. Analysis of aberrantly spliced transcripts of a novel de novo GNAS mutant in a male with albright hereditary

- osteodystrophy and PHP1A. *Hormone and Metabolic Research* 2015 **47** 585–590. (doi:10.1055/s-0034-1395678)
- 39 Lin MH, Numbenjapon N, Germain-Lee EL & Pitukcheewanont P. Progressive osseous heteroplasia, as an isolated entity or overlapping with Albright hereditary osteodystrophy. *Journal of Pediatric Endocrinology and Metabolism* 2015 **28** 911–918. (doi:10.1515/jpem-2014-0435)
 - 40 Geneviève D, Sanlaville D, Faivre L, Kottler M-L, Jambou M, Gosset P, Boustani-Samara D, Pinto G, Ozilou C, Abeguilé G *et al.* Paternal deletion of the GNAS imprinted locus (including Gnasxl) in two girls presenting with severe pre- and post-natal growth retardation and intractable feeding difficulties. *European Journal of Human Genetics* 2005 **13** 1033–1039. (doi:10.1038/sj.ejhg.5201448)
 - 41 Fernandez-Rebollo E, García-Cuartero B, Garin I, Largo C, Martínez F, García-Lacalle C, Castaño L, Bastepe M & Pérez de Nanclares G. Intragenic GNAS deletion involving exon A/B in pseudohypoparathyroidism type 1A resulting in an apparent loss of exon A/B methylation: potential for misdiagnosis of pseudohypoparathyroidism type 1B. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 765–771. (doi:10.1210/jc.2009-1581)
 - 42 Fernandez-Rebollo E, Barrio R, Pérez-Nanclares G, Carcavilla A, Garin I, Castaño L & de Nanclares GP. New mutation type in pseudohypoparathyroidism type 1a. *Clinical Endocrinology* 2008 **69** 705–712. (doi:10.1111/cen.2008.69.issue-5)
 - 43 Mitsui T, Nagasaki K, Takagi M, Narumi S, Ishii T & Hasegawa T. A family of pseudohypoparathyroidism type 1a with an 850-kb submicroscopic deletion encompassing the whole GNAS locus. *American Journal of Medical Genetics Part A* 2012 **158A** 261–264. (doi:10.1002/ajmg.a.34393)
 - 44 Garin I, Elli FM, Linglart A, Silve C, de Sanctis L, Bordogna P, Pereda A, Clarke JT, Kannengiesser C, Coutant R *et al.* Novel microdeletions affecting the GNAS locus in pseudohypoparathyroidism: characterization of the underlying mechanisms. *Journal of Clinical Endocrinology and Metabolism* 2015 **100** E681–E687. (doi:10.1210/jc.2014-3098)
 - 45 Adegbite NS, Xu M, Kaplan FS, Shore EM & Pignolo RJ. Diagnostic and mutational spectrum of progressive osseous heteroplasia (POH) and other forms of GNAS-based heterotopic ossification. *American Journal of Medical Genetics Part A* 2008 **146A** 1788–1796. (doi:10.1002/ajmg.a.32346)
 - 46 Shore EM, Ahn J, Jan de Beur S, Li M, Xu M, Gardner RJM, Zasloff MA, Whyte MP, Levine MA & Kaplan FS. Paternally inherited inactivating mutations of the GNAS1 gene in progressive osseous heteroplasia. *New England Journal of Medicine* 2002 **346** 99–106. (doi:10.1056/NEJMoa011262)
 - 47 Hayward B & Bonthron D. An imprinted antisense transcript at the human GNAS1 locus. *Human Molecular Genetics* 2000 **9** 835–841. (doi:10.1093/hmg/9.5.835)
 - 48 Yu S, Yu D, Lee E, Eckhaus M, Lee R, Corria Z, Accili D, Westphal H & Weinstein LS. Variable and tissue-specific hormone resistance in heterotrimeric Gs protein alpha-subunit (Gsalph) knockout mice is due to tissue-specific imprinting of the gsalph gene. *PNAS* 1998 **95** 8715–8720.
 - 49 Mantovani G, Ballare E, Giammona E, Beck-Peccoz P & Spada A. The gsalph gene: predominant maternal origin of transcription in human thyroid gland and gonads. *Journal of Clinical Endocrinology and Metabolism* 2002 **87** 4736–4740. (doi:10.1210/jc.2002-020183)
 - 50 Liu J, Litman D, Rosenberg MJ, Yu S, Biesecker LG & Weinstein LS. A GNAS1 imprinting defect in pseudohypoparathyroidism type 1B. *Journal of Clinical Investigation* 2000 **106** 1167–1174. (doi:10.1172/JCI10431)
 - 51 Bastepe M, Fröhlich LF, Hendy GN, Indridason OS, Josse RG, Koshiyama H, Körkö J, Nakamoto JM, Rosenbloom AL, Slyper AH *et al.* Autosomal dominant pseudohypoparathyroidism type 1b is associated with a heterozygous microdeletion that likely disrupts a putative imprinting control element of GNAS. *Journal of Clinical Investigation* 2003 **112** 1255–1263. (doi:10.1172/JCI19159)
 - 52 Linglart A, Gensure RC, Olney RC, Jüppner H & Bastepe M. A novel STX16 deletion in autosomal dominant pseudohypoparathyroidism type 1b redefines the boundaries of a cis-acting imprinting control element of GNAS. *American Journal of Human Genetics* 2005 **76** 804–814. (doi:10.1086/429932)
 - 53 Bastepe M, Fröhlich LF, Linglart A, Abu-Zahra HS, Tojo K, Ward LM & Jüppner H. Deletion of the NESP55 differentially methylated region causes loss of maternal GNAS imprints and pseudohypoparathyroidism type 1b. *Nature Genetics* 2005 **37** 25–27. (doi:10.1038/ng1560)
 - 54 Richard N, Abeguilé G, Coudray N, Mittre H, Gruchy N, Andrieux J, Cathebras P & Kottler ML. A new deletion ablating NESP55 causes loss of maternal imprint of A/B GNAS and autosomal dominant pseudohypoparathyroidism type 1b. *Journal of Clinical Endocrinology and Metabolism* 2012 **97** E863–E867. (doi:10.1210/jc.2011-2804)
 - 55 Elli FM, de Sanctis L, Peverelli E, Bordogna P, Pivetta B, Miolo G, Beck-Peccoz P, Spada A & Mantovani G. Autosomal dominant pseudohypoparathyroidism type 1b: a novel inherited deletion ablating STX16 causes loss of imprinting at the A/B DMR. *Journal of Clinical Endocrinology and Metabolism* 2014 **99** E724–E728. (doi:10.1210/jc.2013-3704)
 - 56 Chillambhi S, Turan S, Hwang D-Y, Chen H-C, Jüppner H & Bastepe M. Deletion of the noncoding GNAS antisense transcript causes pseudohypoparathyroidism type 1b and biparental defects of GNAS methylation in cis. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 3993–4002. (doi:10.1210/jc.2009-2205)
 - 57 Rezwan FI, Poole RL, Prescott T, Walker JM, Karen Temple I & Mackay DJG. Very small deletions within the NESP55 gene in pseudohypoparathyroidism type 1b. *European Journal of Human Genetics* 2015 **23** 494–499. (doi:10.1038/ejhg.2014.133)
 - 58 Takatani R, Molinaro A, Grigelioniene G, Tafaj O, Watanabe T, Reyes M, Sharma A, Singhal V, Raymond FL, Linglart A *et al.* Analysis of multiple families with single individuals affected by pseudohypoparathyroidism type 1b (PHP1B) reveals only one novel maternally inherited GNAS deletion. *Journal of Bone and Mineral Research* 2016 **31** 796–805. (doi:10.1002/jbmr.2731)
 - 59 Bastepe M, Lane AH & Jüppner H. Paternal uniparental isodisomy of chromosome 20q—and the resulting changes in GNAS1 methylation—as a plausible cause of pseudohypoparathyroidism. *American Journal of Human Genetics* 2001 **68** 1283–1289.
 - 60 Fernández-Rebollo E, Lecumberri B, Garin I, Arroyo J, Bernal-Chico A, Goñi F, Orduña R, Spanish PHP Group, Castaño L & de Nanclares GP. New mechanisms involved in paternal 20q disomy associated with pseudohypoparathyroidism. *European Journal of Endocrinology* 2010 **163** 953–962. (doi:10.1530/EJE-10-0435)
 - 61 Dixit A, Chandler KE, Lever M, Poole RL, Bullman H, Mughal MZ, Steggall M & Suri M. Pseudohypoparathyroidism type 1b due to paternal uniparental disomy of chromosome 20q. *Journal of Clinical Endocrinology and Metabolism* 2013 **98** E103–E108. (doi:10.1210/jc.2012-2639)
 - 62 Takatani R, Minagawa M, Molinaro A, Reyes M, Kinoshita K, Takatani T, Kazukawa I, Nagatsuma M, Kashimada K, Sato K *et al.* Similar frequency of paternal uniparental disomy involving chromosome 20q (patUPD20q) in Japanese and Caucasian patients affected by sporadic pseudohypoparathyroidism type 1b (sporPHP1B). *Bone* 2015 **79** 15–20. (doi:10.1016/j.bone.2015.05.011)
 - 63 Maupetit-Méhouas S, Azzi S, Steunou V, Sakakini N, Silve C, Reynes C, Perez de Nanclares G, Keren B, Chantot S, Barlier A *et al.* Simultaneous hyper- and hypomethylation at imprinted loci in a subset of patients with GNAS epimutations underlies a complex and different mechanism of multilocus methylation defect in pseudohypoparathyroidism type 1b. *Human Mutation* 2013 **34** 1172–1180. (doi:10.1002/humu.22352)
 - 64 Van Dijk FS, Pals G, Van Rijn RR, Nikkels PGJ & Cobben JM. Classification of Osteogenesis Imperfecta revisited. *European Journal of Medical Genetics* 2010 **53** 1–5. (doi:10.1016/j.ejmg.2009.10.007)

Position Statement	S Thiele and others	ippSD, a novel classification for PHP	175:6	P16
--------------------	---------------------	---------------------------------------	-------	-----

- 65 June RR & Aggarwal R. The use and abuse of diagnostic/classification criteria. *Best Practice & Research. Clinical Rheumatology* 2014 **28** 921–934. (doi:10.1016/j.berh.2015.04.004)
- 66 Bonafe L, Cormier-Daire V, Hall C, Lachman R, Mortier G, Mundlos S, Nishimura G, Sangiorgi L, Savarirayan R, Sillence D *et al.* Nosology and classification of genetic skeletal disorders: 2015 revision. *American Journal of Medical Genetics Part A* 2015 **167A** 2869–2892. (doi:10.1002/ajmg.a.37365)
- 67 Eckardt K-U, Alper SL, Antignac C, Bleyer AJ, Chauveau D, Dahan K, Deltas C, Hosking A, Kmoch S, Rampoldi L *et al.* Autosomal dominant tubulointerstitial kidney disease: diagnosis, classification, and management-A KDIGO consensus report. *Kidney International* 2015 **88** 676–683. (doi:10.1038/ki.2015.28)
- 68 Marcucci G, Cianferotti L, Beck-Peccoz P, Capezzone M, Cetani F, Colao A, Davì MV, degli Uberti E, Del Prato S, Elisei R *et al.* Rare diseases in clinical endocrinology: a taxonomic classification system. *Journal of Endocrinological Investigation* 2015 **38** 193–259. (doi:10.1007/s40618-014-0202-6)
- 69 Picard C, Al-Herz W, Bousfiha A, Casanova J-L, Chatila T, Conley ME, Cunningham-Rundles C, Etzioni A, Holland SM, Klein C *et al.* Primary immunodeficiency diseases: an update on the classification from the International Union of Immunological Societies Expert Committee for Primary Immunodeficiency 2015. *Journal of Clinical Immunology* 2015 **35** 696–726. (doi:10.1007/s10875-015-0201-1)
- 70 Alberti KG & Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabetic Medicine: A Journal of the British Diabetic Association* 1998 **15** 539–553. (doi:10.1002/(SICI)1096-9136(199807)15:7<539::AID-DIA668>3.0.CO;2-S)
- 71 de Nancrales GP, Fernandez-Rebollo E, Santin I, Garcia-Cuartero B, Gaztambide S, Menendez E, Morales MJ, Pombo M, Bilbao JR, Barros F *et al.* Epigenetic defects of GNAS in patients with pseudohypoparathyroidism and mild features of Albright's hereditary osteodystrophy. *Journal of Clinical Endocrinology and Metabolism* 2007 **92** 2370–2373. (doi:10.1210/jc.2006-2287)
- 72 Unluturk U, Harmanci A, Babaoglu M, Yasar U, Varli K, Bastepe M & Bayraktar M. Molecular diagnosis and clinical characterization of pseudohypoparathyroidism type-Ib in a patient with mild Albright's hereditary osteodystrophy-like features, epileptic seizures, and defective renal handling of uric acid. *American Journal of the Medical Sciences* 2008 **336** 84–90. (doi:10.1097/MAJ.0b013e31815b218f)
- 73 Mariot V, Maupetit-Méhous S, Sinding C, Kottler M-L & Linglart A. A maternal epimutation of GNAS leads to Albright osteodystrophy and parathyroid hormone resistance. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** 661–665. (doi:10.1210/jc.2007-0927)
- 74 Mantovani G, de Sanctis L, Barbieri AM, Elli FM, Bollati V, Vaira V, Labarile P, Bondioni S, Peverelli E, Lania AG *et al.* Pseudohypoparathyroidism and GNAS epigenetic defects: clinical evaluation of albright hereditary osteodystrophy and molecular analysis in 40 patients. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 651–658. (doi:10.1210/jc.2009-0176)
- 75 Izzi B, Decallonne B, Devriendt K, Bouillon R, Vanderschueren D, Levchenko E, de Zegher F, Van den Bruel A, Lambrechts D, Van Geet C *et al.* A new approach to imprinting mutation detection in GNAS by Sequenom EpiTYPER system. *International Journal of Clinical Chemistry* 2010 **411** 2033–2039. (doi:10.1016/j.cca.2010.08.034)
- 76 Zazo C, Thiele S, Martín C, Fernandez-Rebollo E, Martinez-Indart L, Werner R, Garin I, Spanish PHP Group, Hiort O & Perez de Nancrales G. Gsα activity is reduced in erythrocyte membranes of patients with pseudohypoparathyroidism due to epigenetic alterations at the GNAS locus. *Journal of Bone and Mineral Research* 2011 **26** 1864–1870. (doi:10.1002/jbmr.369)
- 77 Turan S, Thiele S, Tafaj O, Brix B, Atay Z, Abali S, Haliloglu B, Bereket A & Bastepe M. Evidence of hormone resistance in a pseudohypoparathyroidism patient with a novel paternal mutation in GNAS. *Bone* 2015 **71** 53–57. (doi:10.1016/j.bone.2014.10.006)
- 78 Linglart A, Carel JC, Garabédian M, Lé T, Mallet E & Kottler ML. GNAS1 lesions in pseudohypoparathyroidism Ia and Ic: genotype phenotype relationship and evidence of the maternal transmission of the hormonal resistance. *Journal of Clinical Endocrinology and Metabolism* 2002 **87** 189–197. (doi:10.1210/jcem.87.1.8133)
- 79 Lebrun M, Richard N, Abeguilé G, David A, Coëslie Dieux A, Journel H, Lacombe D, Pinto G, Odent S, Salles JP *et al.* Progressive osseous heteroplasia: a model for the imprinting effects of GNAS inactivating mutations in humans. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 3028–3038. (doi:10.1210/jc.2009-1451)
- 80 Kaplan FS & Shore EM. Progressive osseous heteroplasia. *Journal of Bone and Mineral Research* 2000 **15** 2084–2094. (doi:10.1359/jbmr.2000.15.11.2084)
- 81 Eddy MC, De Beur SM, Yandow SM, McAlister WH, Shore EM, Kaplan FS, Whyte MP & Levine MA. Deficiency of the alpha-subunit of the stimulatory G protein and severe extraskeletal ossification. *Journal of Bone and Mineral Research* 2000 **15** 2074–2083. (doi:10.1359/jbmr.2000.15.11.2074)
- 82 Michot C, Le Goff C, Goldenberg A, Abhyankar A, Klein C, Kinning E, Guerrot AM, Flahaut P, Duncombe A, Baujat G *et al.* Exome sequencing identifies PDE4D mutations as another cause of acrodysostosis. *American Journal of Human Genetics* 2012 **90** 740–745. (doi:10.1016/j.ajhg.2012.03.003)
- 83 Lee H, Graham JM Jr, Rimoin DL, Lachman RS, Krejci P, Tompson SW, Nelson SF, Krakow D & Cohn DH. Exome sequencing identifies PDE4D mutations in acrodysostosis. *American Journal of Human Genetics* 2012 **90** 746–751. (doi:10.1016/j.ajhg.2012.03.004)
- 84 Maroteaux P & Malamut G. [Acrodysostosis]. *Presse Médicale* 1968 **76** 2189–2192.
- 85 Robinow M, Pfeiffer RA, Gorlin RJ, McKusick VA, Renuart AW, Johnson GF & Summitt RL. Acrodysostosis. A syndrome of peripheral dysostosis, nasal hypoplasia, and mental retardation. *American Journal of Diseases of Children* 1971 **121** 195–203. (doi:10.1001/archpedi.1971.02100140061002)
- 86 Reiter S. Acrodysostosis. A case of peripheral dysostosis, nasal hypoplasia, mental retardation and impaired hearing. *Pediatric Radiology* 1978 **7** 53–55. (doi:10.1007/BF00975340)
- 87 Davies SJ & Hughes HE. Familial acrodysostosis: can it be distinguished from Albright's hereditary osteodystrophy? *Clinical Dysmorphology* 1992 **1** 207–215.
- 88 Silve C, Le-Stunff C, Motté E, Gunes Y, Linglart A & Clauser E. Acrodysostosis syndromes. *BoneKey Reports* 2012 **1** 225. (doi:10.1038/bonekey.2012.225)
- 89 Ablow RC, Hsia YE & Brandt IK. Acrodysostosis coinciding with pseudohypoparathyroidism and pseudo-pseudohypoparathyroidism. *American Journal of Roentgenology* 1977 **128** 95–99. (doi:10.2214/ajr.128.1.95)
- 90 Linglart A, Fryssira H, Hiort O, Holterhus P-M, Perez de Nancrales G, Argente J, Heinrichs C, Kuechler A, Mantovani G, Leheup B *et al.* PRKAR1A and PDE4D mutations cause acrodysostosis but two distinct syndromes with or without GPCR-signaling hormone resistance. *Journal of Clinical Endocrinology and Metabolism* 2012 **97** E2328–E2338. (doi:10.1210/jc.2012-2326)
- 91 Nagasaki K, Iida T, Sato H, Ogawa Y, Kikuchi T, Saitoh A, Ogata T & Fukami M. PRKAR1A mutation affecting cAMP-mediated G protein-coupled receptor signaling in a patient with acrodysostosis and hormone resistance. *Journal of Clinical Endocrinology and Metabolism* 2012 **97** E1808–E1813. (doi:10.1210/jc.2012-1369)
- 92 Maass PG, Aydin A, Luft FC, Schächterle C, Weise A, Stricker S, Lindschau C, Vaegler M, Qadri F, Toka HR *et al.* PDE3A mutations cause autosomal dominant hypertension with brachydactyly. *Nature Genetics* 2015 **47** 647–653. (doi:10.1038/ng.3302)
- 93 Mantovani G, Elli FM & Spada A. GNAS epigenetic defects and pseudohypoparathyroidism: time for a new classification? *Hormone and Metabolic Research* 2012 **44** 716–723. (doi:10.1055/s-000000025)

- 94 Bastepe M. Genetics and epigenetics of parathyroid hormone resistance. *Endocrine Development* 2013 **24** 11–24. (doi:10.1159/000342494)
- 95 Pereda A, Garin I, Garcia-Barcina M, Gener B, Beristain E, Ibañez AM & Perez de Nanclares G. Brachydactyly E: isolated or as a feature of a syndrome. *Orphanet Journal of Rare Diseases* 2013 **8** 141. (doi:10.1186/1750-1172-8-141)
- 96 Romanet P, Osei L, Netchine I, Pertuit M, Enjalbert A, Reynaud R & Barlier A. Case report of GNAS epigenetic defect revealed by a congenital hypothyroidism. *Pediatrics* 2015 **135** e1079–e1083. (doi:10.1542/peds.2014-2806)
- 97 Germain-Lee EL, Groman J, Crane JL, Jan de Beur SM & Levine MA. Growth hormone deficiency in pseudohypoparathyroidism type 1a: another manifestation of multihormone resistance. *Journal of Clinical Endocrinology and Metabolism* 2003 **88** 4059–4069. (doi:10.1210/jc.2003-030028)
- 98 Mantovani G, Maghnie M, Weber G, De Menis E, Brunelli V, Cappa M, Loli P, Beck-Peccoz P & Spada A. Growth hormone-releasing hormone resistance in pseudohypoparathyroidism type ia: new evidence for imprinting of the Gs alpha gene. *Journal of Clinical Endocrinology and Metabolism* 2003 **88** 4070–4074. (doi:10.1210/jc.2002-022028)
- 99 de Sanctis L, Bellone J, Salerno M, Faleschini E, Caruso-Nicoletti M, Cicchetti M, Concolino D, Balsamo A, Buzi F, Ghizzoni L *et al.* GH secretion in a cohort of children with pseudohypoparathyroidism type Ia. *Journal of Endocrinological Investigation* 2007 **30** 97–103. (doi:10.1007/BF03347406)
- 100 Namnoum AB, Merriam GR, Moses AM & Levine MA. Reproductive dysfunction in women with Albright's hereditary osteodystrophy. *Journal of Clinical Endocrinology and Metabolism* 1998 **83** 824–829. (doi:10.1210/jcem.83.3.4652)
- 101 Carel JC, Le Stunff C, Condamine L, Mallet E, Chaussain JL, Adnot P, Garabédian M & Bougnères P. Resistance to the lipolytic action of epinephrine: a new feature of protein Gs deficiency. *Journal of Clinical Endocrinology and Metabolism* 1999 **84** 4127–4131. (doi:10.1210/jcem.84.11.6145)
- 102 Kadilli I, Colicchio S, Guglielmo R, Vollono C, Della Marca G & Janiri L. Clinical insights by the presence of bipolar disorder in pseudohypoparathyroidism type 1A. *General Hospital Psychiatry* 2015 **37** 497.e3–497.e5. (doi:10.1016/j.genhosppsych.2015.06.008)
- 103 Richard N, Molin A, Coudray N, Rault-Guillaume P, Jüppner H & Kottler M-L. Paternal GNAS mutations lead to severe intrauterine growth retardation (IUGR) and provide evidence for a role of XLas in fetal development. *Journal of Clinical Endocrinology and Metabolism* 2013 **98** E1549–E1556. (doi:10.1210/jc.2013-1667)
- 104 Bréhin A-C, Colson C, Maupetit-Méhouas S, Grybek V, Richard N, Linglart A, Kottler ML & Jüppner H. Loss of methylation at GNAS exon A/B is associated with increased intrauterine growth. *Journal of Clinical Endocrinology and Metabolism* 2015 **100** E623–E631. (doi:10.1210/jc.2014-4047)
- 105 Germain-Lee EL. Short stature, obesity, and growth hormone deficiency in pseudohypoparathyroidism type 1a. *Pediatric Endocrinology Reviews* 2006 **3** (Supplement 2) 318–327.
- 106 Shoemaker AH, Lomenick JP, Saville BR, Wang W, Buchowski MS & Cone RD. Energy expenditure in obese children with pseudohypoparathyroidism type 1a. *International Journal of Obesity* 2013 **37** 1147–1153. (doi:10.1038/ijo.2012.200)
- 107 Roizen JD, Danzig J, Groleau V, McCormack S, Casella A, Harrington J, Sochett E, Tershakovec A, Zemel BS, Stallings VA *et al.* Resting energy expenditure is decreased in pseudohypoparathyroidism type 1A. *Journal of Clinical Endocrinology and Metabolism* 2015 **101** 880–888. (doi:10.1210/jc.2015-3895)
- 108 Lynch DC, Dymont DA, Huang L, Nikkel SM, Lacombe D, Campeau PM, Lee B, Bacino CA, Michaud JL, Bernier FP *et al.* Identification of novel mutations confirms PDE4D as a major gene causing acrodysostosis. *Human Mutation* 2013 **34** 97–102. (doi:10.1002/humu.22222)
- 109 Bilezikian JP, Khan A, Potts JT, Brandi ML, Clarke BL, Shoback D, Jüppner H, D'Amour P, Fox J, Rejnmark L *et al.* Hypoparathyroidism in the adult: epidemiology, diagnosis, pathophysiology, target-organ involvement, treatment, and challenges for future research. *Journal of Bone and Mineral Research* 2011 **26** 2317–2337. (doi:10.1002/jbmr.483)
- 110 Klopocki E, Hennig BP, Dathe K, Koll R, de Ravel T, Baten E, Blom E, Gillerot Y, Weigel JF, Krüger G *et al.* Deletion and point mutations of PTHLH cause brachydactyly type E. *American Journal of Human Genetics* 2010 **86** 434–439. (doi:10.1016/j.ajhg.2010.01.023)
- 111 Thomas-Teinturier C, Pereda A, Garin I, Diez-Lopez I, Linglart A, Silve C & de Nanclares GP. Report of two novel mutations in PTHLH associated with brachydactyly type E and literature review. *American Journal of Medical Genetics Part A* 2016 **170** 734–742. (doi:10.1002/ajmg.a.37490)
- 112 Schwindinger WF, Miric A, Zimmerman D & Levine MA. A novel Gs alpha mutant in a patient with Albright hereditary osteodystrophy uncouples cell surface receptors from adenylyl cyclase. *Journal of Biological Chemistry* 1994 **269** 25387–25391.
- 113 Srivastava T, Krudys J, Mardis NJ, Sebestyen-VanSickle J & Alon US. Cinacalcet as adjunctive therapy in pseudohypoparathyroidism type 1b. *Pediatric Nephrology* 2016 **31** 795–800. (doi:10.1007/s00467-015-3271-7)
- 114 Bakker B, Sonneveld LJH, Woltering MC, Bikker H & Kant SG. A girl with Beckwith-Wiedemann syndrome and pseudohypoparathyroidism type 1B due to multiple imprinting defects. *Journal of Clinical Endocrinology and Metabolism* 2015 **100** 3963–3966. (doi:10.1210/jc.2015-2260)
- 115 Perez-Nanclares G, Velayos T, Vela A, Muñoz-Torres M & Castaño L. Pseudohypoparathyroidism type Ib associated with novel duplications in the GNAS locus. *PLoS ONE* 2015 **10** e0117691. (doi:10.1371/journal.pone.0117691)
- 116 Mehraein Y, Pfob M, Steinlein O, Aichinger E, Eggert M, Bubendorff V, Mannhart A & Müller S. 2q37.3 deletion syndrome: two cases with highly distinctive facial phenotype, discordant association with schizophrenic psychosis, and shared deletion breakpoint region on 2q37.3. *Cytogenetic and Genome Research* 2015 **146** 33–38. (doi:10.1159/000431389)
- 117 Muhn F, Klopocki E, Graul-Neumann L, Uhrig S, Colley A, Castori M, Lankes E, Henn W, Gruber-Sedlmayr U, Seifert W *et al.* Novel mutations of the PRKAR1A gene in patients with acrodysostosis. *Clinical Genetics* 2013 **84** 531–538. (doi:10.1111/cge.2013.84.issue-6)
- 118 Kaname T, Ki C-S, Niikawa N, Baillie GS, Day JP, Yamamura K-I, Ohta T, Nishimura G, Mastuura N, Kim OH *et al.* Heterozygous mutations in cyclic AMP phosphodiesterase-4D (PDE4D) and protein kinase A (PKA) provide new insights into the molecular pathology of acrodysostosis. *Cellular Signalling* 2014 **26** 2446–2459. (doi:10.1016/j.cellsig.2014.07.025)
- 119 Lindstrand A, Grigelioniene G, Nilsson D, Pettersson M, Hofmeister W, Anderlid B-M, Kant SG, Ruivenkamp CA, Gustavsson P, Valta H *et al.* Different mutations in PDE4D associated with developmental disorders with mirror phenotypes. *Journal of Medical Genetics* 2014 **51** 45–54. (doi:10.1136/jmedgenet-2013-101937)
- 120 Mitsui T, Kim O-H, Hall CM, Offiah A, Johnson D, Jin D-K, Toh TH, Soneda S, Keino D, Matsubayashi S *et al.* Acroscaphodysplasia as a phenotypic variation of pseudohypoparathyroidism and acrodysostosis type 2. *American Journal of Medical Genetics Part A* 2014 **164A** 2529–2534. (doi:10.1002/ajmg.a.36669)

Received 8 February 2016

Revised version received 5 July 2016

Accepted 7 July 2016