Anthropometric factors have significant influence on the outcome of the GHRH-arginine test: establishment of normative data for an automated immunoassay specifically measuring 22 kDa human growth hormone

Timo Deutschbein¹, Martin Bidlingmaier², Jochen Schopohl²,
Christian J Strasburger³ and Stephan Petersenn⁴

¹Department of Internal Medicine I, Division of Endocrinology and Diabetes, University Hospital Würzburg, University of Würzburg, Würzburg, Germany, ²Medizinische Klinik und Poliklinik IV, Klinikum der LMU München, Munich, Germany, ³Department of Medicine for Endocrinology, Diabetes and Nutritional Medicine, Charité Universitätsmedizin, Campus Mitte, Berlin, Germany, and ⁴ENDOC Center for Endocrine Tumors, Hamburg, Germany

Correspondence
should be addressed
to S Petersenn

Email
[stephan.petersenn@
endoc-med.de](mailto:stephan.petersenn@endoc-med.de)

Abstract

Context: Adult growth hormone (GH) deficiency (GHD) is diagnosed by provocative testing of GH secretion.

Objective: To improve the diagnostic accuracy of GH-releasing hormone (GHRH) plus arginine (GARG) testing, we evaluated the influence of age, BMI and sex and established normative data for an automatic immunoassay specifically measuring 22 kDa human GH.

Design/setting: Prospective multicenter study.

Participants: Eighty-seven patients with hypothalamic–pituitary disease and 200 healthy controls. Patients were classified according to the number of pituitary hormone deficiencies (PHD). GHD was assumed when ≥ 2 PHD (in addition to GH) were present ($n=51$); 36 patients with < 2 PHD were considered GH sufficient (GHS). ROC analysis identified cutoffs with $\geq 95\%$ specificity for GHD. Controls were prospectively stratified for sex, age and BMI.

Interventions: All participants received GHRH and L-arginine.

Main outcome measures: GH was measured by immunoassay (iSYS, IDS).

Results: In controls, multiple stepwise regression analysis showed that BMI (21%, $P<0.0001$), sex (20%, $P<0.0001$) and age (5%, $P<0.001$), accounted for 46% of GH peak level variability during GARG. Comparison of peak GH during GARG (GHD vs GHS+controls) revealed an overall cutoff of 3.9 ng/mL (sensitivity 86%, specificity 95%). After adjustment for BMI and sex, optimal cutoffs (male vs female) were 6.5 vs 9.7 ng/mL in lean, 3.5 vs 8.5 ng/mL in overweight and 2.2 vs 4.4 ng/mL in obese subjects respectively.

Conclusion: BMI and sex account for most of the variability of peak GH levels during GARG. Consequently, diagnostic accuracy of the GARG test is significantly improved by use of adjusted cutoffs.

European Journal of
Endocrinology
(2017) **176**, 273–281

Introduction

Human growth hormone (GH) deficiency (GHD) is associated with significant morbidity and mortality (1, 2). For instance, large series have repeatedly shown

that hypopituitarism is generally associated with an increase in overall mortality, cardiovascular events and malignancies compared with the normal population (3, 4).

Apart from that, GH-deficient individuals (especially those without replacement therapy) experience a decrease in quality of life, bone mineral density and lean body mass. Accordingly, careful diagnostic workup of suspected patients is imperatively required.

In patients with proven panhypopituitarism (i.e., at least 3 pituitary hormone deficiencies (PHD)), low levels of circulating insulin-like growth factor 1 (IGF-I), the primary mediator of the effects of GH, are highly suspicious of somatotroph insufficiency (1, 2, 5, 6, 7). However, as illustrated in current guidelines, most patients should undergo provocative testing to establish the diagnosis of GHD (1, 2, 7). Although the insulin tolerance test (ITT) is still regarded as gold standard (1, 2, 7), it is usually not widely applied (6). This is mainly explained by its need for serious hypoglycemia, resulting in undoubted limitations due to potential risks and several contraindications (e.g. ischemic heart disease, seizures).

As a consequence, several other dynamic endocrine tests have been evaluated to explore somatotroph function, for instance, single or combined stimulation with GH secretagogues like arginine, clonidine, GH-releasing hormone (GHRH), GH-releasing peptide 6 (GHRP6), glucagon and levodopa (L-DOPA) (5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16). Some of these tests have severe limitations (e.g. impaired sensitivity, high costs); however, the GHRH plus arginine (GARG) test has repeatedly been described as a good alternative to the ITT (1, 2, 5, 7, 17, 18, 19). Of note, as GH responses during the GARG test decline by increasing body mass index (BMI) and age, adjusted cutoffs have been proposed (13, 14). Unfortunately, however, the weighted influence of these two variables (either alone or in combination) as well as the influence of sex have not been adequately evaluated so far.

Recently, desirable characteristics for GH measurement have been stated by a consensus recommendation (20). These aspects are fulfilled by a modern automated immunoassay specifically measuring 22 kDa human GH (21).

Our study was therefore designed to empirically define reference data for stimulated GH responses in adults, improving the diagnostic accuracy of the GARG test using the formerly mentioned immunoassay. We investigated a well-defined cohort of healthy subjects who were prospectively recruited in such a way that anthropometric factors (e.g. age, BMI, and sex) were reflected on a sufficiently weighted basis. Multiple stepwise regression analysis was performed to identify their potential influence on GH levels. To identify cutoffs with at least 95% specificity for GHD, patients

suffering from hypothalamic–pituitary diseases were included, using the degree of pituitary impairment as gold standard and applying receiver-operating characteristics (ROC) analysis.

Subjects and methods

Subjects

The study protocol was approved by the local ethics committee (approval number: 04-2390), and all study participants provided written informed consent.

Controls

Two hundred healthy subjects were recruited from the general population. None of these subjects had a history of severe and/or chronic illness (especially of endocrine origin). Females on hormonal contraceptives or hormone replacement therapy as well as all subjects who were currently or in the prior months taking drugs known to interfere with the synthesis or metabolism of endocrine parameters were excluded. To evaluate the influence of anthropometric variables, subjects were prospectively stratified for sex (male vs female), age (18–30 years, 31–50 years and >50 years) and BMI (<25 kg/m², 25–30 kg/m² and >30 kg/m²). Study participants were selected in such a way that each group comprised at least 10 subjects (i.e. 10 males aged 18–30 years with a BMI <25 kg/m²; 10 males aged 31–50 years with a BMI <25 kg/m²; 10 males older than 50 years with a BMI <25 kg/m²; and so on). Unfortunately, one control subject was misclassified at screening. Therefore, the group ‘females, 31–50 years, BMI >30 kg/m²’ contains only 9 subjects.

Patients

Eighty-seven patients with hypothalamic–pituitary diseases were enrolled (details are given in Table 1). Subjects who had already been surgically treated had their first regular endocrine workup at least three months after surgery. Two of the 87 patients (2.3%) had been treated by irradiation 1 and 9 years prior to GARG testing respectively. Both of them had insufficiencies of all pituitary axes at the time of GARG testing. Patients with suspected or proven pituitary insufficiency received hormonal replacement therapy (i.e. glucocorticoids, gonadal steroids, thyroxine and desmopressin) with the exception of growth hormone. In detail, 45 patients with corticotrophic insufficiency received hydrocortisone orally

Clinical Study	T Deutschbein and others	BMI and sex: impact on GH after GHRH-arginine	176:3	275

Table 1 Clinical characteristics of patients.

	GHD			GHS		
	Females	Males	All	Females	Males	All
Number (n)	21	30	51	15	21	36
Age (years)	49.6±2.8	49.9±2.4	49.8±1.8	51.0±4.0	44.4±3.1	47.2±2.5
BMI (kg/m ²)	28.2±1.3	28.5±0.9	28.3±0.7	24.4±1.3	27.9±1.6	26.4±1.0
Sellar mass with surgery (n)	14	14	28	9	10	19
Sellar mass without surgery (n)	5	11	16	2	5	7
Hormonal impairment without sellar mass (n)	2	5	7	4	6	10
Pituitary hormone deficiencies (n)	3.0±0.1	3.1±0.1	3.1±0.1	0.6±0.1	0.3±0.1	0.4±0.1

BMI, body mass index; GH, growth hormone; GHD, growth hormone deficiency.

(median dose per day 15 mg, range 10–40 mg), with dose adjustment depending on signs and symptoms of under- or over-replacement. Of note, only 2 patients received doses >25 mg (i.e. 30 and 40 mg respectively). Fifty-four patients with thyrotropic insufficiency were substituted with L-thyroxin, with doses adjusted according to both peripheral hormone levels and clinical symptoms. At the time of GARG testing, fT4 levels were all in the normal range (median fT4 1.37 ng/L, range 0.8–2.1 ng/L). Thirty-four male subjects received testosterone replacement, with individual doses depending on hormone levels, clinical symptoms and age. In these subjects treated with testosterone, a median testosterone of 3.3 ng/mL (range 1.1–10.7 ng/mL) was measured at the time of GARG testing. Of note, one subject had levels below the normal range of our assay. Premenopausal females with gonadotrophic insufficiency were substituted with oral contraceptives or hormone replacement doses, depending on individual preferences.

Methods

Collection of samples

Blood sampling was generally performed between 08:00 h and 10:00 h after an overnight fast. Samples were taken by venipuncture of a forearm vein. If measurement was not performed directly afterward, blood samples were centrifuged and stored at –20°C.

Dynamic testing procedures for the assessment of somatotroph function

During GARG, patients received GHRH (1 µg/kg) by an i.v. bolus, followed by a 30-min i.v. infusion of L-arginine (30 g). Blood samples were taken at 0, 30, 45, 60, 90 and 120 min. In the initial phase of the study, we also evaluated the diagnostic performance of the ITT. In 44 patients,

administration of insulin (0.10–0.15 U/kg) was followed by blood sampling at 0, 15, 30, 45, 60, 90 and 120 min. Both target serum glucose of ≤40 mg/dL and symptoms of neuroglucopenia were required as indicators of sufficient stress response.

Additional hormonal workup

In 44 patients, corticotroph function was investigated by cortisol peak levels during ITT. In the remaining patients, adrenal insufficiency was established either by repeatedly low early morning serum cortisol or by corticotropin-releasing hormone (CRH) testing. Parameters used for characterization of gonadotroph function were testosterone, luteinizing hormone (LH) and follicle-stimulating hormone (FSH) (males); history of regular menses along with estradiol, progesterone, LH and FSH (premenopausal females); LH and FSH only (postmenopausal females). Selected cases also underwent gonadotropin-releasing hormone (GnRH) testing. Thyrotroph function was assessed by thyroid-stimulating hormone (TSH) and peripheral thyroid hormones, with additional thyrotropin-releasing hormone (TRH) testing in some cases.

Interpretation of endocrine parameters

Similar to the approach published by Biller *et al.*, reduced responses to testing of the corticotrophic, gonadotrophic and thyrotrophic axes as well as evidence of diabetes insipidus were separately considered as PHD (5). Therefore, the possible range for grading the severity of PHD was one to four, with two or more insufficiencies (i.e., other than GHD) being classified as multiple PHD (MPHD). As already outlined by Biller *et al.*, prior studies have shown that the latter group has an approximately 90% chance of having severe GHD (22, 23). Patients with less than 2 PHD were classified as GH sufficient (GHS).

Clinical Study	T Deutschbein and others	BMI and sex: impact on GH after GHRH-arginine	176:3	276

Measurement of samples and calculation of estimates

Serum GH was measured by an automated immunoassay (iSYS, Immunodiagnostic Systems (IDS), Boldon, UK) calibrated against the latest International Standard code 98/574 (NIBSC, Hertfordshire, UK). Validation and characterization of the assay had been performed according to recommendations by the clinical laboratory standards institute (CLSI), and results were published in detail elsewhere (21). The epitopes of the monoclonal antibodies used in this assay ensure specific measurement of the 22kDa isoform of GH. Performance of the assay during the study was routinely monitored by internal quality control, longitudinal assessment of serum pools and participation in external quality control programs.

Statistical analysis

Multiple stepwise regression analysis was performed with the following covariates in the model: age, BMI and sex. Groups were compared by Mann-Whitney test, where appropriate. The diagnostic accuracy of GARG and ITT was determined by ROC analysis and its corresponding area under the curve (AUC). That way, we aimed to establish cutoffs (with a specificity of $\geq 95\%$ for GHD) under special consideration of the most relevant anthropometric factors. Statistical significance was taken as $P < 0.05$. Results are expressed as the mean $\pm$ standard error of the mean (S.E.M.) unless otherwise stated. GraphPad Prism 6.0 software (GraphPad Software) and JMP10 (SAS Institute Inc., Cary, NC, USA) were used for statistical calculations.

Results

Influence of age, BMI and sex on GH peak levels in healthy subjects

The control group comprised 200 healthy subjects: 102 males (age 39.7 ± 1.5 years (range 19–75 years), BMI $27.8 \pm 0.5 \text{ kg/m}^2$ (range 19.0–40.9 kg/m^2)) and 98 females (age 39.2 ± 1.3 years (range 18–68 years), BMI $27.7 \pm 0.6 \text{ kg/m}^2$ (range 18.0–46.5 kg/m^2)). If male and female controls were compared, no significant differences between mean age and mean BMI were detected. Multiple stepwise regression analysis revealed that both BMI (21%, $P < 0.0001$) and sex (20%, $P < 0.0001$) accounted for most of the variability of GH peak levels during the GARG test, whereas the influence of age (5%, $P < 0.001$) was much less dominant. Mean peak GH levels were significantly higher in females compared with males ($27.4 \pm 1.73 \text{ ng/mL}$ vs

Figure 1

Peak GH levels during the GARG test in 200 healthy subjects, stratified by sex and BMI. Dark lines represent the means for each group. GH, growth hormone.

$13.7 \pm 0.95 \text{ ng/mL}$, $P < 0.0001$) and higher in lean ($34.1 \pm 2.46 \text{ ng/mL}$ in females vs $20.7 \pm 1.67 \text{ ng/mL}$ in males, $P < 0.0001$) compared with overweight ($31.4 \pm 3.38 \text{ ng/mL}$ in females vs $12.1 \pm 1.34 \text{ ng/mL}$ in males, $P < 0.0001$) and obese subjects ($14.4 \pm 1.52 \text{ ng/mL}$ in females vs $7.19 \pm 0.68 \text{ ng/mL}$ in males, $P < 0.0001$) (Fig. 1).

As changes in sex hormones during menopause could be interfering and potentially explain some of the sex differences in peak GH levels, we performed additional analyses in 139 healthy subjects (age 18–50 years) of whom females were presumably premenopausal and 60 subjects (age > 50 years) of whom females were presumably menopausal. Peak GH remained significantly higher in 67 females between 18 and 50 years of age compared with 72 males ($29.8 \pm 2.19 \text{ ng/mL}$ in females vs $15.2 \pm 1.14 \text{ ng/mL}$ in males, $P < 0.0001$), and in 31 females who were > 50 years old compared with 29 males ($22.1 \pm 2.5 \text{ ng/mL}$ in females vs $9.8 \pm 1.52 \text{ ng/mL}$ in males, $P < 0.0001$).

Patient cohort

Out of the 87 subjects with hypothalamic–pituitary diseases (51 males, 36 females; age 48.7 ± 1.5 years (range 18–74 years), BMI $27.6 \pm 0.6 \text{ kg/m}^2$ (range 16.7–43.2 kg/m^2)), 17 patients were investigated specifically due to suspicion of pituitary insufficiency (7 $\times$ clinical symptoms; 3 $\times$ traumatic brain injury; 2 $\times$ congenital panhypopituitarism; each 1 $\times$ cerebral lymphoma after cranial radiotherapy, congenital isolated growth hormone deficiency, empty sella syndrome,

Table 2 Sex-specific BMI-dependent cutoffs in the combined arginine-GHRH test for GH as measured by the IDS iSYS assay with a specificity of $\geq 95\%$ for GHD.

	Overall	Males			Females		
		Lean (BMI <25 kg/m ²)	Overweight (BMI ≥ 25 –<30 kg/m ²)	Obese (BMI ≥ 30 kg/m ²)	Lean (BMI <25 kg/m ²)	Overweight (BMI ≥ 25 –<30 kg/m ²)	Obese (BMI ≥ 30 kg/m ²)
Cutoff (ng/mL)	3.9	6.5	3.5	2.2	9.7	8.5	4.4
Sensitivity (%)	86	100	92	91	83	100	67
Specificity (%)	95	95	96	97	96	97	97
ROC AUC	0.97	1.00	0.99	0.99	0.97	1.00	0.95
P value	<0.0001	<0.0001	<0.0001	<0.0001	<0.0005	<0.0001	<0.001

BMI, body mass index; GH, growth hormone; GHD, growth hormone deficiency; ROC, receiver-operating characteristics; AUC, area under the curve.

Sheehan's syndrome and subarachnoid hemorrhage), whereas the remaining 70 patients had been tested because of proven sellar masses. Forty-seven of the latter subjects had already been surgically treated (30× nonfunctioning pituitary adenoma; 8× ACTH secreting pituitary adenoma; 3× meningioma; 2× prolactinoma; each 1× arachnoid cyst, colloid cyst, craniopharyngioma and growth hormone-secreting pituitary adenoma), whereas 23 patients had not yet been operated (8× craniopharyngioma; 7× nonfunctioning pituitary adenoma; 4× prolactinoma; 2× meningioma; each 1× astrocytoma and suprasellar myxoid chondroma). Fifty-one of the 87 patients had MPHD as defined previously and were therefore assumed GHD, whereas 36 patients had ≤ 1 PHD and were therefore considered to be GHS. The median number of PHD was 3 in the GHD and 1 in the GHS group. Clinical characteristics of patients are provided in Table 1.

Establishment of new cutoffs for peak GH levels during the GHRH plus arginine test by comparison of growth hormone-deficient patients and control subjects

Considering the significant influence of BMI and sex for the GARG in normal subjects, we aimed to establish diagnostic cutoffs that were adjusted accordingly. Comparison of peak GH levels during the GARG test of patients with MPHD to those of a control group (consisting of patients with PHD <2 and healthy subjects) allowed calculation of cutoffs for peak GH (either as an overall threshold or as BMI-dependent thresholds with at least 95% specificity for GHD, with the latter being separately provided for both sexes). As shown in Table 2 as well as in Fig. 2, the overall cutoff for both sexes was 3.9 ng/mL (sensitivity 86% (95% confidence interval (CI) 74–94%), specificity 95% (95% CI 92–98%));

the BMI-adjusted thresholds were 6.5, 3.5 and 2.2 ng/mL for males and 9.7, 8.5, and 4.4 ng/mL for females (always for lean, overweight and obese subjects respectively). Of note, our group of patients with less than 2 PHD may contain some subjects with isolated GHD, which could influence our analysis. We therefore repeated our ROC analysis focusing on patients with MPHD in comparison to healthy

Figure 2

Peak GH levels and corresponding ROC curves during (upper panel) GARG test and (lower panel) ITT. The broken lines indicate the overall GH cutoff (3.9 ng/mL during the GARG test, 2 ng/mL during the ITT). AUC, area under the curve; C, healthy subjects; GARG, growth hormone-releasing hormone plus arginine; GH, growth hormone; GHD, growth hormone deficiency; GHS, growth hormone sufficiency; ITT, insulin tolerance test.

controls alone. Interestingly, the results were very similar: the overall cutoff for both sexes was 3.8 ng/mL (sensitivity 86% (95% CI 73–94%), specificity 95% (95% CI 92–98%)).

Diagnostic performance of the insulin tolerance test

Forty-four of the 87 patients (51%) also underwent an ITT as additional reference test. ROC analysis suggested a GH cutoff of 2.0 ng/mL that categorized 29 patients as GHD and 15 patients as GHS (sensitivity 95% and specificity 93%). Hence, two cases were discordant to the classification by the degree of pituitary impairment (i.e., 1 GHS patient with 3 PHD and 1 GHD patient with 0 PHD).

If the 'classical' GH cutoff of 3.0 ng/mL was applied, 32 patients were regarded as GHD and 12 patients as GHS. Using ITT, 1 individual with ≥ 2 PHD was considered GHS, whereas 4 individuals (each with 0 PHD) were classified as GHD. As a result, sensitivity remained at 95%, whereas specificity dropped to 73%.

Discussion

The current study established reference data for peak GH during the GARG test, using a modern automated immunoassay, which had been developed in compliance with recent consensus guidelines and specifically measures the 22 kDa isoform of human GH. To establish diagnostic cutoffs, prospectively recruited subjects from the general population underwent careful questioning and examination, thereby including only well-defined healthy controls. Multiple stepwise regression analysis was performed to determine the influence of various anthropometric variables on GH, identifying BMI and sex as the most relevant parameters. Consequently, adjusted cutoffs with at least 95% specificity for GHD were derived from ROC analysis, investigating patients with proven hypothalamic–pituitary diseases (using the number of PHD as gold standard).

Although measurement of GH and IGF-I is a cornerstone of the diagnostic and therapeutic monitoring of diseases associated with an impaired somatotroph function, considerable difficulties exist with respect to the comparability of currently available measurement techniques and their respective analytical performance (20, 24). Therefore, the recent consensus guidelines on GH assay performance have called for method-specific cutoffs for each GH assay (20). Noteworthy, the need to establish cutoffs not only for each individual hormone and dynamic test but also for each of the different

measurement techniques used to quantify the hormones has already been demonstrated for other hormones (25). The cutoff data presented here apply for the specific GH assay used, and conversion of data obtained by one GH assay into another GH assay is explicitly discouraged by the guidelines. However, especially in a pediatric population where ethical reasons might prohibit the re-establishment of cutoffs with each new assay, it is helpful to know that a recent publication has proposed formulas allowing a comparison of GH concentrations obtained by the IDS iSYS GH assay to other commercially available immunoassays (26). To date, robust reference ranges for other parameters associated with somatotroph function (IGF1 and IGF-BP3) measured on the same analytical platform have already been established (27, 28, 29), but the present study is the first to provide assay-specific cutoffs for GH stimulation tests for this assay.

In patients with suspicion for GHD, current guidelines regard testing with well-established dynamic endocrine procedures as mandatory (1, 2, 7). Over the last years, the GARG test has repeatedly been described as a reliable alternative to the traditional gold standard ITT, which is nowadays regarded as unpleasant, cumbersome and potentially life-threatening (1, 2, 5, 7, 17, 18, 19).

Of note, a significant relationship between anthropometric parameters and GH levels during dynamic endocrine testing has been described, e.g. for the oral glucose tolerance test (30) and the GARG test (5, 13, 14). With respect to the latter test, BMI-dependent cutoffs for peak GH have been suggested, mainly to overcome overlapping GH responses between GHD and obese patients (13, 14). We also identified BMI as the most relevant confounder in our current analysis, accounting for 21% of the variability of peak GH during the GARG test. Unfortunately, though, a further analysis of the underlying pathophysiological mechanism for this finding was not feasible, as additional parameters of metabolic disturbance (e.g. visceral fat mass, visceral fat mass, fasting insulin levels and waist circumference) were not routinely determined.

Interestingly, multiple stepwise regression analysis revealed that sex had a very comparable effect to BMI (i.e., accounting for 20% of peak GH variability). We therefore decided to establish combined BMI- and sex-adjusted cutoffs rather than various cutoffs only adjusted for one of these potential confounders.

It has to be pointed out that these newly established thresholds were markedly lower in males (6.5, 3.5 and 2.2 ng/mL for lean, overweight and obese subjects) than those in females (9.7, 8.5 and 4.4 ng/mL) and not well

comparable to BMI-dependent thresholds published by Corneli *et al.* (i.e., 11.5, 8.0 and 4.2 ng/mL for males and females combined) or Colao *et al.* (who did not observe a gender effect and provided age-specific cutoffs instead, ranging from 12.0 to 21.5 ng/mL, 11.1 to 16.5 ng/mL and 4.0 to 15.0 ng/mL respectively) (threshold values in both studies always given for lean, overweight and obese subjects) (13, 14). As discussed previously, variations in the assay methods used in the former studies certainly contribute to the differences in concentrations reported. However, these differences might also point to other confounders in the respective cohorts investigated (e.g. anthropometric factors other than BMI, which appears to have a relatively robust inverse correlation to GH). With respect to age, some groups have reported that peak GH levels during the GARG test were substantially decreasing in the elderly (14), whereas others did not observe such a decline (addressing either the GARG test only (31) or several GH stimulation tests at the same time) (5). Divergent results have also been published for the role of sex, with reports ranging from no (14, 31) to a significant (5) impact on peak GH levels during the GARG test. Our study confirms the influence of sex, which appears to be plausible in view of the known, significantly stimulating effect of estradiol on GH secretion (32, 33). Our current multiple stepwise regression analysis and the results published by Biller *et al.* (5) strongly suggest that sex may have a more dominant effect on GH peak levels than age, with GH peak values more than twice as high in females compared with males. As a result, we believe that the use of cutoffs adjusted for BMI and sex could be a reasonable approach to improve diagnostic accuracy.

Our study has limitations. For instance, although we included a comparably large cohort of healthy subjects, our patient cohort was smaller than that in some of the previous studies on this topic. Although the rarity of GH-related disorders needs to be considered, larger investigations are certainly needed to better define the impact of sex and BMI on patients. Such studies should ideally be designed as large prospective multicenter studies to also account for other potential confounders. Nevertheless, all of our patients were carefully clinically reviewed, with special emphasis on symptoms and biochemical signs suggestive of (additional) PHD. In line with current recommendations, we strongly believe that this approach allows for a reliable identification of hypopituitarism. This assumption is also supported by the good agreement between clinical annotations and the corresponding ITT in a subgroup of the patients. Another drawback of our current analysis is that previous

studies of GARG have demonstrated limited sensitivity to diagnose GHD in patients with prior irradiation (34, 35). It has been speculated that the GH response to arginine may be somehow resistant to the effects of radiation (36). As our patient cohort contained only 2 patients after radiation, we believe that the results are not biased by this treatment, but like to emphasize that the results may not apply to patients after cranial irradiation. Furthermore, the GARG test itself has certain pitfalls. For instance, GHRH is not easily available, making the test quite cumbersome. Moreover, as many potential confounders need to be considered, different cutoffs values have to be memorized (which are only valid for the specific assay applied). Consequently, each center needs to establish its own normative data.

In conclusion, here we provide method-specific cutoffs for a new automated immunoassay specifically measuring the 22 kDa isoform of human GH. Furthermore, we defined the effects of anthropometric factors on peak GH levels during the GARG test. We demonstrated that BMI and sex account for most of the variability in peak GH, thereby having important impact on the diagnosis of GHD. Consequently, the diagnostic accuracy of the GARG test is significantly improved by application of BMI- and sex-adjusted cutoffs.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported. Stephan Petersenn has received research support from IDS, and Martin Bidlingmaier has received research support and consultancy fees from IDS.

Funding

Timo Deutschbein received a grant from the Interdisciplinary Center for Clinical Research (IZKF) of the University of Würzburg (grant number Z-2/57).

Author contribution statement

Timo Deutschbein and Stephan Petersenn conducted the statistical analyses. All authors collected samples, provided patient data and wrote the manuscript.

References

- 1 Cook DM, Yuen KC, Biller BM, Kemp SF, Vance ML & American Association of Clinical Endocrinologists. American Association of Clinical Endocrinologists medical guidelines for clinical practice for growth hormone use in growth hormone-deficient adults and transition patients – 2009 update: executive summary of recommendations. *Endocrine Practice* 2009 **15** 580–586. (doi:10.4158/EP.15.6.580)
- 2 Molitch ME, Clemmons DR, Malozowski S, Merriam GR, Vance ML & Endocrine Society. Evaluation and treatment of adult growth

- hormone deficiency: an Endocrine Society clinical practice guideline. *Journal of Clinical Endocrinology and Metabolism* 2011 **96** 1587–1609. (doi:10.1210/jc.2011-0179)
- 3 Bates AS, Van't Hof W, Jones PJ & Clayton RN. The effect of hypopituitarism on life expectancy. *Journal of Clinical Endocrinology and Metabolism* 1996 **81** 1169–1172. (doi:10.1210/jcem.81.3.8772595)
  - 4 Svensson J, Bengtsson BA, Rosén T, Odén A & Johannsson G. Malignant disease and cardiovascular morbidity in hypopituitary adults with or without growth hormone replacement therapy. *Journal of Clinical Endocrinology and Metabolism* 2004 **89** 3306–3312. (doi:10.1210/jc.2003-031601)
  - 5 Biller BM, Samuels MH, Zagar A, Cook DM, Arafah BM, Bonert V, Stavrou S, Kleinberg DL, Chipman JJ & Hartman ML. Sensitivity and specificity of six tests for the diagnosis of adult GH deficiency. *Journal of Clinical Endocrinology and Metabolism* 2002 **87** 2067–2079. (doi:10.1210/jcem.87.5.8509)
  - 6 Hartman ML, Crowe BJ, Biller BMK, Ho KKY, Clemmons DR, Chipman JJ, HyposCCS Advisory Board & U.S. HypoCCS Study Group. Which patients do not require a growth hormone (GH) stimulation test for the diagnosis of adult GH deficiency? *Journal of Clinical Endocrinology and Metabolism* 2002 **87** 477–485. (doi:10.1210/jcem.87.2.8216)
  - 7 Ho KK & 2007 GH Deficiency Consensus Workshop Participants. Consensus guidelines for the diagnosis and treatment of adults with GH deficiency II: a statement of the GH Research Society in association with the European Society for Pediatric Endocrinology, Lawson Wilkins Society, European Society of Endocrinology, Japan Endocrine Society, and Endocrine Society of Australia. *European Journal of Endocrinology* 2007 **157** 695–700. (doi:10.1530/EJE-07-0631)
  - 8 Rahim A, Toogood AA & Shalet SM. The assessment of growth hormone status in normal young adult males using a variety of provocative agents. *Clinical Endocrinology* 1996 **45** 557–562. (doi:10.1046/j.1365-2265.1996.00855.x)
  - 9 Aimaretti G, Baffoni C, DiVito L, Bellone S, Grottoli S, Maccario M, Arvat E, Camanni F & Ghigo E. Comparisons among old and new provocative tests of GH secretion in 178 normal adults. *European Journal of Endocrinology* 2000 **142** 347–352. (doi:10.1530/eje.0.1420347)
  - 10 Hoeck HC, Vestergaard P, Jakobsen PE, Falhof J & Laurberg P. Diagnosis of growth hormone (GH) deficiency in adults with hypothalamic-pituitary disorders: comparison of test results using pyridostigmine plus GH-releasing hormone (GHRH), clonidine plus GHRH, and insulin-induced hypoglycemia as GH secretagogues. *Journal of Clinical Endocrinology and Metabolism* 2000 **85** 1467–1472. (doi:10.1210/jcem.85.4.6538)
  - 11 Popovic V, Leal A, Micic D, Koppeschaar HP, Torres E, Paramo C, Obradovic S, Dieguez C & Casanueva FF. GH-releasing hormone and GH-releasing peptide-6 for diagnostic testing in GH-deficient adults. *Lancet* 2000 **356** 1137–1142. (doi:10.1016/S0140-6736(00)02755-0)
  - 12 Leong KS, Walker AB, Martin I, Wile D, Wilding J & MacFarlane IA. An audit of 500 subcutaneous glucagon stimulation tests to assess growth hormone and ACTH secretion in patients with hypothalamic-pituitary disease. *Clinical Endocrinology* 2001 **54** 463–468. (doi:10.1046/j.1365-2265.2001.01169.x)
  - 13 Corneli G, Di Somma C, Baldelli R, Rovere S, Gasco V, Croce CG, Grottoli S, Maccario M, Colao A, Lombardi G *et al.* The cut-off limits of the GH response to GH-releasing hormone-arginine test related to body mass index. *European Journal of Endocrinology* 2005 **153** 257–264. (doi:10.1530/eje.1.01967)
  - 14 Colao A, Di Somma C, Savastano S, Rota F, Savanelli MC, Aimaretti G & Lombardi G. A reappraisal of diagnosing GH deficiency in adults: role of gender, age, waist circumference, and body mass index. *Journal of Clinical Endocrinology and Metabolism* 2009 **94** 4414–4422. (doi:10.1210/jc.2009-1134)
  - 15 Alaoui B, Mann K & Petersenn S. Diagnosis of adrenal insufficiency using the GHRP-6 test: comparison with the insulin tolerance test in patients with hypothalamic-pituitary-adrenal disease. *Hormone and Metabolic Research* 2010 **42** 198–203. (doi:10.1055/s-0029-1243184)
  - 16 Berg C, Meinel T, Lahner H, Yucel A, Mann K & Petersenn S. Diagnostic utility of the glucagon stimulation test in comparison to the insulin tolerance test in patients following pituitary surgery. *European Journal of Endocrinology* 2010 **162** 477–482. (doi:10.1530/EJE-09-0824)
  - 17 Chanson P, Cailleux-Bounacer A, Kuhn JM, Weryha G, Chabre O, Borson-Chazot F, Dubois S, Vincent-Dejean C, Brue T, Fedou C *et al.* Comparative validation of the growth hormone-releasing hormone and arginine test for the diagnosis of adult growth hormone deficiency using a growth hormone assay conforming to recent international recommendations. *Journal of Clinical Endocrinology and Metabolism* 2010 **95** 3684–3692. (doi:10.1210/jc.2010-0295)
  - 18 Markkanen HM, Pekkarinen T, Välimäki MJ, Alfthan H, Hämäläinen E & Stenman UH. Comparison of two growth hormone stimulation tests and their cut-off limits in healthy adults at an outpatient clinic. *Growth Hormone and IGF Research* 2013 **23** 165–169. (doi:10.1016/j.ghir.2013.06.002)
  - 19 Andersen M. The robustness of diagnostic tests for GH deficiency in adults. *Growth Hormone and IGF Research* 2015 **25** 108–114. (doi:10.1016/j.ghir.2015.03.001)
  - 20 Clemmons DR. Consensus statement on the standardization and evaluation of growth hormone and insulin-like growth factor assays. *Clinical Chemistry* 2011 **57** 555–559. (doi:10.1373/clinchem.2010.150631)
  - 21 Manolopoulou J, Alami Y, Petersenn S, Schopohl J, Wu Z, Strasburger J & Bidlingmaier M. Automated 22-kDa growth hormone-specific assay without interference from pegvisomant. *Clinical Chemistry* 2012 **58** 1446–1456. (doi:10.1373/clinchem.2012.188128)
  - 22 Toogood AA, Beardwell CG & Shalet SM. The severity of growth hormone deficiency in adults with pituitary disease is related to the degree of hypopituitarism. *Clinical Endocrinology* 1994 **41** 511–516. (doi:10.1111/j.1365-2265.1994.tb02583.x)
  - 23 Bates AS, Evans AJ & Clayton RN. Assessment of GH status in adults with GH deficiency using serum growth hormone, serum insulin-like growth factor-I and urinary growth hormone excretion. *Clinical Endocrinology* 1995 **42** 425–430. (doi:10.1111/j.1365-2265.1995.tb02652.x)
  - 24 Bidlingmaier M & Freda PU. Measurement of human growth hormone by immunoassays: current status, unsolved problems and clinical consequences. *Growth Hormone and IGF Research* 2009 **20** 19–25. (doi:10.1016/j.ghir.2009.09.005)
  - 25 Deutschbein T & Petersenn S. Screening for Cushing's syndrome: new immunoassays require adequate normative data. *Hormone and Metabolic Research* 2013 **45** 118–123. (doi:10.1055/s-0032-1331745)
  - 26 Müller A, Scholz M, Blankenstein O, Binder G, Pfäffle R, Körner A, Kiess W, Heider A, Bidlingmaier M, Thiery J *et al.* Harmonization of growth hormone measurements with different immunoassays by data adjustment. *Clinical Chemistry and Laboratory Medicine* 2011 **49** 1135–1142. (doi:10.1515/cclm.2011.201)
  - 27 Bidlingmaier M, Friedrich N, Emeny RT, Spranger J, Wolthers OD, Roswall J, Körner A, Obermayer-Pietsch B, Hübener C, Dahlgren J *et al.* Reference intervals for insulin-like growth factor-1 (igf-I) from birth to senescence: results from a multicenter study using a new automated chemiluminescence IGF-I immunoassay conforming to recent international recommendations. *Journal of Clinical Endocrinology and Metabolism* 2014 **99** 1712–1721. (doi:10.1210/jc.2013-3059)
  - 28 Friedrich N, Wolthers OD, Arafat AM, Emeny RT, Spranger J, Roswall J, Kratzsch J, Grabe HJ, Hübener C, Pfeiffer AF *et al.* Age and sex specific reference intervals across lifespan for insulin-like growth factor binding protein 3 (IGFBP-3) and the IGF-I/IGFBP-3 ratio measured by new automated chemiluminescence assays. *Journal of Clinical Endocrinology and Metabolism* 2014 **99** 1675–1686. (doi:10.1210/jc.2013-3060)

- 29 Wagner IV, Paetzold C, Gausche R, Vogel M, Koerner A, Thierry J, Arsene CG, Henrion A, Guettler B, Keller E *et al.* Clinical evidence-based cutoff limits for GH stimulation tests in children with a backup of results with reference to mass spectrometry. *European Journal of Endocrinology* 2014 **171** 389–397. (doi:10.1530/EJE-14-0165)
- 30 Arafat AM, Möhlig M, Weickert MO, Perschel FH, Purschwitz J, Spranger J, Strasburger CJ, Schöfl C & Pfeiffer AF. Growth hormone response during oral glucose tolerance test: the impact of assay method on the estimation of reference values in patients with acromegaly and in healthy controls, and the role of gender, age, and body mass index. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** 1254–1262. (doi:10.1210/jc.2007-2084)
- 31 Ghigo E, Aimaretti G, Arvat E & Camanni F. Growth hormone releasing hormone combined with arginine or growth hormone secretagogues for the diagnosis of growth hormone deficiency in adults. *Endocrine* 2001 **15** 29–38. (doi:10.1385/ENDO:15:1:029)
- 32 Ho KY, Evans WS, Blizzard RM, Veldhuis JD, Merriam GR, Samojlik E, Furlanetto R, Kaiser DL & Thorner MO. Effects of sex and age on the 24-hour profile of growth hormone secretion in man: importance of endogenous estradiol concentrations. *Journal of Clinical Endocrinology and Metabolism* 1987 **64** 51–58. (doi:10.1210/jcem-64-1-51)
- 33 Faria AC, Bekenstein LW, Booth RA Jr, Vaccaro VA, Asplin CM, Veldhuis JD, Thorner MO & Evans WS. Pulsatile growth hormone release in normal woman during the menstrual cycle. *Clinical Endocrinology* 1992 **36** 591–596. (doi:10.1111/j.1365-2265.1992.tb02270.x)
- 34 Björk J, Link K & Erfurth EM. The utility of the growth hormone (GH) releasing hormone-arginine test for diagnosing GH deficiency in adults with childhood acute lymphoblastic leukemia treated with cranial irradiation. *Journal of Clinical Endocrinology and Metabolism* 2005 **90** 6048–6054. (doi:10.1210/jc.2005-0304)
- 35 Ham JN, Ginsberg JP, Hendell CD & Moshang T Jr. Growth hormone releasing hormone plus arginine stimulation testing in young adults treated in childhood with cranio-spinal radiation therapy. *Clinical Endocrinology* 2005 **62** 628–632. (doi:10.1111/j.1365-2265.2005.02272.x)
- 36 Lissett CA, Saleem S, Rahim A, Brennan BM & Shalet SM. The impact of irradiation on growth hormone responsiveness to provocative agents is stimulus dependent: results in 161 individuals with radiation damage to the somatotrophic axis. *Journal of Clinical Endocrinology and Metabolism* 2001 **86** 663–668. (doi:10.1210/jcem.86.2.7235)

Received 7 August 2016

Revised version received 24 November 2016

Accepted 8 December 2016