DIAGNOSIS OF ENDOCRINE DISEASE

Bone turnover markers: are they clinically useful?

Richard Eastell¹, Tom Pigott², Fatma Gossiel¹, Kim E Naylor¹, Jennifer S Walsh¹ and Nicola F APeel²

¹Academic Unit of Bone Metabolism, University of Sheffield, Sheffield, UK and ²Metabolic Bone Centre, Sheffield Teaching Hospitals NHS Trust, Sheffield, UK

Correspondence should be addressed to R Eastell
Email
r.eastell@sheffield.ac.uk

Abstract

Bone turnover markers (BTMs) are useful in clinical practice as they are inexpensive, and they have proven useful for treatment monitoring and identification of poor adherence. BTMs cannot be used in individual patients for identifying accelerated bone loss or an increase in fracture risk or in deciding on the optimal therapy. They are useful for monitoring both anti-resorptive and anabolic treatment. Response can be defined as a result that exceeds an absolute target, or by a change greater than the least significant change; if such a response is not present, then poor compliance or secondary osteoporosis are likely causes. A baseline BTM measurement is not always made; in that case, a value of BTM on anti-resorptive treatment that is low or low normal or above the reference interval for anabolic therapy may be taken to indicate a satisfactory response. We provide an approach to using these bone turnover markers in clinical practice by describing algorithms for anti-resorptive and anabolic therapy and describing the changes we observe in the clinical practice setting.

European Journal of Endocrinology
(2018) **178**, R19–R31

Introduction

The fractures that result from osteoporosis are a major public health problem (1). Osteoporosis is characterised by reduced bone mass and microarchitectural deterioration leading to increased bone fragility and may be diagnosed by measurement of bone mineral density (BMD) using dual-energy X-ray absorptiometry. A BMD value at the spine or hip that is 2.5 s.d. or more below the average value for healthy young women is considered to represent osteoporosis, according to the WHO Working Group. Several treatments have been licensed for use in osteoporosis that are effective in reducing the risk of fracture.

This article focuses on the use of bone turnover markers (BTM) in osteoporosis. BTM can be measured in serum, plasma and urine, and their levels relate to the activity of osteoblasts (bone formation markers) and

osteoclasts (bone resorption markers). Bone formation markers include proteins that are specific to bone (osteocalcin) or not so specific to bone such as fragments of type I procollagen released during formation of type I collagen (N-propeptide of type I collagen, PINP) and the bone isoform of alkaline phosphatase (bone ALP). Bone resorption markers include fragments released from the telopeptide (end) region of type I collagen following its enzymatic degradation, including the N-telopeptide of type I collagen (NTX) and the C-telopeptide of type I collagen (CTX), deoxypyridinoline and the enzyme tartrate-resistant acid phosphatase (Table 1).

In women, the BTMs increase after the menopause and in other situations of accelerated bone loss. In men, there is little increase with age. In cohort studies of women (but not of men), the higher the BTM, the more rapid the

Review	R Eastell and others	Bone turnover markers	178:1	R20
--------	----------------------	-----------------------	-------	-----

Table 1 Bone turnover markers commonly used in clinical practice.

Bone formation	N-propeptide of type I collagen, PINP*	Serum or plasma	Radioimmunoassay, automated assay (Roche Cobas, IDS iSYS)
	Osteocalcin (OC)	Serum or plasma	ELISA, automated assay (Roche Cobas, IDS iSYS)
Bone resorption	Bone alkaline phosphatase (bone ALP)	Serum	IRMA, ELISA, automated assay (IDS iSYS)
	C-terminal telopeptide of type I collagen (CTX)*	Serum, plasma or urine	ELISA, automated assay (Roche Cobas, IDS iSYS)
	N-terminal telopeptide of type I collagen (NTX)	Serum, plasma or urine	ELISA, automated assay (Ortho Clinical Diagnostics)
	Deoxypyridinoline (DPD)	Urine	HPLC for total, ELISA for free
	Tartrate-resistant acid phosphatase, 5b	Serum or plasma	ELISA, automated assay (IDS iSYS)

bone loss and the greater the risk of fracture. Thus, the measurement of BTM may have clinical relevance to the individual. Currently, the main clinical use for BTM is for the monitoring of response to therapy. A typical goal of therapy might be to lower BTM to values found in women before the menopause.

History, assays and validation

Bone histomorphometry is the gold standard for assessment of bone turnover, but it is invasive, cannot be repeated many times in an individual and requires specialist laboratory interpretation. Bone turnover can also be quantified with calcium balance and kinetic studies, but they are time-consuming, use radio-isotopes and again need specialist interpretation.

Therefore, for clinical use in large numbers of patients, there is a need for measures that can be made on easily accessible samples (single measurements of blood or urine), inexpensively, do not require time-consuming specialist processing and give results that can be interpreted by non-specialist health care practitioners.

The BTM that were developed initially were not bone specific (for example, hydroxyproline and total alkaline phosphatase), the assays were technically challenging (HPLC for total deoxypyridinoline) and therefore costly and difficult to implement widely. The newer BTM are more bone specific and the use of enzyme-linked immunosorbent assays (ELISA) and autoanalyser techniques have made them widely available and more affordable (Table 1).

Although assays for total alkaline phosphatase (ALP) were available in the 1920s, only about half of the total ALP is from bone. Hydroxyproline assays were developed in the 1950s, but again were not specific for bone and were laborious and dangerous (they resulted in explosions). There were significant developments in the 1980s and 1990s with assays for pyridinium crosslinks

(deoxypyridinoline and pyridinoline), bone ALP, PINP and osteocalcin and progression from HPLC to immunoassays.

The introduction of automated immunoassay analysers in 2000 was a major technical advance. These are widely used in clinical practice for measuring many analytes, including hormones, as well as BTM, and they do so with high precision (CV less than 5%) and reliability.

BTM have been validated against gold standard methods for studying bone turnover such as a comparison with tracer kinetics and bone histomorphometry, both in health (2) and in response to osteoporosis treatments (3). Currently used BTMs were evaluated in a study of 370 women with osteoporosis (4). BTM were assessed against dynamic histomorphometry of iliac crest biopsies. There were weak-to-moderate correlations (highest *r* value was 0.41) between the bone formation markers PINP or bone ALP and bone formation estimates and between CTX and bone resorption estimates.

Practical aspects

There are different requirements for the use of BTM in clinical practice compared to the research setting. In clinical practice, patients may attend appointments at any time of day, and there may be a delay before samples are transported to the laboratory. Patients often have complex medical problems and take multiple medications. In this context, some properties of BTM present challenges to their clinical use.

BTM need to be measured reliably and easily, be locally accessible, inexpensive and be unaffected by the time of day the samples are obtained.

The bone resorption markers show a strong circadian rhythm and decrease shortly after feeding. Thus, it is recommended that blood samples for CTX be drawn from the patient following an overnight fast between 07:30 and 10:00 (5). The sample can be collected as EDTA plasma or serum; EDTA is preferable if the sample cannot

be processed within 2 h. The sample can be stored frozen until measured; if the storage is likely to be more than 12 weeks, it is recommended that this is at -70 to -80°C (6). It is recommended for urinary NTX that the sample is taken as the second morning void, that excessive fluid consumption is avoided and preservative is not added. Variability can be further reduced by obtaining urine samples on three consecutive days, pooling the samples and just making one measurement. When measuring bone resorption markers (NTX, CTX, DPD), it is usual to measure urinary creatinine and to express the result as the BTM-to-creatinine ratio to correct for urinary dilution.

For bone formation markers, there is a weaker circadian rhythm and so the sample can be drawn at any time of the day (5). Serum or plasma should be measured the same day or stored in the freezer until measured (5). EDTA plasma should not be used for bone ALP measurement (5). Osteocalcin is affected by haemolysis, which can lead to a falsely low result. BTM, especially bone formation markers, are increased following fracture and are affected by some medical conditions and treatments, as discussed in the following sections (5).

Choice of BTM

It is logical to include a bone resorption and a bone formation assay when evaluating bone turnover. The choice of BTM will be determined by local availability and cost. It will also be determined by the clinical picture. Thus, in chronic kidney disease, the markers that are usually excreted by the kidney circulate at very high levels and so markers that are not excreted by the kidney are best used, e.g. bone ALP and intact PINP. In the evaluation of glucocorticoid treatment on bone, markers that are sensitive to the bone effects of these drugs may be most useful, e.g. osteocalcin and PINP, which are affected in a dose-dependent manner. These markers are not, however, useful to evaluate the effect of anti-resorptive therapy in these patients.

The IOF has proposed serum CTX and PINP as the two reference markers; they propose that all research studies should include these two at a bare minimum (7), but for clinical practice, it may suffice to have one marker only.

Potential clinical utility

BTM have proven useful in evaluating the relationship between bone turnover and rates of bone loss, fracture risk

and treatment effect in groups of patients using statistical approaches including linear and logistic regression or repeated measures analysis of variance. However, different statistical approaches are needed to evaluate the utility of BTM in the individual. These include tests of diagnostic accuracy (sensitivity, specificity) and assessment of the least significant change to identify response. The least significant change is a change (expressed in absolute units or percentage) that is beyond the day-to-day changes observed in untreated individuals. It is a statistical approach and is often defined and allows for change up or down and beyond the change expected 95% of the time. It is widely used in clinical chemistry where it is also referred to in some texts as the 'reference critical difference' (8). It is calculated as 2.77 times the coefficient of variation; the latter includes both assay and within-subject variability.

Prediction of bone loss

High bone turnover is associated with more rapid bone loss in postmenopausal women (9) and BTMs have been studied in evaluation of this relationship. Higher BTMs are associated with bone loss from both trabecular and cortical bone at the hip and also relate to greater periosteal expansion in the femoral neck (10). The assessment is improved by making more than one BTM measurement (11). Estimation of the rate of bone loss in a postmenopausal woman when deciding about her need for anti-resorptive treatment would potentially be useful. Unfortunately, the association between BTM and bone loss is not sufficient to classify individuals reliably by their BTM level (12).

Prediction of fracture

It would also be of interest to estimate the risk of fracture in the individual postmenopausal woman when deciding about the need for anti-resorptive treatment, and high bone turnover is associated with increased risk of several types of fracture in both men and women (7, 13). In a recent meta-analysis of 6 studies that had measurements of bone resorption (CTX) and bone formation (PINP), the hazard ratio per SD increase was similar for CTX (1.18, 95% CI 1.05–1.34) and PINP (1.23, 95% CI 1.09–1.39) (14). These results were not adjusted for BMD. However, not all studies find an association between BTM and fracture risk (10) and the FRAX Position Development Conference members were unable to find sufficient

Review	R Eastell and others	Bone turnover markers	178:1	R22
--------	----------------------	-----------------------	-------	-----

evidence for inclusion of BTM into the FRAX fracture risk prediction algorithm (15).

Selection of therapy

Intuitively, we would like to choose our therapies based on the mechanism of bone loss underlying the osteoporosis. Thus, we might use anti-resorptive therapies (bisphosphonates, raloxifene, denosumab) in patients with high BTM and anabolic therapies (teriparatide, abaloparatide) in patients with low BTM. Unfortunately, this approach is not supported by the results of clinical trials. In the Fracture Intervention Trial, treatment with alendronate was more effective at reducing non-vertebral fracture in those women with higher PINP, but this was not true for other BTM or other fracture types (16). Similarly, the baseline BTM did not predict the fracture benefit with teriparatide (17). In general, a low PINP is associated with lower rates of bone loss and lower response to zoledronic acid (18). Further research is needed.

Treatment used for osteoporosis

Despite having several treatments that reduce the risk of fracture in osteoporosis, it is well established that adherence to these treatments can be poor, especially in the case of oral bisphosphonates for which the dosing instructions are complex. There is therefore a need to identify optimal treatment response in individual patients. It has been proposed that treatment failure may be considered if two or more fractures occur on treatment (19) based on evidence from clinical trials of drugs for osteoporosis in which there is a large reduction in risk for spine and hip fracture, although the reduction in risk of other fractures is lower. In practice, the occurrence of two or more fractures during treatment is a very rare event. Bone mineral density is commonly used as a tool to monitor treatment in the individual and an increase that exceeds the least significant change, for example, an increase in lumbar spine or total hip BMD more than 4% (19) may be considered a response. However, such changes occur over many months and persistence with medication declines very early in treatment (less than 50% after 12 months) (20) so an earlier response marker would be preferred. The International Osteoporosis Foundation has proposed that a BTM such as PINP or

CTX measured within 3 months of starting therapy would help identify poor adherence with the commonest osteoporosis therapy, oral bisphosphonates (21). Another advantage to using BTM rather than bone mineral density is that measurements are less expensive. In our hospital setting, a PINP measurement costs less than 20% that of a bone mineral density measurement. Finally, BTM may be a better surrogate for fracture risk reduction than BMD. The proportion of treatment effect explained by BTM has usually been higher than that for BMD (7).

Bisphosphonate

Bisphosphonates are the most commonly used drugs for osteoporosis. There are three oral bisphosphonates that are licensed in most countries, namely alendronate, risedronate and ibandronate. The absorption of the oral bisphosphonates is very poor and as the dosing regimen is complex, many patients do not comply fully with the instructions so do not achieve an optimal response even though they may take their medication regularly. The oral bisphosphonates have been compared in the TRIO study (Clinical Trial Number: NCT00666627) (22) to evaluate the clinical utility of BTM to assess response. Alendronate and ibandronate decreased BTM (CTX, NTX) more than risedronate. In this study, more than 80% of patients responded to treatment as defined by a decrease more than the LSC for CTX (56%) and PINP (38%) after 3 months of treatment. Response can also be defined as a reduction to a level below the mean found in healthy young women (22). In one study, the mean values were given as 217–317 ng/L for CTX and 32–38 µg/L for PINP (23). In the assessment of treatment response in the individual, the magnitude of the decrease has also been found to be important; for example, with alendronate (24) and risedronate (25) the greater the reduction in BTM, the greater the reduction in vertebral fracture risk.

Zoledronic acid is given by annual intravenous infusion, thus avoiding concerns about poor absorption. It results in a reduction in CTX by 2 weeks and when it is given for 6 years as in the Horizon Study, the suppression of CTX and PINP is maintained (26). PINP was found to be even better than CTX and BMD in the horizon study at identifying clinical (fracture) efficacy and responders (27). As with the oral treatments, the greater the reduction in PINP with zoledronic acid, the greater the reduction in the risk of vertebral fractures (28).

Denosumab

Denosumab inhibits bone resorption, leading to an early and large decrease in bone resorption markers followed by a later and smaller decrease in bone formation markers. Bone resorption markers (such as CTX) decrease within 24 h of treatment. In the FREEDOM Study, there was no overlap in CTX levels between treated and control subjects at one month indicating that everyone appears to respond (29). Denosumab results in a greater inhibition of bone resorption than zoledronic acid (30). PINP decreases over several months to a lesser extent than the bone resorption markers and remains suppressed with continued dosing for up to 8 years (31). Once the drug is stopped, the BTM overshoot so that their levels are increased compared to baseline (32). These high BTM results are associated with accelerated bone loss, and there are recent reports of multiple vertebral fractures associated with this high BTM (33).

SERMs

Selective receptor oestrogen agonists (SERM) such as raloxifene have a weaker effect on bone turnover than bisphosphonates and denosumab. Even so, their effect can be monitored using BTM. In 60–65% of women with osteopenia, a significant response could be demonstrated using the LSC approach with CTX or PINP (34). The BTM response to raloxifene was greatest in those with greatest adherence (35) providing further support for use of BTM as a means of identifying poor adherence to therapy (see below).

Teriparatide

Teriparatide is an anabolic agent administered as a daily subcutaneous injection and bone formation markers increase within days of starting treatment (36), peaking by 3 months. PINP has proven to be the most responsive BTM to this treatment. Most patients have a significant response using PINP and an increase greater than the LSC (of more than 10 µg/L) (37) may be used to identify responders. The change in BTM relates to the later change in BMD (38). Poor BMD response is associated with low BTM at baseline (PINP, NTX) or smaller increase in BTM after 4 months on treatment (39).

The licence for teriparatide is for 2 years as there is a concern about osteosarcoma with long-term use and the

effect of the drug wanes after three years of therapy. There is accelerated bone loss after stopping teriparatide, but this can be prevented by administering bisphosphonates, raloxifene or denosumab (40).

Abaloparatide is a new licensed anabolic therapy for osteoporosis (41). It is a synthetic peptide analogue of the human parathyroid hormone-related protein and works through the PTH receptor as does teriparatide. However, the increase in PINP is less than that with teriparatide (42). The clinical utility of BTMs for monitoring abaloparatide therapy has not yet been fully reported.

Practical approach to monitoring

We have been using BTM to monitor osteoporosis therapy in our secondary care practice for 20 years. We have observed that many patients commencing treatment and having a poor BTM response are identified as having minor errors in following the dosing instructions that may not be picked up by a brief medication review. This is particularly important as most osteoporosis medication prescribing takes place in general practice by non-specialists who may not appreciate the limited absorption of oral bisphosphonates and the need for complete and consistent adherence to the dose regime. In primary care, time and resource to undertake early assessment of compliance is also limited and so we felt it appropriate to roll out the approach of monitoring osteoporosis therapy using BTM into general practice.

Figure 1 illustrates the algorithm that has been implemented in clinical practice. The physician decides to treat; most commonly, this would be with an oral bisphosphonate such as alendronate. At this point, PINP is measured. Our local laboratory uses the automated immunoassay (Roche Cobas) for this measurement; the results are similar for other automated immunoassays (IDS iSYS) (23) (Table 2); we need more data on the Orion PINP assay in comparison to the other assays. In one study, the results were similar (43) but in another the Orion assay give results lower than either Roche or IDS (Cavalier, personal communication). A discussion is held with the patient after one month to assess compliance and any problems or concerns with their treatment. This discussion is often held over the phone and may be initiated by their doctor, nurse or a pharmacist. The PINP measurement is repeated after 6 months to assess response. Treatment response is defined as a decrease in PINP that exceeds the least significant change of 10 µg/L or a decrease to below the geometric mean for young women (35 µg/L). As we

Figure 1

Sheffield PINP monitoring algorithm for anti-resorptive treatment. Optimal treatment response with PINP is a decrease of 10 µg/L to below 35 µg/L. Optimal treatment response with CTX is a decrease of 100 ng/L to below 280 ng/L.

describe later, the management advice accompanying the algorithm highlights factors such as incident fracture that may affect the interpretation of the PINP result and the actions to take if a clear response is not identified.

CTX can be used in the same way as for PINP for monitoring in practice, and it has the advantage of the change being earlier than for PINP (22). The mean CTX value is 280 ng/L (23) and the least significant change value is about 100 ng/L (in Table 2 it is 60–80 ng/L depending on the method).

The estimates of least significant change (and the geometric means) are based on the assay and within-subject variability and so are a statistical approach and

can be used for the monitoring of any intervention on bone turnover.

The mean values for PINP and CTX for the population were based on values obtained from studies in Italy, UK, France, Belgium, USA, Saudi Arabia and Denmark (23); there doesn't seem to be major differences between countries (44) and so they should be suitable for international use. The mean values for PINP using Roche and IDS assays are probably similar (23) (Table 2). However, the mean values for CTX used to be higher for IDS (Table 2) (43) but more recent reports show them to be lower (23). There is a clear need for harmonisation of assay results for BTMs (7).

Treatment targets

The rationale for choosing a least significant change of 10 µg/L is that changes up to this level could occur by chance in up to 95% of people, whereas a change greater than this is relatively uncommon, and it is based on untreated postmenopausal women with low bone mineral density (45). The least significant change is also similar for both the Roche Cobas and the IDS iSYS assays as is the mean response to oral bisphosphonates (Table 2).

The rationale for choosing a target PINP value of ≤35 µg/L is that in clinical trials of anti-resorptive drugs, the lowest fracture risk is found in those women with bone turnover marker levels on treatment below the average value for young women (46, 47). Bone turnover markers have a skewed distribution, so it is best to take the geometric mean or the median. This value is around 35 µg/L for the Roche Cobas assay (23). The critical values for PINP (and CTX) are supported by results from those obtained in 50 women from the TRIO study with postmenopausal osteoporosis treated with oral bisphosphonate and compared to 200 healthy young control women. It can be seen from Table 2 that the mean values for young women are similar for PINP and close to 35 µg/L for both assays. It can also be seen that the least significant change is similar and as the baseline PINP was around 46 µg/L, an LSC of 23% is equivalent to about 10 µg/L. It is also notable that the mean reduction on treatment after 12–13 weeks is very similar. The attraction of using PINP rather than CTX can be seen from this table – the LSC is lower for PINP than CTX and PINP does not need to be taken in the fasting state.

The concepts of least significant change and target for treatment are not unique to the use of bone turnover markers in osteoporosis and so are already familiar to colleagues in primary care. In type 2 diabetes, it is usual to

Table 2 The critical values for PINP and CTX are supported by results in 50 women from the TRIO study with postmenopausal osteoporosis and treated with oral bisphosphonate and 200 healthy young control women.

	PINP		CTX	
	IDS iSYS	Roche Cobas e411	IDS iSYS	Roche Cobas e411
Controls, mean	31 µg/L	33 µg/L	327 ng/L	221 ng/L
LSC	29 %	23 %	54 %	50 %
LSC	6.2 µg/L	5.7 µg/L	80 ng/L	60 ng/L
Effect	–51 %	–54 %	–75 %	–74 %
Effect	–28 µg/L	–32 µg/L	–490 ng/L	–360 ng/L

The means in the control group are geometric means; LSC estimates were based on the CV of samples taken at 12 and 13 weeks on treatment and multiplied by 2.77; effects are change from baseline to the average of the 12 and 13 week values. The full study has been described before (20).

monitor with haemoglobin A_{1c} and consider the reference change value (the same as least significant change) and the target; the critical values used are 0.5% change and a target of 7.0% (48).

Sources of variability in BTMs

Many clinical factors influence BTM measurements, but we pay particular attention to the occurrence of a fracture or to treatment with glucocorticoids as these are common confounders and have a clinically important impact. There is a large increase in PINP after a fracture, with a mean increase of 55% six weeks after wrist fracture (49), 96% six weeks after ankle fracture (50) and 100% 12 weeks after tibial shaft fracture (51). BTMs have also been reported to be increased after vertebral fracture (52). The magnitude of the increase appears to relate to the bone size at the site of fracture and may be greater if the fracture is managed surgically. Many patients initiate osteoporosis therapy in the weeks following a fracture and in the GLOW study, 7% of patients taking osteoporosis medication for 3 years sustained an incident fracture, illustrating the importance of this effect (53).

Glucocorticoid therapy reduces the level of PINP in a dose-responsive manner, and this also makes the interpretation of PINP difficult – is any reduction due to a beneficial effect of bisphosphonate treatment or due to the harmful effect of the glucocorticoid? For example, a daily dose of 10mg prednisone resulted in a 20% reduction in PINP over a week. Thus, PINP is only helpful in monitoring treatment of glucocorticoid-induced osteoporosis if the glucocorticoid dose is established and remains stable. If accessible, it may be preferable to use another BTM in this clinical situation, such as bone ALP, CTX or NTX.

Evaluation of use in clinical practice

Antiresorptive therapy

We introduced the monitoring algorithm (Fig. 1) into primary care in Sheffield in September 2011 and conducted an audit on all patients being evaluated for osteoporosis at the Metabolic Bone Centre in Sheffield in July 2012. New treatment was recommended to the general practitioner in 108 cases (mean age 65 years, 86% female) and baseline PINP was obtained by the GP in 76 of these. Follow-up measurement was made in 34 of these. We found that at follow-up, 27 (79%) met the

Serum PINP, µg/L

Figure 2

Serum PINP (µg/L) measured using Roche Cobas in patients starting anti-resorptive therapy in 34 patients based in general practice setting. Solid lines indicate response with a significant change in PINP to a level below 35 µg/L, and dotted lines no response. The lines are coloured red to indicate those patients who had a significant change in PINP but did not reach the target, yellow to indicate those who reached the target but PINP did not change and green to show an increase in PINP. The second PINP value should have been taken after 6 months of treatment and the broken horizontal line shows the critical value of 35 µg/L (the mean value of PINP in healthy young women).

criteria for treatment response (Fig. 2). Among the 7 people with poor response, we found reasons for this in 3 cases (poor compliance, intercurrent surgery and the sample measured too early). We were encouraged by this early uptake of PINP monitoring and are working further to develop awareness among general practitioners and to develop confidence in its use.

We acknowledge that monitoring using PINP in clinical practice is by necessity pragmatic, needing to minimise cost and patient inconvenience and differs considerably from the research approach. Nonetheless, the approach has been welcomed by GP colleagues and we believe is preferable to no monitoring or reassessment of BMD after 2 years, by which time a high proportion of patients have stopped treatment.

The interpretation of PINP results and the need to change treatment needs to be considered on an individual basis using clinical judgement and considering factors

Review	R Eastell and others	Bone turnover markers	178:1	R26
--------	----------------------	-----------------------	-------	-----

such as the severity of the osteoporosis, likelihood of poor compliance (e.g. presence of dementia) and presence of known confounders.

In all patients with suboptimal response, especially if repeat PINP remains increased, change in treatment needs to be considered. We would often move onto parenteral treatment at this point to eliminate problems due to poor compliance and/or poor absorption.

Our initial evaluation of the use of PINP for monitoring anti-resorptive therapy in clinical practice (Fig. 2) drew our attention to several types of responses.

Black – a significant decrease to below 35 µg/L

This is consistent with a good response to a level that is associated with low fracture risk and so is the optimal response. We would recommend the physician confirms compliance, enquires about any drug side effects and encourages the patient to continue therapy and report any new issues. Medications should be reviewed at least annually and risk assessment including DXA planned at 5 years. Further PINP measurement is not considered necessary unless the clinical situation alters.

Red – a significant decrease but not to below 35 µg/L

Many patients like this will have high baseline BTM. The high PINP on treatment may indicate that fracture risk is still somewhat elevated (46, 47) and the patient may benefit from treatment with a parenteral drug such as zoledronic acid or denosumab.

- A high baseline PINP may be due to fracture within the preceding 6–12 months, especially if very recent. In that case, any subsequent decrease may be due to the healing fracture and so does not indicate good response. Thus, it is important to check compliance and if this is adequate then continue with treatment and repeat PINP in another 3 months. If PINP is still above 35 then consider investigation for secondary osteoporosis.
- A high baseline PINP may occur in a patient with another disease affecting PINP e.g. skin disease, liver dysfunction. Consider if an additional confounder has developed and if investigations are indicated; check compliance and repeat in another 3 months. If uncertainty remains at that point, consider change in treatment.

Yellow – no significant change and remains under 35 µg/L

- Low PINP values are associated with a lower risk of fracture. It is not certain whether patients with low BTM have fracture benefit from treatment – one study with alendronate would indicate that there is no benefit (16), whereas a study with risedronate indicates there may be some benefit (54). Certainly, patients with low PINP tend to have low rates of bone loss when left untreated and low BMD increases in response to anti-resorptive treatment (18). Nonetheless, continued treatment would generally be considered appropriate, and the patient may be advised that the BTM result indicates a low risk of fracture.

Green – a significant increase in PINP, to above 35 µg/L

- This is likely to reflect either an intercurrent event leading to an increase in bone turnover or presence of a disease that provides a non-bone source of PINP.
- A common cause of increased PINP is a new fracture (magnitude of increase related to size of bone affected) with an additional impact from surgical intervention. If a fracture is confirmed, then check the compliance, reassure the patient that a fracture occurring very early in treatment is not due to treatment failure and reassess after another 3–6 months.
- Other causes include a reduction in glucocorticoid dose since treatment initiation, i.e. with greater suppression of PINP at baseline when on a higher dose – if the 6-month result remains above 35 µg/L, this still suggests poor treatment response. In practice, we would measure another marker unaffected by glucocorticoid.
- At this point e.g. NTX and if result is low then presume response.
- May be due to new secondary osteoporosis e.g. development of thyrotoxicosis – clinical evaluation and relevant investigation should be undertaken and any underlying cause should be treated. Compliance with the osteoporosis treatment should also be checked and encouraged.
- We may observe fluctuations in PINP (and other BTM) in patients with co-morbidities especially those affecting liver, kidney and skin. For example, increases in PINP are observed after non-bone surgical intervention reflecting PINP from type I collagen in skin. In these patients, check compliance and consider using another marker.

If there remains a suboptimal response, then a change to an alternative agent such as a parenteral drug (zoledronic acid or denosumab) may be considered. If there is a good BTM response, but the patient has one or more new vertebral fractures, then a change to an anabolic drug (teriparatide) may be considered.

Monitoring offset of effect in the individual

It is often recommended that oral bisphosphonate therapy is stopped after 5 years in milder forms of osteoporosis (55). The rationale for this is that longer-term therapy may increase the risk of atypical femur fractures, and once treatment is stopped, there is continued benefit with little bone loss from the spine and continued (if mild) suppression of bone turnover (56). Attempts have been made to monitor the offset of effect with BMD, but the changes at the hip are quite small relative to the least significant change and so only a small proportion of patients are identified as having offset of effect, with just 29% having more than 5% bone loss from the total hip 5 years after stopping alendronate (57). BTM could be used for this purpose, perhaps using the LSC approach to examine for an increase or the threshold approach to identify a value that is above the mean for young women and so merits re-starting therapy. There has been little

published on this topic and there appears to be little association between change in BMD or BTM and fracture risk off treatment with alendronate (58). We await further research before making any recommendations.

Anabolic therapy

We also use PINP to assess response to teriparatide treatment (Fig. 3). Teriparatide is used in patients with severe and complicated osteoporosis, so it is important to consider if response is optimal as early as possible, particularly as

Figure 3

Sheffield PINP monitoring algorithm for anabolic treatment. An optimal response would be an increase in PINP of more than 10 µg/L to above 69 µg/L.

Figure 4

The absolute value of PINP (µg/L) measured using Roche Cobas at baseline, one and three months after starting teriparatide in 91 people for osteoporosis. The blue dashed horizontal line represents the upper limit of the reference interval for healthy young women (69 µg/L). Overall, 95% responded with an increase of more than 10 µg/L above baseline at both 1 and 3 months and 66% exceeded the upper limit of the reference interval in at least one timepoint. Patients meeting both response criteria, i.e. demonstrating an optimal response, are shown in black; those who had a significant increase in PINP but not exceeding the reference interval are shown in red. Patients in whom no BTM response was demonstrated are shown in green. The extremely high value of PINP of more than 500 µg/L was observed in a patient with autoimmune hepatitis.

Review	R Eastell and others	Bone turnover markers	178:1	R28
--------	----------------------	-----------------------	-------	-----

treatment is limited to 24 months. Suboptimal response may be due to issues with compliance, drug storage or injection technique.

We evaluated 91 patients monitored using PINP. All had previously been treated with anti-resorptive therapy, mean age 71 years (89% female). The baseline PINP was 35 µg/L using the Roche Cobas automated immunoassay analyser, reflecting the effect of the prior anti-resorptive treatment. We took our treatment targets as an increase of more than the least significant change in PINP at months one and three (10 µg/L) (45) and an increase to above the reference interval of 69 µg/L (59) on at least one occasion. We found that by 3 months of treatment 93% exceeded the least significant change and 66% exceeded the upper limit of the reference interval (Fig. 4). This responder rate of 93% was similar to that found in clinical trials of teriparatide of 87, 77 and 87% (60, 61, 62). There was no significant correlation between PINP results and change in lumbar spine BMD at two years, but this was difficult to evaluate as only 29% of our patients had reliable spine scans due to very high prevalence of vertebral fracture and degenerative change. The baseline PINPs in this evaluation were low as all patients had previously been treated with anti-resorptive drugs and so these findings are not relevant to patients starting teriparatide with no such prior therapy.

In our experience, treatment is often commenced without measurement of a baseline PINP, especially in primary care. In this situation, it is particularly important to undertake a thorough evaluation of adherence to treatment, and we find it remains valuable to make the 6-month measurement. A PINP value on treatment that is low or low normal for anti-resorptive treatment (i.e. <35 µg/L) or above the reference interval for anabolic therapy (i.e. >69 µg/L) may be presumed to indicate adequate response. However, this approach is less well documented than the least significant change approach.

Current recommendations: examples

The IOF and IFCC made recommendations concerning BTM and reviewed national guidelines; five out of nine national societies or organisations recommended the use of BTM for treatment monitoring, although recognising that further research and evaluation remains necessary (7).

The IOF proposed that a PINP or CTX value at 12 weeks on treatment with oral bisphosphonate can identify poor response and be used to identify patients who are unlikely

to be adhering to therapy (21) or who have failed therapy (53). The IOF proposed using the BTMs at 12 weeks rather than 3 months on treatment and the responder rate in the TRIO study was similar for 12 and 48 weeks (22), and so this approach is appropriate. National guidelines supporting the use of bone turnover markers are also available, such as those from the Austrian Group and from the Japanese Osteoporosis Society (63, 64).

Declaration of interest and Funding

R Eastell reports grants and personal fees from Amgen, grants from AstraZeneca, grants and personal fees from Alexion, grants, personal fees and non-financial support from Immunodiagnostic Systems, grants, personal fees and non-financial support from Roche, personal fees from Lilly, personal fees from D3 Biomedical Science Institutes, personal fees from GSK Nutrition, within 3 years of writing this review. K E Naylor, T Pigott and F Gossiel have no disclosures. J S Walsh has received speaker's honoraria from Lilly and the donation of drug and placebo from Prostrakan. N F A Peel has received speaker's honoraria and funding to attend educational events from Warner-Chilcott, Lilly, Amgen, GSK and Prostrakan and consultancy fees from Internis Pharma and Lilly.

Acknowledgements

The authors thank Roy Wheeler for his contribution to the service evaluation of anti-resorptive therapies, Roche Diagnostics and Immunodiagnostics Systems for research support.

References

- Eastell R, O'Neill TW, Hofbauer LC, Langdahl B, Reid IR, Gold DT & Cummings SR. Postmenopausal osteoporosis. *Nature Reviews Disease Primers* 2016 **2** 16069. (<https://doi.org/10.1038/nrdp.2016.69>)
- Eastell R, Delmas PD, Hodgson SF, Eriksen EF, Mann KG & Riggs BL. Bone formation rate in older normal women: concurrent assessment with bone histomorphometry, calcium kinetics, and biochemical markers. *Journal of Clinical Endocrinology and Metabolism* 1988 **67** 741–748. (<https://doi.org/10.1210/jcem-67-4-741>)
- Eastell R, Colwell A, Hampton L & Reeve J. Biochemical markers of bone resorption compared with estimates of bone resorption from radiotracer kinetic studies in osteoporosis. *Journal of Bone and Mineral Research* 1997 **12** 59–65. (<https://doi.org/10.1359/jbmr.1997.12.1.59>)
- Chavassieux P, Portero-Muzy N, Roux JP, Garnero P & Chapurlat R. Are biochemical markers of bone turnover representative of bone histomorphometry in 370 postmenopausal women? *Journal of Clinical Endocrinology and Metabolism* 2015 **100** 4662–4668. (<https://doi.org/10.1210/jc.2015-2957>)
- Szulc P, Naylor K, Hoyle NR, Eastell R, Leary ET & National Bone Health Alliance Bone Turnover Marker Project. Use of CTX-I and PINP as bone turnover markers: National Bone Health Alliance recommendations to standardize sample handling and patient preparation to reduce pre-analytical variability. *Osteoporosis International* 2017 **28** 2541–2556. (<https://doi.org/10.1007/s00198-017-4082-4>)
- Okabe R, Nakatsuka K, Inaba M, Miki T, Naka H, Masaki H, Moriguchi A & Nishizawa Y. Clinical evaluation of the Elecsys beta-CrossLaps serum assay, a new assay for degradation products of type I collagen C-telopeptides. *Clinical Chemistry* 2001 **47** 1410–1414.
- Vasikaran S, Eastell R, Bruyere O, Foldes AJ, Garnero P, Griesmacher A, McClung M, Morris HA, Silverman S, Trenti T *et al.* Markers of bone turnover for the prediction of fracture risk and

- monitoring of osteoporosis treatment: a need for international reference standards. *Osteoporosis International* 2011 **22** 391–420. (<https://doi.org/10.1007/s00198-010-1501-1>)
- 8 Fraser CG. Changes in serial results. In *Biological Variation: From Principles to Practice*, pp. 67–90. Ed CG Fraser. USA: AACC Press, 2001.
 - 9 Shieh A, Ishii S, Greendale GA, Cauley JA, Lo JC & Karlamangla AS. Urinary N-telopeptide and rate of bone loss over the menopause transition and early postmenopause. *Journal of Bone and Mineral Research* 2016 **31** 2057–2064. (<https://doi.org/10.1002/jbmr.2889>)
 - 10 Marques EA, Gudnason V, Lang T, Sigurdsson G, Sigurdsson S, Aspelund T, Siggeirsdottir K, Launer L, Eiriksdottir G & Harris TB. Association of bone turnover markers with volumetric bone loss, periosteal apposition, and fracture risk in older men and women: the AGES-Reykjavik Longitudinal Study. *Osteoporosis International* 2016 **27** 3485–3494. (<https://doi.org/10.1007/s00198-016-3675-7>)
 - 11 Ivaska KK, Lenora J, Gerdhem P, Akesson K, Vaananen HK & Obrant KJ. Serial assessment of serum bone metabolism markers identifies women with the highest rate of bone loss and osteoporosis risk. *Journal of Clinical Endocrinology and Metabolism* 2008 **93** 2622–2632. (<https://doi.org/10.1210/jc.2007-1508>)
 - 12 Rogers A, Hannon R & Eastell R. Biochemical markers as predictors of rates of bone loss after menopause. *Journal of Bone and Mineral Research* 2000 **15** 1398–1404. (<https://doi.org/10.1359/jbmr.2000.15.7.1398>)
 - 13 Vilaca T, Gossiel F & Eastell R. Bone turnover markers: use in fracture prediction. *Journal of Clinical Densitometry* 2017 **20** 346–352. (<https://doi.org/10.1016/j.jocd.2017.06.020>)
 - 14 Johansson H, Oden A, Kanis JA, McCloskey EV, Morris HA, Cooper C, Vasikaran S & IFCC-IOF Joint Working Group on Standardisation of Biochemical Markers of Bone Turnover. A meta-analysis of reference markers of bone turnover for prediction of fracture. *Calcified Tissue International* 2014 **94** 560–567. (<https://doi.org/10.1007/s00223-014-9842-y>)
 - 15 McCloskey EV, Vasikaran S, Cooper C & FRAX(R) Position Development Conference Members. Official positions for FRAX(R) clinical regarding biochemical markers from Joint Official Positions Development Conference of the International Society for Clinical Densitometry and International Osteoporosis Foundation on FRAX(R). *Journal of Clinical Densitometry* 2011 **14** 220–222. (<https://doi.org/10.1016/j.jocd.2011.05.008>)
 - 16 Bauer DC, Garnero P, Hochberg MC, Santora A, Delmas P, Ewing SK & Black DM. Pretreatment levels of bone turnover and the antifracture efficacy of alendronate: the fracture intervention trial. *Journal of Bone and Mineral Research* 2006 **21** 292–299. (<https://doi.org/10.1359/JBMR.051018>)
 - 17 Delmas PD, Licata AA, Reginster JY, Crans GG, Chen P, Misurski DA, Wagman RB & Mitlak BH. Fracture risk reduction during treatment with teriparatide is independent of pretreatment bone turnover. *Bone* 2006 **39** 237–243. (<https://doi.org/10.1016/j.bone.2006.02.003>)
 - 18 Eastell R, Boonen S, Cosman F, Reid IR, Palermo L, Cummings SR & Black DM. Relationship between pretreatment rate of bone loss and bone density response to once-yearly ZOL: HORIZON-PFT extension study. *Journal of Bone and Mineral Research* 2015 **30** 570–574. (<https://doi.org/10.1002/jbmr.2361>)
 - 19 Diez-Perez A, Adachi JD, Agnusdei D, Bilezikian JP, Compston JE, Cummings SR, Eastell R, Eriksen EF, Gonzalez-Macias J, Liberman UA *et al.* Treatment failure in osteoporosis. *Osteoporosis International* 2012 **23** 2769–2774. (<https://doi.org/10.1007/s00198-012-2093-8>)
 - 20 Netelenbos JC, Geusens JP, Ypma G & Buijs SJ. Adherence and profile of non-persistence in patients treated for osteoporosis – a large-scale, long-term retrospective study in The Netherlands. *Osteoporosis International* 2011 **22** 1537–1546. (<https://doi.org/10.1007/s00198-010-1372-5>)
 - 21 Diez-Perez A, Naylor KE, Abrahamsen B, Agnusdei D, Brandi ML, Cooper C, Dennison E, Eriksen EF, Gold DT, Guanabens N *et al.* International Osteoporosis Foundation and European Calcified Tissue Society Working Group. Recommendations for the screening of adherence to oral bisphosphonates. *Osteoporosis International* 2017 **28** 767–774. (<https://doi.org/10.1007/s00198-017-3906-6>)
 - 22 Naylor KE, Jacques RM, Paggiosi M, Gossiel F, Peel NF, McCloskey EV, Walsh JS & Eastell R. Response of bone turnover markers to three oral bisphosphonate therapies in postmenopausal osteoporosis: the TRIO study. *Osteoporosis International* 2015 **27** 21–31. (<https://doi.org/10.1007/s00198-015-3145-7>)
 - 23 Morris HA, Eastell R, Jorgensen NR, Cavalier E, Vasikaran S, Chubb SA, Kanis JA, Cooper C, Makris K & IFCC-IOF Working Group for Standardisation of Bone Marker Assays (WG-BMA). Clinical usefulness of bone turnover marker concentrations in osteoporosis. *Clinica Chimica Acta* 2016 **467** 34–41. (<https://doi.org/10.1016/j.cca.2016.06.036>)
 - 24 Bauer DC, Black DM, Garnero P, Hochberg M, Ott S, Orloff J, Thompson DE, Ewing SK & Delmas PD. Change in bone turnover and hip, non-spine, and vertebral fracture in alendronate-treated women: the fracture intervention trial. *Journal of Bone and Mineral Research* 2004 **19** 1250–1258. (<https://doi.org/10.1359/JBMR.040512>)
 - 25 Eastell R, Barton I, Hannon R, Chines A, Garnero P & Delmas P. Relationship of early changes in bone resorption to the reduction in fracture risk with risedronate. *Journal of Bone and Mineral Research* 2003 **18** 1051–1056. (<https://doi.org/10.1359/jbmr.2003.18.6.1051>)
 - 26 Black DM, Reid IR, Cauley JA, Cosman F, Leung PC, Lakatos P, Lippuner K, Cummings SR, Hue TF, Mukhopadhyay A *et al.* The effect of 6 versus 9 years of zoledronic acid treatment in osteoporosis: a randomized second extension to the HORIZON-Pivotal Fracture Trial (PFT). *Journal of Bone and Mineral Research* 2015 **30** 934–944. (<https://doi.org/10.1002/jbmr.2442>)
 - 27 Bell KJ, Hayen A, Glasziou P, Irwig L, Eastell R, Harrison SL, Black DM & Bauer DC. Potential usefulness of BMD and bone turnover monitoring of zoledronic acid therapy among women with osteoporosis: secondary analysis of randomized controlled trial data. *Journal of Bone and Mineral Research* 2016 **31** 1767–1773. (<https://doi.org/10.1002/jbmr.2847>)
 - 28 Jacques RM, Boonen S, Cosman F, Reid IR, Bauer DC, Black DM & Eastell R. Relationship of changes in total hip bone mineral density to vertebral and nonvertebral fracture risk in women with postmenopausal osteoporosis treated with once-yearly zoledronic acid 5 mg: the HORIZON-Pivotal Fracture Trial (PFT). *Journal of Bone and Mineral Research* 2012 **27** 1627–1634. (<https://doi.org/10.1002/jbmr.1644>)
 - 29 Eastell R, Christiansen C, Grauer A, Kutilek S, Libanati C, McClung MR, Reid IR, Resch H, Siris E, Uebelhart D *et al.* Effects of denosumab on bone turnover markers in postmenopausal osteoporosis. *Journal of Bone and Mineral Research* 2011 **26** 530–537. (<https://doi.org/10.1002/jbmr.251>)
 - 30 Miller PD, Pannacciulli N, Brown JP, Czerwinski E, Nedergaard BS, Bolognese MA, Malouf J, Bone HG, Reginster JY, Singer A *et al.* Denosumab or zoledronic acid in postmenopausal women with osteoporosis previously treated with oral bisphosphonates. *Journal of Clinical Endocrinology and Metabolism* 2016 **101** 3163–3170. (<https://doi.org/10.1210/jc.2016-1801>)
 - 31 Papapoulos S, Lippuner K, Roux C, Lin CJ, Kendler DL, Lewiecki EM, Brandi ML, Czerwinski E, Franek E, Lakatos P *et al.* The effect of 8 or 5 years of denosumab treatment in postmenopausal women with osteoporosis: results from the FREEDOM Extension study. *Osteoporosis International* 2015 **26** 2773–2783. (<https://doi.org/10.1007/s00198-015-3234-7>)
 - 32 Asociación Mexicana de Metabolismo Oseo y Mineral (AMMOM). Mexican consensus on osteoporosis. Mexican Bone and Mineral Metabolism Association. *Revista De Investigacion Clinica* 2001 **53** 469–495.
 - 33 Lamy O, Gonzalez-Rodriguez E, Stoll D, Hans D & Aubry-Rozier B. Severe rebound-associated vertebral fractures after denosumab

Review	R Eastell and others	Bone turnover markers	178:1	R30
--------	----------------------	-----------------------	-------	-----

- discontinuation: 9 clinical cases report. *Journal of Clinical Endocrinology and Metabolism* 2017 **102** 354–358. (<https://doi.org/10.1210/jc.2016-3170>)
- 34 Naylor KE, Jacques RM, Peel NF, Gossiel F & Eastell R. Response of bone turnover markers to raloxifene treatment in postmenopausal women with osteopenia. *Osteoporosis International* 2016 **27** 2585–2592. (<https://doi.org/10.1007/s00198-016-3573-z>)
 - 35 Finigan J, Naylor K, Paggiosi MA, Peel NF & Eastell R. Adherence to raloxifene therapy: assessment methods and relationship with efficacy. *Osteoporosis International* 2013 **24** 2879–2886. (<https://doi.org/10.1007/s00198-013-2386-6>)
 - 36 Glover SJ, Eastell R, McCloskey EV, Rogers A, Garnero P, Lowery J, Belleli R, Wright TM & John MR. Rapid and robust response of biochemical markers of bone formation to teriparatide therapy. *Bone* 2009 **45** 1053–1058. (<https://doi.org/10.1016/j.bone.2009.07.091>)
 - 37 Kregg JH, Lane NE, Harris JM & Miller PD. PINP as a biological response marker during teriparatide treatment for osteoporosis. *Osteoporosis International* 2014 **25** 2159–2171. (<https://doi.org/10.1007/s00198-014-2646-0>)
 - 38 Niimi R, Kono T, Nishihara A, Hasegawa M, Matsumine A, Nakamura T, Kono T & Sudo A. An algorithm using the early changes in PINP to predict the future BMD response for patients treated with daily teriparatide. *Osteoporosis International* 2014 **25** 377–384. (<https://doi.org/10.1007/s00198-013-2426-2>)
 - 39 Niimi R, Kono T, Nishihara A, Hasegawa M, Kono T & Sudo A. A retrospective analysis of nonresponse to daily teriparatide treatment. *Osteoporosis International* 2016 **27** 2845–2853. (<https://doi.org/10.1007/s00198-016-3581-z>)
 - 40 Ebina K, Hashimoto J, Kashii M, Hirao M, Kaneshiro S, Noguchi T, Tsukamoto Y & Yoshikawa H. The effects of switching daily teriparatide to oral bisphosphonates or denosumab in patients with primary osteoporosis. *Journal of Bone and Mineral Metabolism* 2016 **35** 91–98. (<https://doi.org/10.1007/s00774-015-0731-x>)
 - 41 Shirley M. Abaloparatide: first global approval. *Drugs* 2017 **77** 1363–1368. (<https://doi.org/10.1007/s40265-017-0780-7>)
 - 42 Miller PD, Hattersley G, Riis BJ, Williams GC, Lau E, Russo LA, Alexandersen P, Zerbini CA, Hu MY, Harris AG *et al.* Effect of abaloparatide vs placebo on new vertebral fractures in postmenopausal women with osteoporosis: a randomized clinical trial. *JAMA* 2016 **316** 722–733. (<https://doi.org/10.1001/jama.2016.11136>)
 - 43 Eastell R, Garnero P, Audebert C & Cahall DL. Reference intervals of bone turnover markers in healthy premenopausal women: results from a cross-sectional European study. *Bone* 2012 **50** 1141–1147. (<https://doi.org/10.1016/j.bone.2012.02.003>)
 - 44 Glover SJ, Gall M, Schoenborn-Kellenberger O, Wagener M, Garnero P, Boonen S, Cauley JA, Black DM, Delmas PD & Eastell R. Establishing a reference interval for bone turnover markers in 637 healthy, young, premenopausal women from the United Kingdom, France, Belgium, and the United States. *Journal of Bone and Mineral Research* 2009 **24** 389–397. (<https://doi.org/10.1359/jbmr.080703>)
 - 45 Eastell R, Kregg J, Chen P, Glass E & Reginster J. Development of an algorithm for using PINP to monitor treatment of patients with teriparatide. *Current Medical Research and Opinion* 2006 **22** 61–66. (<https://doi.org/10.1185/030079905X75096>)
 - 46 Eastell R, Hannon RA, Garnero P, Campbell MJ & Delmas PD. Relationship of early changes in bone resorption to the reduction in fracture risk with risendronate: review of statistical analysis – response. *Journal of Bone and Mineral Research* 2007 **22** 1656–1660. (<https://doi.org/10.1359/jbmr.07090b>)
 - 47 Delmas PD, Munoz F, Black DM, Cosman F, Boonen S, Watts NB, Kendler D, Eriksen EF, Mesenbrink PG, Eastell R *et al.* Effects of yearly zoledronic acid 5 mg on bone turnover markers and relation of PINP with fracture reduction in postmenopausal women with osteoporosis. *Journal of Bone and Mineral Research* 2009 **24** 1544–1551. (<https://doi.org/10.1359/jbmr.090310>)
 - 48 Little RR, Rohlfing CL, Sacks DB & National Glycohemoglobin Standardization Program Steering Committee. Status of hemoglobin A1c measurement and goals for improvement: from chaos to order for improving diabetes care. *Clinical Chemistry* 2011 **57** 205–214. (<https://doi.org/10.1373/clinchem.2010.148841>)
 - 49 Ingle B, Hay S, Bottjer H & Eastell R. Changes in bone mass and bone turnover following distal forearm fracture. *Osteoporosis International* 1999 **10** 399–407. (<https://doi.org/10.1007/s001980050246>)
 - 50 Ingle BM, Hay SM, Bottjer HM, Eastell R. Changes in bone mass and bone turnover following ankle fracture. *Osteoporosis International* 1999 **10** 408–416. (<https://doi.org/10.1007/s001980050247>)
 - 51 Veitch S, Findlay S, Hamer A, Blumsohn A, Eastell R & Ingle B. Changes in bone mass and bone turnover following tibial shaft fracture. *Osteoporosis International* 2006 **17** 364–372. (<https://doi.org/10.1007/s00198-005-2025-y>)
 - 52 Hashidate H, Kamimura M, Nakagawa H, Takahara K, Ikegami S, Uchiyama S & Kato H. Early changes in bone specific turnover markers during the healing process after vertebral fracture. *Open Orthopaedics Journal* 2011 **5** 32–36. (<https://doi.org/10.2174/1874325001105010032>)
 - 53 Diez-Perez A, Adachi JD, Adami S, Anderson FA Jr, Boonen S, Chapurlat R, Compston JE, Cooper C, Gehlbach SH, Greenspan SL *et al.* Risk factors for treatment failure with antiosteoporosis medication: the global longitudinal study of osteoporosis in women (GLOW). *Journal of Bone and Mineral Research* 2014 **29** 260–267. (<https://doi.org/10.1002/jbmr.2023>)
 - 54 Seibel MJ, Naganathan V, Barton I & Grauer A. Relationship between pretreatment bone resorption and vertebral fracture incidence in postmenopausal osteoporotic women treated with risendronate. *Journal of Bone and Mineral Research* 2004 **19** 323–329. (<https://doi.org/10.1359/JBMR.0301231>)
 - 55 Adler RA, El-Hajj Fuleihan G, Bauer DC, Camacho PM, Clarke BL, Clines GA, Compston JE, Drake MT, Edwards BJ, Favus MJ *et al.* Managing osteoporosis in patients on long-term bisphosphonate treatment: report of a task force of the American Society for Bone and Mineral Research. *Journal of Bone and Mineral Research* 2016 **31** 16–35. (<https://doi.org/10.1002/jbmr.2708>)
 - 56 Black DM, Schwartz AV, Ensrud KE, Cauley JA, Levis S, Quandt SA, Satterfield S, Wallace RB, Bauer DC, Palermo L *et al.* Effects of continuing or stopping alendronate after 5 years of treatment: the Fracture Intervention Trial Long-term Extension (FLEX): a randomized trial. *JAMA* 2006 **296** 2927–2938. (<https://doi.org/10.1001/jama.296.24.2927>)
 - 57 McNabb BL, Vittinghoff E, Schwartz AV, Eastell R, Bauer DC, Ensrud K, Rosenberg E, Santora A, Barrett-Connor E & Black DM. BMD changes and predictors of increased bone loss in postmenopausal women after a 5-year course of alendronate. *Journal of Bone and Mineral Research* 2013 **28** 1319–1327. (<https://doi.org/10.1002/jbmr.1864>)
 - 58 Bauer DC, Schwartz A, Palermo L, Cauley J, Hochberg M, Santora A, Cummings SR & Black DM. Fracture prediction after discontinuation of 4 to 5 years of alendronate therapy: the FLEX study. *JAMA Internal Medicine* 2014 **174** 1126–1134. (<https://doi.org/10.1001/jamainternmed.2014.1232>)
 - 59 Glover SJ, Gamero P, Naylor K, Rogers A & Eastell R. Establishing a reference range for bone turnover markers in young, healthy women. *Bone* 2008 **42** 623–630. (<https://doi.org/10.1016/j.bone.2007.12.218>)
 - 60 McClung MR, San Martin J, Miller PD, Civitelli R, Bandeira F, Omizo M, Donley DW, Dalsky GP & Eriksen EF. Opposite bone remodeling effects of teriparatide and alendronate in increasing bone mass. *Archives of Internal Medicine* 2005 **165** 1762–1768. (<https://doi.org/10.1001/archinte.165.15.1762>)
 - 61 Neer RM, Arnaud CD, Zanchetta JR, Prince R, Gaich GA, Reginster JY, Hodman AB, Eriksen EF, Ish-Shalom S, Genant HK *et al.* Effect of parathyroid hormone (1–34) on fractures and bone mineral density in postmenopausal women with osteoporosis. *New England*

- Journal of Medicine* 2001 **344** 1434–1441. (<https://doi.org/10.1056/NEJM200105103441904>)
- 62 Tsujimoto M, Chen P, Miyauchi A, Sowa H & Kregge JH. PINP as an aid for monitoring patients treated with teriparatide. *Bone* 2011 **48** 798–803. (<https://doi.org/10.1016/j.bone.2010.12.006>)
- 63 Bieglmayer C, Dimai HP, Gasser RW, Kudlacek S, Obermayer-Pietsch B, Woloszczuk W, Zwettler E & Griesmacher A. Biomarkers of bone turnover in diagnosis and therapy of osteoporosis:

a consensus advice from an Austrian working group. *Wiener Medizinische Wochenschrift* 2012 **162** 464–477. (<https://doi.org/10.1007/s10354-012-0133-9>)

- 64 Nishizawa Y, Ohta H, Miura M, Inaba M, Ichimura S, Shiraki M, Takada J, Chaki O, Hagino H, Fujiwara S *et al.* Guidelines for the use of bone metabolic markers in the diagnosis and treatment of osteoporosis (2012 edition). *Journal of Bone and Mineral Metabolism* 2013 **31** 1–15. (<https://doi.org/10.1007/s00774-012-0392-y>)

Received 14 July 2017

Revised version received 11 October 2017

Accepted 17 October 2017