Primary adrenal insufficiency is associated with impaired natural killer cell function: a potential link to increased mortality

Irina Bancos^{1,2,*}, Jon Hazeldine^{3,4,*}, Vasileios Chortis^{1,5}, Peter Hampson^{3,4}, Angela E Taylor^{1,5}, Janet M Lord^{3,4} and Wiebke Arlt^{1,5}

¹Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK, ²Division of Endocrinology, Diabetes, Metabolism, and Nutrition, Department of Internal Medicine, Mayo Clinic, Rochester, Minnesota, USA, ³Institute of Inflammation and Ageing, ⁴Medical Research Council-Arthritis Research UK (MRC-ARUK), Centre for Musculoskeletal Ageing Research, University of Birmingham, Birmingham, UK, and ⁵Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, UK

*(I Bancos and J Hazeldine contributed equally to this work)

Correspondence should be addressed to W Arlt
Email
w.arlt@bham.ac.uk

The journal and the authors apologise for an error in Table 1 of this article that was published in the April 2017 issue (volume: 176, issue: 4, pages: 471–480) of *the European Journal of Endocrinology*. The unit for fludrocortisone should read as micrograms (µg) and not as reported.