Androgen excess and diagnostic steroid biomarkers for nonclassic 21-hydroxylase deficiency without cosyntropin stimulation

Adina F Turcu¹, Diala El-Maouche², Lili Zhao³, Aya T Nanba¹, Alison Gaynor², Padma Veeraraghavan², Richard J Auchus^{1,4} and Deborah P Merke^{2,5}

¹Division of Metabolism, Endocrinology and Diabetes, University of Michigan, Ann Arbor, Michigan, USA, ²National Institutes of Health (NIH) Clinical Center, Bethesda, Maryland, USA, ³School of Public Health, ⁴Department of Pharmacology, University of Michigan, Ann Arbor, Michigan, USA, and ⁵Eunice Kennedy Shriver National Institute of Child Health and Human Development, Bethesda, Maryland, USA

Correspondence should be addressed to A F Turcu
Email
aturcu@umich.edu

The authors apologise for an error in Table 2 of the above article published in the July 2020 issue (vol 183 Iss 1 pp 63–71) of the *European Journal of Endocrinology*. The values for 17OHP4 should be as given in the table below and not as published. The correct Table 2 is published in full below.

Table 2 Comparison of steroid concentrations between patients with nonclassic 21-hydroxylase deficiency (N21OHD) and controls. Data are expressed as median (interquartile range).

Steroid (nmol/L)	N21OHD (n = 32)	Controls (n = 54)	Ratio of medians N21OHD/controls	P
17OHP4	18.1 (8.7–33.0)	2.8 (0.9–4.3)	6.5	<0.0001
16OHP4	2.4 (1.4–4.9)	0.6 (0.4–0.9)	4.1	<0.0001
21dF	1.8 (1.3–5.3)	0.0 (0.0–0.3)	N/A	<0.0001
Progesterone	0.8 (0.5–1.4)	0.4 (0.3–0.8)	1.8	0.0018
Cortisol	202.5 (125.9–442.8)	249.1 (208.9–373.7)	0.8	0.197
Cortisone	53.7 (30.5–72.3)	51.5 (40.3–60.8)	1.0	0.855
Corticosterone	1.6 (0.8–3.5)	5.9 (3.5–13.6)	0.3	<0.0001
11dF	0.8 (0.5–1.4)	0.9 (0.4–1.6)	0.8	0.626
DOC	0.3 (0.3–0.4)	0.5 (0.4–0.9)	0.6	0.113
A4	4.3 (2.0–8.1)	3.9 (2.2–8.5)	1.1	0.798
Testosterone	1.1 (0.5–2.7)	1.6 (0.8–4.3)	0.7	0.195
11OHA4	9.6 (5.1–25.9)	4.5 (2.9–7.1)	2.1	<0.0001
11KA4	1.1 (0.7–1.7)	0.6 (0.4–0.8)	1.7	0.0001
11OHT	1.0 (0.3–1.7)	0.4 (0.3–0.6)	2.2	0.0019
11KT	1.8 (1.2–3.3)	0.9 (0.6–1.3)	2.0	<0.0001
11KT/Testosterone	1.8 (1.2–3.0)	0.6 (0.2–0.9)	3.3	<0.0001
11OHA4/A4	2.6 (2.1–4.6)	1.2 (0.7–2.1)	2.0	<0.0001
A4/testosterone	3.8 (2.8–4.5)	3.5 (2.4–4.2)	1.3	0.274

A4, androstenedione; DOC, 11-deoxycorticosterone; 17OHP4, 17 α -hydroxyprogesterone; 16OHP4, 16 α -hydroxyprogesterone; 21dF, 21-deoxycortisol; 11dF, 11-deoxycortisol; 11OHA4, 11 β -hydroxyandrostenedione; 11KA4, 11-ketoandrostenedione; 11OHT 11 β -hydroxytestosterone; 11KT, 11-ketotestosterone.